

Original Research Article

COMPARATIVE PHYTOCHEMICAL, PROXIMATE, AND *in vitro* ANTIHYPERGLYCEMIC ANALYSES OF THE STEM BARK EXTRACTS OF *Kigelia africana* (Lam.) Benth. AND *Annona senegalensis* Pers.

Abstract

The present research aimed to investigate the phytochemical bioactive constituents and proximate composition of the methanol extract of *Kigelia africana* and *Annona senegalensis* stem bark, as well as its *in vitro* antidiabetic potential. The results suggest that the presence of bioactive compounds may be responsible for the numerous medicinal capacities of these plants, including their antidiabetic abilities. The methanol stem bark extracts of *K. africana* and *A. senegalensis* demonstrated strong α -amylase and yeast cell glucose absorption inhibition, respectively. The stem bark methanol extract of *A. senegalensis* exhibited the highest level of inhibition (87.61%) at 0.1 mg/ml concentration when compared to conventional acarbose medication (56.0%) on the α -amylase enzyme. The percentage of glucose uptake by both methanol stem bark extract of *Kigelia africana* and *Annona senegalensis* was recorded maximum (31.40%) at a concentration of 0.5 mg/ml. The present research establishes the antidiabetic effect of methanol stem bark extract of *Kigelia africana* and *Annona senegalensis* by *in vitro* assays.

Keywords: *Kigelia africana*; *Annona senegalensis*; Phytochemicals; Antihyperglycemic; Stem Bark; Alpha-amylase inhibition; Yeast Cell Glucose Uptake.

Introduction

Diabetes mellitus is a complex chronic metabolic disorder that affects carbohydrate, fat, and protein metabolism. It is a non-infectious illness that affects 5% of the worldwide population [1-4] and accounts for nearly 10% of total healthcare costs in several countries [5]. It is caused by the pancreas producing insufficient or inadequate insulin, resulting in an increase or decrease in glucose concentrations. Many biological systems, including blood vessels and nerves, have been shown to be harmed by it [6]. One of the treatment techniques is to minimize postprandial hyperglycemia, by retarding the absorption of glucose by inhibiting carbohydrate-hydrolyzing enzymes, such as α -amylase and α -glucosidase [7, 8]. Herbal medications have recently been used in the treatment of diabetes since they have fewer side effects than synthetic hyperglycemic agents; therefore, there is a need to create strong, safe, and cost-effective drugs for diabetes management. These effective, safe, and inexpensive drugs might be developed by using

medicinal plants that have been used by humans to treat or cure disorders such as diabetes [9, 10, 11]. There is available literature that shows more than 400 plant species with hypoglycemic activity [12, 13] however, only a few of these medicinal plants have received scientific scrutiny, even though the World Health Organization has recommended that medical and scientific examinations of such plants should be conducted [14]. For this study, we will be focused on two plant species; *Kigelia africana* (Lam.) Benth and *Annona senegalensis* Pers.

Kigelia africana (Lam.) Benth (also known as the sausage tree) is one of the tree species that has been extensively used for its medicinal properties [15]. It is the only species in the genus *Kigelia* and belongs to the family Bignoniaceae. It is also known to have several local names such as Rawuya (Hausa), Pandoro (Yoruba), Uturubein (Igbo), Mranaa (Swahili), Worsboom (Afrikaans) etc. This plant has been used to treat a variety of ailments, including skin diseases like eczema, fungal infections, psoriasis, and boils, as well as more serious diseases like leprosy, impetigo, syphilis, skin cancer, elephantiasis, dysentery, swellings, constipation, piles, malaria, diabetes, pneumonia, worm infestations, venereal diseases, rheumatism, psoriasis [16, 17, 18, 19, 20, 21, 22]. Several chemical constituents have been isolated from various parts of *Kigelia africana* by several researchers. Secondary metabolites such as alkaloids, steroids, saponins, tannins, flavonoids, etc. [23] have been identified in the plant. Also, phenylpropanoid, stigmasterol, β -sitosterol, ferulic acid, lapachol, iridoids, kigelinole, kigelin, pinnatal, isopinnatal, verminoside, and other chemical compounds have been found in *K. africana* [24, 25, 26, 27].

Annona senegalensis Pers. commonly known as wild custard apple, is a member of the Annonaceae family and is widely distributed throughout Nigeria and is known locally as Gwándàndàj (Hausa), Dukuu-hi (Fulani), Uburuocha (Igbo), Arere or Abo (Yoruba) [28]. Numerous ethnomedicinal applications have been linked to various parts of *A. senegalensis*, as well as its usage as food and food additives. All parts of the plant have been found useful for traditional medicine applications in the treatment of yellow fever, tuberculosis, and smallpox [29]; snakebite and hernia [30]; difficulty in swallowing, gastritis, male sexual impotence, erectile dysfunction, tuberculosis [31,32,33]; chicken pox, kwashiorkor, and marasmus [34]; antidotes for venomous bites and diabetes [35]; malaria and open sores [36], etc. According to some research, the primary bioactive constituents of the plant include tannins, flavonoids, saponins, alkaloids, glycosides, steroids, volatile acids, and anthocyanin [37, 38, 39, 40, 41]. It was also discovered to include minerals such as Ca, K, Mg, Zn, Fe, Cu, Mn, Pb, and Cr as well as ascorbic acid and amino acids, making it a valuable source of nutrition [42].

The present study aims to investigate the *in vitro* antidiabetic activity of methanol extracts of *A. senegalensis* and *K. africana* stem bark and to understand the effect of the plant extracts on the yeast cell glucose uptake and α -amylase inhibitory activities.

MATERIALS AND METHODS

Both plant samples were collected from the International Center for Ethnomedicinal and Drug Development in Nsukka and were identified by Mr. Alfred Ozioko, a senior staff member of the herbarium. The stem barks of *A. senegalensis* and *K. africana* were washed, sliced into very small sizes, sun-dried, grounded into a fine powder, and stored in air-tight containers for further extraction.

Extraction of Phytochemicals

The air-dried and coarse powdered plant samples of *A. senegalensis* and *K. africana* (500g) were extracted with 500ml of methanol, n-hexane, and ethyl acetate each by maceration. Each of the solutions was agitated vigorously for 48 hours at room temperature. The extracts were filtered through Whatman No. 1 filter paper. The filtered extracts were concentrated using a rotary evaporator, under decreased pressure at about 55°C, and kept in a labeled sterile screw-sealed container at -20°C.

Phytochemical screening

Freshly prepared extracts were subjected to quantitative and qualitative phytochemical analysis to determine the presence or absence of flavonoids, alkaloids, terpenoids, saponins, carbohydrates, resins, tannins, reducing sugars, glycosides, and proteins using standard phytochemical procedures described by [43-45].

in vitro Antihyperglycemic Evaluations

Alpha-amylase Inhibitory Assay

According to a procedure previously described by [46] with slight modification, the alpha-amylase inhibitory test of methanol extracts of *A. senegalensis* and *K. africana* was examined. In summary, 0.5 ml of extracts was combined with 0.5 ml of α -amylase solution (0.5 mg/ml) with 0.02 M sodium phosphate buffer (pH 6.9 with 0.006 M NaCl). The mixture was incubated at room temperature for 10 min, and 0.5 ml of starch solution (1%) was added. The resulting mixture was incubated at room temperature for 10 min, and the reaction was stopped using 1 ml of dinitro salicylic acid colour reagent. The test tubes were then placed in a water bath (100°C for 5 minutes) and cooled until room temperature was achieved. The mixture was then diluted with 10 ml of deionized water, and absorbance was determined at 540 nm. The absorbance of blank (buffer instead of extract and amylase solution) and control (buffer instead of extract) samples were also determined. Acarbose was used as a standard drug. The inhibition of α -amylase was determined using the following equation:

%inhibition of α -Amylase = Abs (control) – Abs (Sample) / Abs (control) x 100.

Where;

Abs_{control} corresponds to the absorbance of the solution without extract (buffer instead of extract) and with α -amylase solution, and

Abs_{sample} corresponds to the solution with extract and α -amylase solution.

***in vitro* Evaluation of Yeast Cell Uptake of Glucose**

Commercial baker's yeast was washed by repeated centrifugation (3,000 \times g, 5 min) in distilled water until the supernatant fluid was clear and a 10% (v/v) suspension was prepared in distilled water. Different concentrations of extracts (1-5 mg/ml) were added to 1 ml of glucose solution and further incubated for 10 min at 37⁰ C. The reaction was started by adding 100 μ l of yeast suspension, vortexing it, and incubating it at 37⁰C for 60 minutes. The tubes were centrifuged (2,500 \times g, 5 min) and glucose was estimated in the supernatant. Metformin is taken as a standard antidiabetic drug. All tests were carried out in triplicates, and absorbance was measured at 540 nm. The formula below was used to determine the percentage increase in glucose uptake by yeast cells.

$$\text{Activity \%} = \frac{\text{Abs}_{\text{control}} - \text{Abs}_{\text{sample}}}{\text{Abs}_{\text{control}}} \times 100$$

Increase in glucose uptake

Where,

Abs_{control} is the absorbance of the control reaction (containing all reagents except the test sample), and

Abs_{sample} is the absorbance of the test sample.

STATISTICAL ANALYSIS

All experimental data were subjected to analysis of variance (ANOVA) using Statistical package for social science (SPSS) software (version 23.0).Duncan's new multiple range tests (DNMRT) were used to compare the group means obtained after each treatment with control measurements. Differences were considered significant when $p < 0.05$.All analyses were carried out in triplicates.

Results And Discussion

Table 1: Proximate Composition of *A. senegalensis* and *K. africana* Stem Bark Extracts

PARAMETERS	<i>Annona senegalensis</i>	<i>Kigelia africana</i>
Moisture Content (%)	10.0000 ^d ± 0.00000	10.6667 ^{c a} ± 1.15470
Ash Content (%)	6.0000 ^e ± 0.00000	6.0000 ^f ± 0.00000
Crude Fiber (%)	35.0000 ^b ± 0.0000	26.2500 ^b ± 1.25000
Protein (%)	6.8667 ^e ± 0.04041	9.7533 ^{cd} ± 0.04509
Oil Content (%)	8.5000 ^c ± 0.50000	8.0000 ^e ± 0.0000
Carbohydrate (%)	33.9300 ^a ± 0.74940	38.9933 ^a ± 1.29501

Table 2: Qualitative Phytochemical of *K. africana* and *A. senegalensis* Stem Bark Extracts

Phytochemicals	KaSH	KaSE	KaSM	AsSH	AsSE	AsSM
Saponins	-	+	+	+	-	++
Flavonoids	+	++	-	-	-	-
Alkaloids	-	-	+	+	+	+
Tannin	-	-	+++	-	++	+++
Steroids	++	+++	-	-	-	-
Terpenoids	+++	+++	-	+	++	++
Glycosides	++	++	-	++	++	+++

Key

+++ = Present in high concentration.

++ = Present in moderate concentration.

+ = Slightly or sparingly present.

- = Absent.

KaSH = n-hexane stem bark extract of *K. africana*

KaSE = Ethyl acetate stem bark extract of *K. africana*

KaSM = Methanol stem bark extract of *K. africana*

AsSH = n-hexane Stem bark extract of *A. senegalensis*

AsSE = Ethyl acetate Stem bark extract of *A. senegalensis*

AsSM = Methanol Stem bark extract of *A. senegalensis*

Table 3: Quantitative Phytochemical Composition of *K. africana* and *A. senegalensis* Stem Bark

Parameter	<i>Kigelia africana</i>	<i>Annona senegalensis</i>
Alkaloids (%)	12.0000 ^b ± 0.00000	2.5667 ^c ± 0.51316
Flavonoids (%)	21.1667 ^a ± 0.28868	13.0000 ^a ± 0.00000
Phenols (mg/g)	2.1867 ^e ± 0.02309	2.5567 ^c ± 0.01155
Saponins (%)	4.0000 ^d ± 0.00000	3.5833 ^b ± 0.14434
Tannins (mg/100g)	6.6367 ^c ± 0.51481	0.7967 ^d ± 0.06429

The proximate composition of *K. africana* and *A. senegalensis* samples presented above in **Table 1** showed that carbohydrates (38.99%) were higher in concentration in *K. africana* extract than in *A. senegalensis* extract with a concentration of 33.93% and crude fiber (35.00%) was seen to have a higher concentration in *A. senegalensis* extract than in *K. africana* with a concentration of 26.25%. With a concentration of 6.00%, ash content was the least abundant in both plant samples.

From the result of the qualitative phytochemical analysis (**Table 2**), it can be deduced that the methanol extract of *K. africana* stem bark extract showed the presence of secondary metabolites such as saponin, alkaloid, and tannins, which is in agreement with the phytochemical screening reported on the same plant by [47], while n-hexane and ethyl acetate extracts showed the presence of glycoside, terpenoid, steroids, and flavonoids, respectively. In n-hexane and ethyl acetate extracts, alkaloids and tannins were not detectable. Likewise, flavonoids, steroids, terpenoids, and glycosides were not detected in the methanol extract. The lack of steroids in the methanol extract is not unexpected, given they are lipophilic compounds [43]. On the other hand, *A. senegalensis* stem bark qualitative phytochemical screening confirmed the presence of alkaloids, terpenoids, and glycosides in all three extraction solvents. The methanol extract of the plant showed the presence of saponins, alkaloids, tannins, terpenoids, and glycosides, which is in agreement with the phytochemical screening described by [48]. Flavonoids and steroids were absent in all the extraction solvents. The absence of tannins in n-hexane extract contradicts the finding [40]. The higher amount of tannins in the methanol extract of both the plant samples is in agreement with our previous report, as tannins, being a polar compound, should be more in methanol. The presence of cardiac glycosides in all extracts of both plant samples except in the methanol extract of *K. africana* stem bark offers credibility to their antifungal, antibacterial, analgesic, anti-inflammation, antihypertensive, anticancer, diuretic, and emetic potency [49, 50]. The result of the quantitative phytochemical analysis of the methanol stem bark extract of *K. Africana* and *A. senegalensis* (**Table 3**) indicated that flavonoids were more abundant in both plants.

Table 4: α -amylase Inhibitory Activities of Methanol Extract of *K. africana* and *A. senegalensis* Stem Bark.

Concentration of samples (mg/mL)	<i>K. africana</i>	<i>A. senegalensis</i>
0.1	30.0000 ^d ± 0.00000	87.6100 ± 0.00000
0.2	23.7667 ^e ± 0.05774	82.0000 ± 0.00000
0.5	38.4333 ^c ± 0.30551	68.5367 ± 0.46479
1.0	39.9067 ^b ± 0.00577	60.9400 ± 0.04000
Standard Drug (Control)	56.0000 ^a ± 0.00000	56.0000 ± 0.00000

α -amylase is a type of intestinal enzyme that plays a vital function in carbohydrate digestion and glucose absorption. Suppression of the activity of digestive enzymes such as α -amylase, would delay the digestion of starch and oligosaccharides, which in turn inhibits the absorption of glucose and consequently reduces blood glucose. This study looked at the in vitro α -amylase inhibitory activity of methanol extracts of *K. africana* and *A. senegalensis* stem bark. The conclusion of the study demonstrated a good inhibitory effect of the plant samples. At a concentration of 0.1mg/ml, the extract exhibited its maximum value of inhibition (87.61%) when compared to Acarbose. Acarbose is a standard drug for α -amylase inhibitors. Acarbose showed α -amylase inhibitory activity of 56.0%. It is also important to note that the various levels of inhibition are not concentration-dependent, as indicated in **Table 4**. The stem bark extract of *A. senegalensis* demonstrated the highest level of inhibition (87.61%) at a 0.1 mg/ml concentration when compared to the standard drug (Acarbose). In this study, the methanol stem bark extract of *A. senegalensis* was found to be the most effective at inhibiting α -amylase enzymes in vitro. The plant extract might be utilized as a starch blocker as it limits or delays the absorption of starch into the body, mainly by preventing the hydrolysis of 1,4-glycosidic linkages of starch and other oligosaccharides into maltose, maltotriose, and other simple sugars. This inhibitory action could be tied to the presence of flavonoids and alkaloids, which play a vital role as antidiabetic agents [51]. The data revealed that the *A. senegalensis* extract is capable of boosting glucose absorption efficiently, which is similar to the result obtained by [52]. Hence, validating the claim of its usage as an antidiabetic drug in traditional medicine.

Table 5: Glucose Uptake in Yeast Cells Treated with Methanol Extract of *K. africana* and *A. senegalensis* Stem Bark.

Concentration of samples (mg/mL)	<i>K. africana</i>	<i>A. senegalensis</i>
0.1	11.3667 ^e ± 0.35119	0.4133 ^b ± 0.34641
0.2	12.0000 ^d ± 0.00000	2.0000 ^b ± 1.15470
0.5	17.6333 ^c ± 0.30551	31.4000 ^{ab} ± 0.47258
1.0	23.3333 ^b ± 0.35119	13.3333 ^b ± 0.17321
Standard Drug (Control)	68.4400 ^a ± 0.00000	68.4400 ^a ± 0.023090

The results of the present study showed that the methanol stem bark extract of *K. africana* and *A. senegalensis* increased glucose uptake in yeast cells at varying glucose concentrations. The result of glucose transport across the cell membrane in the yeast cell system is shown in **Table 5** after treatment of yeast cells with methanol stem bark of *K. africana* and *A. senegalensis*. No dose-dependent glucose uptake was observed in *A. senegalensis*, whereas dose-dependent glucose uptake was seen in *K. africana* which indicates an increase in concentration would enhance the capacity of yeast cells to uptake more glucose. The percentage of glucose uptake by both plant extracts was observed to be highest (31.40%) at a concentration of 0.5 mg/ml. The methanol stem bark extract of *A. senegalensis* showed the maximum glucose uptake (31.40%) at a 0.5 mg/ml concentration, while *K. africana* recorded 23.33 % of glucose uptake at a 1.0 mg/ml concentration. According to previous research [53, 54], the methanol stem bark extracts of *K. africana* and *A. senegalensis* are capable of improving glucose uptake and thus controlling blood glucose levels.

Conclusion

The results of this research suggest that *K. africana* and *A. senegalensis* stem bark have antihyperglycemic potential and might be used as an alternative medicine to treat diabetic patients by significantly decreasing the dose of conventional drugs. The plant extracts under study were shown to exhibit inhibitory activity against the α -amylase enzyme and glucose uptake in yeast cells; as a result, their therapeutic potential can be employed in the management and treatment of diabetes mellitus as well as a dietary health supplement. The bioactive compounds

in the stem bark of *K. africana* and *A. senegalensis* that are responsible for the inhibitory activity must be isolated, purified, and characterized. Further research is needed to determine if the plant has antidiabetic potential *in vivo* as well as a comprehensive pharmacological study to evaluate the likely toxicological implications of these antidiabetic plants.

References

1. Chatzigeorgiou, A., Halapas, A., Kalafatakis, K., and Kamper, E. (2009). The use of animal models in the study of diabetes mellitus. *In Vivo* 23: 245–258.
2. Palsamy, P., and Subramanian, S. (2010). Ameliorative potential of resveratrol on proinflammatory cytokines, hyperglycemia mediated oxidative stress and pancreatic beta-cell dysfunction in streptozotocin-nicotinamide-induced diabetic rats. *Journal of Cellular Physiology*. 224: 423–432.
3. Szkudelska, K., Nogowski, L., and Szkudelski, T. (2014). Adipocyte dysfunction in rats with streptozotocin-nicotinamide induced diabetes. *International Journal of Experimental Pathology*. 95: 86–94.
4. Szkudelski, T. (2012). Streptozotocin-nicotinamide-induced diabetes in the rat. Characteristics of the experimental model. *Experimental Biology and Medicine (Maywood)* 237: 481–490.
5. Pari, L., and Saravanan, R. (2007). Beneficial effect of succinic acid monoethyl ester on erythrocyte membrane bound enzymes and antioxidant status in streptozotocin-nicotinamide induced type 2 diabetes. *Chemico-Biological Interactions*. 169: 15–24
6. Nagappa, A.N., Thakurdesai, P.A., Venkat, R.N., and Jiwan, S. (2003). Antidiabetic activity of *Terminalia catappa* Linn fruits. *Journal of Ethnopharmacology*. 88: 45–50.
7. Bhandari, M.R., Nilubon, J.A. Hong, G., and Kawabata, J. (2008). Alpha-glucosidase and alpha-amylase inhibitory activities of Nepalese medicinal herb Pakhanbhed (*Bergeniaciliata* Haw.). *Food Chemistry*. 106 (1): 247 - 252
8. Perfetti, R., Barnett, P.S., Mathur, R., and Egan, J.M. (1998). Novel therapeutic strategies for the treatment of type 2 diabetes. *Diabetes/Metabolism Research and Review*. 14 (3): 207–225.
9. Surendran, S., Eswaran, M.B., Vijayakumar, M., and Rao, C.V. (2011). *In vitro* and *in vivo* hepatoprotective activity of *Cissampelos pareira* against carbon tetrachloride-induced hepatic damage. *Indian Journal of Experimental Biology*. 49 (12): 939–945.
10. Grover, J.K., Yadav, S., and Vats, V. (2002). Medicinal plants of India with antidiabetic potential. *Journal of Ethnopharmacology*. 81(1): 81–100.

11. Mukherjee, P.K., Maiti, K., Mukherjee, K., and Houghton, P.J. (2006). Leads from Indian medicinal plants with hypoglycemic potentials. *Journal of Ethnopharmacology*. 106(1): 1–28.
12. Ivorra, M.D., Paya, M., and Villar, A. (1989). A review of natural products and plants as potent antidiabetic drugs. *Journal of Ethnopharmacology*. 27(3): 243-276.
13. Konkon, N.G., Adjoungoua, A.L., Manda, P., Simaga, D., N'Guessan, K.E., and Kone B.D. (2010). Toxicological and phytochemical screening study of *Mitragyna Inermis* (willd.) O k tze (Rubiaceae), anti-diabetic plant. *International Journal of Medicinal Plants Research*. 2(10): 279 – 284.
14. WHO (1980) World Health Organization expert committee on Diabetes Mellitus: Second Report. Technical Report Series. WHO, Geneva, 646: 61.
15. Grace, O.M., Light, M.L., Lindsey, K.L., Mulholland, D.A., Van Staden, J., Jager, A.K., and Eloff, J.N. (2002). Antibacterial activity and isolation of active compounds from fruit of the traditional African medicinal tree *Kigelia africana*. *South African Journal of Botany*. 68:220–222.
DOI: 10.1016/S0254-6299(15)30424-5.
16. Adam, I.Y., and Alhameed, I.A. (2013). *Kigelia africana* fruits extract anti-hepato-toxic effects on male Wistar rat's liver destruction induced by CCl₄. *Asian Journal of Medical Sciences*. 5(1):26–32.
17. Chih-Cheng, L., Lu-Te, C., and Yuan-Chen, W. (2012). Polyphenol and antioxidant content of *Kigelia africana* leaf from Ghana. *Food*. 6(1):1–5.
18. Jackson, S., and Beckett, K. (2012). Sausage Tree *Kigelia pinnata*: An Ethnobotanical and Scientific Review. *American Botanical Council*. 48-59.
19. Oyedeji, F.O., Bankole-Ojo, O.S. (2012). Quantitative evaluation of the antipsoriatic activity of sausage tree (*Kigelia africana*). *African Journal of Pure and Applied Chemistry*. 6(13). 214-218.
20. Houghton, P.J., Jäger, A.K. (2002). The sausage tree (*K. pinnata*): Ethnobotany and recent scientific work. *South African Journal of Botany*. 68: 14–20.
21. Akintunde, J.K., Akintunde, D.G., Ironde, E.A.; Babaita, K. (2016). Antioxidants from the stem bark of *K. africana* inhibit free radicals and membrane lipid damage in rat testes in vitro. *Oxidants and Antioxidants in Medical Sciences*. 5: 63–69.
22. Azu, O.O., Duru, F., Abraham, A., Osinubi, A.A., Oremosu, A., Noronha, C.C., Elesha, S.O. (2010). Histomorphometric effects of *K. africana* (Bignoniaceae) fruit extract on the testis following short-term treatment with cisplatin in male Sprague-Dawley rats. *Middle East Fertility Society Journal*. 15: 200–208.
23. Kuete, Victor. (2013). Medicinal Plant Research in Africa: Pharmacology and Chemistry. 10.1016/B978-0-12-405927-6.0000.
24. Patrizia, P., Giuseppina, A., Stelama, M., Marisa, M., Rok Rita P.A (2005). Anti-inflammatory activity of a verumumaside from *Kigelia africana* and evaluation of cumucous imitation in cell cultures and reconstituted human epidemis. *Nat Prod* 68:1610-4

25. Owolabi, O.F., and Omogbai, E.K.I. (2007). Analgesic and anti-inflammatory activities of the ethanolic stem bark extract of *Kigella africana* (Bignoniaceae). *African journal Biotechnology*. 6(5):582-585.
26. Oyelami, O.A., Yusuf, K.O., and Oyelami, A.O. (2012). The Use of *K. africana* in the Management of polycystic Ovary Syndrome (PCOS). *Chinese Medical Journal (Engl.)*. 3: 1-3.
27. Eldeen, I.M.S., and Van Staden, J. (2007). *In vitro* pharmacological investigation of extracts from some trees used in Sudanese traditional medicine. *South African Journal Botany*. 73: 435-440.
28. Samuel, E.O., Edifonfon, A.N., Omolola, T.F., Keuwe, B.E., Oluyemsi, F.K. (2016). *Annona senegalensis*: A review of its ethnomedicinal uses, Bioactivity and Phyto compound. *Journal of Pharmacognosy and phytochemistry* 5(2):211-219.
29. Mustapha, A., Owuna, G., and Uthman, I. (2013). Plant Remedies Practiced by Keffi People in the Management of Dermatitis. *Journal of Medicinal Plant Studies*. 1(5):112-118.
30. Dambatta, S., Aliyu, B.A. (2011). Survey of Major Ethnomedicinal plants of Kano North, Nigeria, their Knowledge and Uses by Traditional Healers. *Bayero Journal of Pure and Applied Sciences*. 4(2):28-34.
31. Ofukwu, R., Ayoola, A., Akwuobu, C. (2008). Medicinal Plants Used in the Management of Tuberculosis in Humans and Animals by Idoma Tribe of North Central Nigeria. *Nigerian Veterinary Journal*. 29(2):25-30.
32. Noumi, E., Safiatou, M. (2015). Some Investigations on the Traditional Pharmacopoeia about Venomous Bites and Stings from Scorpions, Snakes, and Spiders; in the Hina Subdivision, Far North, Cameroon. *British Journal of Pharmaceutical Research*. 344-358. DOI: [10.9734/BJPR/2015/8894](https://doi.org/10.9734/BJPR/2015/8894).
33. Jiofack, T., Fokunang, C., Guedje, N., Kemeuze, V., Fongzossie, E., Nkongmeneck, B., Mapongmetsem, P., and Tsabang, N. (2010). Ethnobotanical Uses of some Plants of Two Ethnoecological Regions of Cameroon. *International Journal of Medicine and Medical Sciences*. 2(3):60-79.
34. Faleyimu, O., Ahmad, I., and Akinyemi, O. (2010). Herbal Approach to the Treatment of Chickenpox: A Case study of Giwa Local Government, Kaduna State. Nigeria. *International Journal of Biomedical and Healthcare Science*. 6(1):13-18.
35. Ahombo, G., Ampa, R., Diatwa, M., Mpati, J., Abena, A., and Ouamba, J. (2012). Investigating related diabetes therapeutic plants used in traditional medicine at Brazzaville. *Journal of Medicinal Plant Research*. 6(44):5630-5639.
36. Traore, M.S., Baldé, M.A., Diallo, M.S., Baldé, E.S., Diané, S., Camara, A., Diallo, A., Balde, A., Keita, A., Keita, S.M., Oulare, K., Magassouba, F.B., Diakite, I., Diallo, A., Pieters, L., and Balde, A.M. (2013). Ethnobotanical survey on medicinal plants used by Guinean traditional healers in the treatment of malaria. *Journal of Ethnopharmacology*. 150(3):1145-1153. DOI: [10.1016/j.jep.2013.10.048](https://doi.org/10.1016/j.jep.2013.10.048).

37. Jada, M., Usman, W., Adamu, Y. (2014). *In vitro* Antimicrobial Effect of Crude Tannins Isolated from the Leaf of *Annona senegalensis*. *International Journal of Biochemistry Research and Review*. 4(6):615-623,
38. Jada, M., Usman, W., and Olabisi, A.(2015). Crude Flavonoids Isolated from the Stem Bark of *Annona senegalensis* have Antimicrobial Activity. *Journal of Advances in Biology and Biotechnology*. 2(1):24-29.
39. Afolabi, F., and Afolabi, O.(2013). Phytochemical Constituents of Some Medicinal Plants in South West, Nigeria. *IOSR Journal of Applied Chemistry*. 4(1):76-78. DOI: [10.9790/5736-0417678](https://doi.org/10.9790/5736-0417678).
40. Ijaiya, I., Arzika, S., Abdulkadir, M. (2014). Extraction and Phytochemical Screening of the Root and Leaf of *Annona senegalensis* (Wild Custard Apple). *Academic Journal of Interdisciplinary Studies*. 3(7):9-15.
41. Mpiana, P. (2012). Anti-sickling properties, thermal and photochemical degradations of anthocyanin extracts from *Annona senegalensis* (Annonaceae). *International Journal of Biological and Chemical Sciences*. 6(5):2241-2251.
42. Yisa, J., Egila, J.N. and Darlinton, A.O. (2010). Chemical composition of *Annona senegalensis* from Nupe land, Nigeria. *African Journal of Biotechnology*. 9: 4106-4109.
43. Anarado, C.E., Ajiwe, V.I.E., Obumselu, O.F., Anarado, C.J.O., Okafor, N.P. (2021). Phytochemical, Proximate, *in-vitro* Anti-malarial, Anti-inflammatory and Antimicrobial Screening of Leaf Extracts of *Combretum platypterum* (Welw) Hutch & Dalziel. *Journal of Pharmaceutical Research International*. 33(52A): 39-61. DOI: [10.9734/JPRI/2021/v33i52A33556](https://doi.org/10.9734/JPRI/2021/v33i52A33556)
44. Anarado, C.E., Ajiwe, V.I.E., Anarado, C.J.O., Obumselu, O.F., Onuegbu, T.U., Okafor, S.E. (2021). Phytochemical, Proximate, *In-vitro* Antimalarial and Antimicrobial screening of leaf Extracts of *Justicia carnea* Lindl. *International Research Journal of Pure and Applied Chemistry*. 22(8), 55-73. DOI: [10.9734/IRJPAC/2021/v22i830428](https://doi.org/10.9734/IRJPAC/2021/v22i830428).
45. Anarado, C.E., Anarado, C.J.O., Umedum, N.L., Ogbodo, Q.M. (2020). Comparative phytochemical and anti-microbial studies of leaf, stem, and root of *Spathodea campanulata*. *Asian Journal of Applied Chemistry Research*. 6(1): 10-20. DOI: [10.9734/AJACR/2020/v6i130148](https://doi.org/10.9734/AJACR/2020/v6i130148).
46. Ranila, L.G., Kwon, Y.I., Apostoldis, E., Shetty, K. (2010). Phenolic Compounds: Antioxidant activity and *in vitro* inhibitory potential against key enzymes relevant for hyperglycemia and hypertension of commonly used medicinal plants, herbs, and spices in Latin America. *Bioresource Technology*. 101(12): 4676-4689.
47. Kaman, J.K, Nthiga, P.M, Safari, V.C, Njagi, S.M., Mwonjaria, J.K., Ngugi, M.P., and Ngeranwa, J.J.N. (2016). Antipyretic properties of methanol stem bark extracts of *Acacia hockii* De wild and *Kigelia africana* (Lam.) Benth in Wistar Rats. *Journal of Pharmacognosy and Natural products*. 2:3. DOI: [10.4172/2472-0992.1000118](https://doi.org/10.4172/2472-0992.1000118).

48. Ajaiyeoba, E., Falade, M., Ogbale, O., Okpako, L., and Akinboye, D. (2006). *In vivo* antimalarial and cytotoxic properties of *Annona senegalensis* extract. *African Journal of Traditional, Complementary and Alternative Medicines*. 3(1): 137-141.
49. Morsy, N. (2017). Cardiac glycosides in medicinal plants. *Aromatic and Medicinal Plants -Back to Nature. Intech Open*. 29-45.
50. Fu, J., Wu, Z., Zhang, L. (2019). Clinical applications of the naturally occurring or synthetic glycosylated low molecular weight drugs, *Progress in Molecular Biology and Translational Science*. Academic Press. 163: pp 487-522. <https://doi.org/10.1016/bs.pmbts.2019.03.005>.
51. Chowdhury, S.S., Tareq, A.M., Tareq, S.M., Farhead, S., and Sayeed, M.A. (2021). Screening of antidiabetic and antioxidant potential along with phytochemicals of *Annona* genus: a review. *Future Journal of Pharmaceutical Sciences*. 7: 144.
52. Beverley, B., and Eschwège, E. (2004). Diagnosis and classification of diabetes and impaired glucose tolerance. *Textbook of Diabetes*. Ed: John C Pickup and Gareth Williams Third edition, 2: 2.1-2.11.
53. Abirami, N., and Natarajan, B. (2014). Isolation and Characterization of (4Z, 12Z)-Cyclopentadeca-4, 12-Dienone from Indian Medicinal Plant *Grewia hirsuta* and its Hyperglycemic Effect on 3T3 and L6 Cell Lines. *International Journal of Pharmacy and Biological Sciences*. 6:393-398.
54. Ahmed, F., Sairam, S., and Urooj, A. (2009). Effect of various Ayurvedic formulations and medicinal plants on carbohydrate hydrolyzing enzymes and glucose uptake by yeast cells - an *in vitro* study. *Journal of Pharmacy Research*. 2:563-8.