

Effect of Dietary Inclusion of Agricultural Waste-Derived Activated Charcoal on Hematological and Serum Biochemical Indices of Layer Chickens

Abstract

The ban on sub-therapeutic dietary inclusion of antibiotics has necessitated the inclusion of natural additives such as activated charcoal (AC) produced from agricultural wastes. One hundred and twenty Isa Brown layer chickens aged 16 weeks with good management and vaccination history were used for this experiment. The birds were divided into four groups (G1-G4) of thirty birds each with each group further replicated three times comprising of 10 birds each. Chickens in G2-G4 were fed layer mash which contained 0.5g/kg, 1.0g/kg and 1.5g/kg of AC, respectively with G1 as control group and reared on deep litter. Blood samples were collected from the chickens at the end of 20th, 28th and 33rd week of the experiment and used for hematological and serum biochemical assays. Data collected were subjected to analysis using ANOVA. Hematological indices (RBC, Hb, PCV and lymphocytes) were significantly improved ($P < 0.05$) in the supplemented treatment group (G2, G3 and G4) at 28th and 33rd week than the control with G3 having the highest value. The MCV of G1 and G4 were both higher ($P < 0.05$) while the MCH, MCHC and neutrophil counts of G1 were significantly higher than other layer groups. The cholesterol level of G1 was significantly higher ($P < 0.05$) than the supplemented groups (G2-G4) whose total proteins were significantly higher ($P < 0.05$) than G1. AC is recommended as feed additive for improvement of hematological indices and for use in cases of hypercholesterolemia.

Keywords: Activated charcoal, agricultural wastes, palm kernel shell, palm fruit fibre and pig dung

Introduction

Activated charcoal (AC) is a solid, porous, tasteless and black carbonaceous material (AAFCO, 2012) produced from a variety of carbon containing materials including agricultural residues and wastes. It is manufactured by a two step process of carbonization followed by oxidation (activation) in a steam chamber (Schmidt *et al.*, 2019). Numerous studies have described the absorbance characteristics and potential clinical benefits of charcoal (Maklad *et al.* 2012) and its widely acceptance as an essential tool in disease management in human and veterinary medicines. Emerging reports however, showed that activated charcoal adsorbs more toxins than any other natural substance thus rendering them ineffective and harmless (Cooney, 1980; Maklad *et al.*, 2012). It has non-nutritive and non-digestible components and is effective in the elimination of mycotoxins, such as aflatoxins as well as pesticide residues that occasionally contaminate feed ingredients (Huwig *et al.*, 2001; Bhatti *et al.*, 2018; Schmidt *et al.*, 2019).

Research in 2005 showed that the quantity of agricultural wastes in Nigeria stood at 61 million tonnes per year of animal waste and 83 million tonnes of crop residues (Agba *et al.*, 2010; Akorede *et al.*, 2017) with

the major agricultural crops biomass feedstocks as millet, yam, cassava, sorghum, rice, groundnut, oil palm, sugar cane and soya-beans (Mohammed *et al.*, 2013). Wastes from livestock activities include solid waste such as pig dung and organic materials in the slaughter houses and liquid waste such as urine which are capable of generating pollutants known for their characteristic offensive odour (Iregbu *et al.*, 2014). Though endowed with these abundant agricultural wastes, Nigeria is yet to fully harness them to play a significant role in the production of value added products such as activated charcoal and renewable energy production (Oyedepo, 2012).

Dim *et al.* (2018) reported that the white blood cell (WBC) count and the packed cell volume (PCV) were not affected while the haemoglobin concentration (Hb) and the red blood cell (RBC) count and cholesterol level were significantly improved in broiler chickens whose diets were supplemented with activated charcoal. Abdel-Latif *et al.* (2021) also observed that the inclusion of activated charcoal did not significantly affect hematological indices of Nile tilapia fish except lymphocytes counts which were significantly increased in all AC supplemented groups compared with control. More so, Abdel-Latif *et al.* (2021) noted that there was a significant increase in serum albumin, globulin and total protein values a significant decrease in ALP, AST and ALT activities in all AC supplemented groups. Mabe *et al.* (2018) reported that the supplementation of juvenile common carp diet with 4% dietary bamboo charcoal improved serum indicators such as ALT, AST, TP, total cholesterol and density lipoprotein (HDL). Transaminases are biomarkers of liver functions and the increase of these enzymes is regarded as an indicator of liver damage while a decrease in their values suggest a healthy status of the liver. Majewska *et al.* (2009) observed that dietary supplementation of charcoal at 0.3% did not have significant effect on the RBC, WBC, Hb and hematocrit values of broilers fed aflatoxin-B1 contaminated feeds. Boonanuntanasarn *et al.* (2014) also observed that the inclusion of activated charcoal did not significantly affect RBC, Hb or hematocrit values of Nile tilapia in a 28-days feeding trial while Bawala *et al.* (2007) and Kana *et al.* (2014) reported low levels for hematological parameters, especially the RBC values with the use of poor quality feed.

Hematological and serum biochemical profile are important physiological indicators for evaluating the overall performances and health status of animals (Dawood *et al.*, 2020; Abdel-Hameed *et al.*, 2021). Hence, in veterinary and medical practice, a diagnosis is considered incomplete if information from history and clinical examination is not combined with laboratory test results, including results of assessment of hematological and serum biochemical indices (Ugochukwu, 2001). The increase leucocyte count reflects humoral and cellular immune response against pathogenic agent that causes diseases (Moye and Schute, 2008). It should be recalled that the animal's health could be measured from the total leucocyte count as increased leucocyte count represents the body's immune response while decreased

WHC may translate into non-existence of infection (Sufiriyanto and Emmy, 2008). Activated charcoal may contribute to enhancement of blood profiles and immune response since it has been reported to increase body weight in animals (Majewska *et al.*, 2009; Jiya *et al.*, 2013). Hence, the need to determine the effects of its inclusion in layer feeds on hematology and serum biochemistry of layer chickens.

Location of the study

The study was carried out at the Teaching and Research Farm of Michael Okpara University of Agriculture Umudike, Umuahia, Abia State, located within the South East agro-ecological zone of Nigeria with geographical coordinates of 5.4801° N and 7.5437° E.

Materials and methods

One hundred and twenty Isa Brown layer chickens aged 16 weeks with good management and vaccination history were used for this experiment. The birds were divided into four groups (G1-G4) of thirty birds each with each group further replicated three times comprising of 10 birds each. Chickens in G2-G4 were fed layer mash which contained 0.5g/kg, 1.0g/kg and 1.5g/kg of AC, respectively with G1 as control group and reared on deep litter. Blood samples were collected at the end of 1st, 4th and 6th week of the experiment from the wing vein of the birds into EDTA (Ethylene-diamine tetra-acetic acid) and plain bottles for hematological and serum biochemical analysis respectively. The erythrocyte was counted using the haemocytometer method as describe by Schalm *et al.* (1975) while the hemoglobin concentration was determined according to the techniques described by Cole (1986). In determining the packed cell volume (PCV), the Wintrob microheamatocrit tube method was used while other hematological indices were calculated according to the formula reported by Schalm *et al.* (1975). Serum biochemical tests were carried out using Randox commercial test kit specific for each biochemical parameter in accordance with standard procedures prescribed by the producer Randox Laboratories (UK). The serum parameters analyzed include the following included total serum protein, serum albumin and globulin, urea, serum creatinine concentration, alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase activity (ALP), total bilirubin content and total serum cholesterol.

Production of activated charcoal

Palm kernel shell and palm fruit fibre were collected from a palm oil mill while freshly voided pig dung was collected from pig farms using plastic container. The materials were carefully collected to avoid contamination with sand or other objects. Each material was sun-dried to constant weight and crushed manually using a wooden pestle and mortar. The materials were then blended together at a ratio of 4:3:3 weight for weight for pig dung, palm kernel shell and palm fruit fibre, respectively and used to produce

the activated charcoal which was subsequently supplemented in layerfeeds. The physical method of activated charcoal production described by Gunamartha and Widana (2018) was employed in this study.

Results

Tables I, II and III presents the hematological indices of the experimental birds at twenty, twenty eight and thirty three weeks of age, respectively while the serum biochemical indices of the experimental birds at week twenty, twenty eight and thirty three weeks of age are presented in Tables IV, V, and VI respectively.

Table I: Hematological indices of 20weeks of age layer chickens fed varying dietary levels of AC

Parameters	Group 1	Group 2	Group 3	Group 4
RBC ($\times 10^6/\text{mm}^3$)	3.61 \pm 0.18	3.38 \pm 0.13	3.44 \pm 0.06	3.49 \pm 0.11
PCV (%)	31.67 \pm 1.53	30.00 \pm 1.00	30.33 \pm 1.53	30.67 \pm 1.16
HB (g/dl)	11.03 \pm 0.25 ^b	10.13 \pm 0.23 ^a	10.17 \pm 0.15 ^a	10.63 \pm 0.32 ^b
WBC ($\times 10^3/\text{mm}^3$)	40.57 \pm 1.12 ^a	42.03 \pm 0.38 ^a	40.53 \pm 1.78 ^a	45.07 \pm 0.32 ^b
Platelets ($\times 10^3/\text{mm}^3$)	113.67 \pm 8.96	126.00 \pm 5.29	122.33 \pm 13.50	119.67 \pm 6.51
MCV (fl)	90.44 \pm 8.82	91.91 \pm 6.30	90.86 \pm 5.43	90.81 \pm 5.57
MCH (pg)	30.56 \pm 1.00	30.02 \pm 0.65	29.53 \pm 0.93	30.52 \pm 1.27
MCHC (g/dl)	34.91 \pm 2.39	33.80 \pm 1.35	33.59 \pm 2.16	34.72 \pm 1.93
Neutrophils (%)	51.67 \pm 1.53 ^a	54.00 \pm 1.00 ^{ab}	54.68 \pm 2.31 ^{ab}	56.66 \pm 1.54 ^b
Lymphocytes (%)	34.67 \pm 5.13	37.00 \pm 1.00	36.33 \pm 0.58	35.33 \pm 1.53
Monocytes (%)	6.67 \pm 0.58 ^b	6.00 \pm 0.00 ^{ab}	6.33 \pm 0.58 ^b	5.33 \pm 0.58 ^a

Eosinophils (%)	3.67±0.58	3.00±0.00	2.67±1.16	2.67±0.58
-----------------	-----------	-----------	-----------	-----------

Table II: Hematological indices of 28 weeks of age layer chickens fed varying dietary levels of AC

Parameters	Group 1	Group 2	Group 3	Group 4
RBC (x10 ⁶ /mm ³)	2.59±0.04 ^a	3.23±0.01 ^b	3.72±0.08 ^c	3.32±0.06 ^b
PCV (%)	25.67±0.58 ^a	30.33±0.58 ^b	34.00±1.00 ^c	33.00±1.00 ^c
Hb (g/dl)	11.40±0.20 ^a	12.20±0.20 ^b	12.97±0.20 ^c	12.73±0.15 ^c
WBC (X10 ³ /mm ³)	31.70±0.98	35.07±1.21	36.30±1.28	35.10±0.70
Platelets (x10 ³ /mm ³)	143.67±3.89	161.67±14.15	134.00±1.36	149.33±6.11
MCV (fl)	100.47±1.75 ^b	94.88±1.49 ^a	92.12±1.72 ^a	100.35±2.85 ^b
Neutrophils (%)	30.67±1.53 ^c	26.67±2.08 ^b	21.67±1.53 ^a	23.33±2.08 ^{a,b}
Lymphocytes (%)	60.00±1.00 ^a	63.67±1.53 ^b	70.00±2.00 ^c	66.33±2.08 ^b
Monocytes (%)	6.67±0.58 ^b	7.33±0.58 ^b	4.67±1.53 ^a	7.67±0.58 ^b
Eosinophils (%)	2.67±0.58	2.33±0.58	2.67±0.58	2.67±0.58
MCH (pg)	44.03±1.43 ^c	37.81±0.50 ^b	34.86±0.16 ^a	38.31±0.23 ^b
MCHC (g/dl)	45.86±3.56 ^b	41.20±2.08 ^{a,b}	38.94±1.51 ^a	39.50±2.38 ^a

Results are presented as mean ± standard deviation (n = 3). The results with different letter superscripts are significantly different (P < 0.05) from any paired mean across the row.

Table III: Hematological indices of 33 weeks of age layer chickens fed varying dietary levels of AC

Parameters	Group 1	Group 2	Group 3	Group 4
RBC (x10 ⁶ /mm ³)	2.73±0.05 ^a	3.31±0.08 ^b	3.88±0.02 ^d	3.75±0.05 ^c
PCV	26.67±0.58 ^a	33.00±1.00 ^b	36.67±1.16 ^c	35.33±0.58 ^c
Hb	12.10±0.10 ^a	12.47±0.49 ^a	13.47±0.06 ^b	13.50±0.10 ^b
WBC (x10 ³ /mm ³)	33.00±0.66 ^a	33.93±1.10 ^{a,b}	36.07±1.04 ^b	34.87±1.45 ^{a,b}
Platelets (x10 ³ /mm ³)	131.67±18.77	139.67±8.51	145.67±6.66 ^a	146.00±5.20
MCV (fl)	97.56±1.19 ^{b,c}	99.59±1.28 ^c	94.58±2.71 ^{a,b}	94.22±0.59 ^a
MCH (pg)	44.27±0.61 ^d	37.62±1.11 ^c	34.74±0.22 ^a	36.00±0.19 ^b
MCHC (g/dl)	45.39±1.06 ^b	37.77±0.65 ^a	36.75±1.07 ^a	37.33±1.16 ^a
Neutrophils (%)	31.33±1.53 ^c	30.33±0.58 ^{b,c}	28.33±1.53 ^{a,b}	27.33±1.16 ^a
Lymphocytes (%)	60.67±1.53	61.67±1.53	63.33±1.53	63.00±2.00

Monocytes (%)	5.67±0.5	5.33±0.58	5.67±0.58	6.33±0.58
Eosinophils (%)	2.33±0.58	2.67±0.58	2.67±0.58	3.33±0.58

Table IV: Serum biochemical indices at 20 weeks of age for layer chicken fed varying dietary levels of activated charcoal

Parameters	Group 1	Group 2	Group 3	Group 4
TP (g/dl)	4.87±0.05 ^a	5.23±0.08 ^b	5.57±0.13 ^{b,c}	5.38±0.22 ^c
Albumin (g/dl)	2.68±0.12 ^a	3.20±0.08 ^b	3.31±0.17 ^b	3.24±0.11 ^b
Globulin (g/dl)	2.19±0.09	2.01±0.13	2.25±0.32	2.14±0.11
ALT (μ/l)	27.67±1.53	25.00±2.65	26.33±1.53	27.33±2.52
AST (μ/l)	43.00±9.85	38.00±2.00	39.67±2.08	38.33±7.37
ALP (μ/l)	101.33±3.06	101.00±2.00	103.33±1.53	103.00±2.65
Cholesterol (mg/dl)	95.45±2.43 ^a	112.72±2.23 ^b	109.01±390 ^b	110.87±1.60 ^b
Urea (mg/dl)	10.01±0.28 ^a	10.01±0.22 ^a	11.11±0.19 ^b	10.08±0.22 ^a
Creatinine (%)	1.01±0.04 ^b	0.94±0.04 ^{a,b}	0.97±0.05 ^{a,b}	0.90±0.04 ^a
Bilirubin (mg/dl)	0.54±0.03 ^{a,b}	0.60±0.03 ^b	0.61±0.04 ^b	0.51±0.07 ^a

Results are presented as mean ± standard deviation (n = 3). The results with different letter superscripts are significantly different (P < 0.05) from any paired mean across the row.

Table V: Serum biochemical indices at 28 weeks of age for layer chicken fed varying dietary levels of activated charcoal

Parameters	Group 1	Group 2	Group 3	Group 4
TP (g/dl)	3.53±0.06 ^a	3.91±0.03 ^a	3.88±0.58 ^a	4.07±0.06 ^a
AST (μ/l)	35.33±2.52 ^b	30.67±1.16 ^a	31.00±1.00 ^a	30.33±0.58 ^a
ALT (μ/l)	21.67±1.53 ^a	20.67±1.16 ^a	19.67±1.53 ^a	20.33±2.08 ^a
ALP (μ/l)	82.00±3.00 ^b	79.67±1.53 ^{a,b}	75.33±3.06 ^a	78.33±2.52 ^{a,b}
Albumin (g/dl)	2.30±0.14 ^a	2.44±0.13 ^a	2.99±0.07 ^b	2.87±0.04 ^b
Globulin (g/dl)	1.24±0.15 ^a	1.48±0.16 ^b	1.22±0.04 ^a	1.20±0.05 ^a
Cholesterol (mg/dl)	104.50±3.20 ^b	99.10±0.84 ^{a,b}	92.47±2.25 ^a	94.66±4.62 ^a
Bilirubin (mg/dl)	0.61±0.02 ^c	0.51±0.02 ^b	0.41±0.01 ^a	0.49±0.03 ^b

Urea (mg/dl)	12.88±0.60 ^b	11.75±0.43 ^a	12.38±0.28 ^{a,b}	12.11±0.12 ^{a,b}
Creatinine (mg/dl)	0.77±0.03 ^{a,b}	0.72±0.03 ^a	0.79±0.03 ^b	0.78±0.03 ^b

Table VI: Serum biochemical indices at 33weeks of age for layer chicken fed varying dietary levels of activated charcoal

Parameters	Group 1	Group 2	Group 3	Group 4
TP (g/dl)	3.63±0.06 ^a	3.94±0.05 ^b	4.19±0.11 ^c	4.26±0.04 ^c
AST (μ/l)	36.67±2.08 ^a	35.00±1.00 ^a	33.33±2.31 ^a	32.67±2.52 ^a
ALT (μ/l)	22.33±2.52 ^a	24.67±1.53 ^a	23.67±1.53 ^a	22.33±3.22 ^a
ALP (μ/l)	80.67±2.08 ^a	86.00±2.65 ^b	79.00±1.73 ^a	79.33±3.06 ^a
Albumin (g/dl)	2.24±0.07 ^a	2.71±0.09 ^b	2.93±0.04 ^c	3.04±0.08 ^c
Globulin (g/dl)	1.38±0.02 ^a	1.23±0.13 ^a	1.26±0.14 ^a	1.22±0.08 ^a
Cholesterol (mg/dl)	108.00±5.92 ^b	101.80±3.92 ^{a,b}	97.88±1.33 ^a	98.27±1.31 ^a
Bilirubin (mg/dl)	0.64±0.18 ^a	0.60±0.01 ^a	0.49±0.02 ^a	0.53±0.01 ^a
Urea (mg/dl)	11.07±0.13 ^b	10.05±0.22 ^a	10.07±0.16 ^a	11.40±0.30 ^b
Creatinine (mg/dl)	0.70±0.02 ^{a,b}	0.71±0.02 ^{a,b}	0.69±0.01 ^a	0.73±0.02 ^b

Results are presented as mean ± standard deviation (n = 3). The results with different letter superscripts are significantly different (P < 0.05) from any paired mean across the row.

Discussion

Hematological Parameters:

More effects of the activated charcoal were noticed on the hematological indices at twenty eight and thirty three weeks of age than at twenty weeks with RBC, Hb and PCV and lymphocytes significantly higher in the supplemented layer groups (G2, G3 and G4) when compared with the control with G3 having the highest value. The MCV of G1 and G4 were both higher while the neutrophil counts of G1 was higher than other groups and closely followed by G2. The monocytes counts of G2, G4 and G1 were significantly higher than G3 while the MCH and MCHC of G1 were greater than other layer groups with no significant differences in other hematological parameters determined.

Notwithstanding, in all the weeks, the results of the PCV, Hb and RBC values were in good agreement with normal ranges of hematological parameters in chicken with only the MCHC, total WBC, lymphocytes and platelets exceeding their normal ranges. The normal ranges of the hematological indices in chickens are RBC $2.5-3.5 \times 10^6/\mu\text{l}$, PCV 22-35%, Hb 7-13g/dl, WBC $12-30 \times 10^3/\mu\text{l}$, MCV 90-140fL, MCH 33-47 Pg/cell and MCHC 26-35g/dl (Bounous & Stedman, 2000). The mean corpuscular hemoglobin concentration (MCHC) measures the concentration of hemoglobin in a given volume of packed red blood cells and is useful in the characterization of erythrocytes, especially in the evaluation of anemia (Igwe *et al.*, 2017). The significant higher MCHC value across the groups (especially in group 1) at week 28 and 33 could be associated with the presence of immature erythrocytes in circulation (Polychromasia) which are produced during erythrocyte regeneration (Irrizarry-Rovira, 2004; Samour, 2009).

Leukocytosis most especially could be due to heterophilia, lymphocytosis and monocytosis which can be caused by inflammation, infections or heat stress (Irrizarry-Rovira, 2004; Igwe *et al.*, 2017). These must have stimulated the hematopoietic production of the granulocyte (lymphocytes and platelets) precursors which occur when there is increased demand for them in the peripheral tissue (Campbell & Cole, 1986; Irrizarry-Rovira, 2004; Juul-Madsen *et al.*, 2008). Hence, the most cause of an increase in the WBC count is the normal response of the body to an infection, certain drugs and release of immature or abnormal WBC from bone marrow into the blood.

Laying hens have been reported to have predominant lymphocytes in good health. The lymphocytes are found in significant number in the ovaries and oviduct where they are indicators of stress (Latimer and Bienzle, 2000). This is true considering the fact that the point of lay birds were brought in November and raised till March when heat stress was at its peak in this location. Lymphopoiesis or enhanced lymphocytes from lymphomyeloid tissue act as a defense mechanism to tolerate infections or environmental insults such as heat stress (Irrizarry-Rovira, 2004).

Serum Biochemical Parameters:

Serum parameters just like hematological indices are used to assess the clinical and physiological responsiveness and well-being of chicken (Sharma *et al.*, 2015) which can be influenced by feed, medication, toxic compounds, infections, age and sex of the birds (Huff *et al.*, 2008). Total protein is made up of albumin and globulin and the normal ranges of total protein, albumin and globulin in chicken's blood are 3.0-4.9mg/dl, 1.17-2.74mg/dl (Meluzzi *et al.*, 1992) and 2.33-3.33 (Makama *et al.*, 2021) respectively. The results of the total protein, albumin and globulin obtained were in good agreement with this range. Albumin is a serum protein synthesized in the liver and is responsible for

transporting insoluble substance in the blood and aids in maintaining oncotic pressure (Fischbatch & Dunning, 2009). Total protein and albumin are indicators of protein reserve in the body (Makama *et al.*, 2021). A higher concentration of albumin usually denotes dehydration while lower concentration may be due to the liver not functioning adequately due to factors such as malnutrition and infection (Esubonteng, 2011).

Liver enzymes namely the alanine transaminase (ALT), alkaline phosphatase (ALP) and aspartate transaminase (AST) are important in determining the proper functioning of the liver (Ambrosy *et al.*, 2015). These enzymes are present in negligible concentration and an increase in the concentration of these enzymes may be suggestive of damaged liver cells which denote the status of the liver function. The normal ranges of the concentration of liver enzymes are AST 70-220 μL , ALP 40-129 μL and ALT 7-55 μL in poultry (Makama *et al.*, 2021) and the results obtained were within these reference values. Increased AST is related with the necrosis of the liver cells that can lead to the escape of this enzyme into the blood while ALT is generally increased in situation where there is damage to the liver membranes. The increased blood total bilirubin concentration observed was compatible with other reported results.

Creatinine is used to determine the status of the kidney. The functions of the kidney include excretion of waste products resulting from protein metabolism and muscle contraction (Ileke *et al.*, 2014). Creatinine is excreted by the kidney as a by-product of creatinine phosphate metabolism which is produced as a result of energy production by the skeletal muscles (Esubonteng, 2011). The creatinine level in this study did not fall within the normal reference value of between 0.88-0.95 mg/dl as established by Wikivet (An online Veterinary encyclopedia). The slight increment observed in the level of creatinine is expected in female birds in view of the metabolic changes as a result of sexual maturity and their involvement in egg lay. High Protein intake increases metabolism, stress and dehydration which influence the concentration of uric acid in the blood (Chernecky & Berger, 2008). The urea concentrations obtained were within the normal range of urea/uric acid of 1.9-125mg/dl in poultry (Clinical diagnostic division, 1990). Age, Sex and diets of birds influence the amount of uric acid. A high level of uric acid (hyperuricemia) is usually evident in female birds due to ovulatory activities (Ibrahim *et al.*, 2012).

Cholesterol is synthesized from fats consumed and could also be synthesized endogenously within the cells. A high level of cholesterol is an indication of high risk to cardiovascular disease (Ugbogo *et al.*, 2017). The standard range of values of cholesterol in domestic fowl is between 87-192mmol/L (Meluzzi *et al.*, 1992). The cholesterol level may increase significantly during vitellogenesis and egg formation in birds (Harr, 2002). The increase in the cholesterol level may be due to increased biosynthesis and accumulation in the egg yolk. Notwithstanding, the supplemented layer groups witnessed lower levels of

cholesterol which seemed to be dose-dependent as against G1 that witnessed higher level of cholesterols at 28 and 33 weeks of age. This result was in good agreement with the results of previous researchers that observed that serum cholesterol levels were reduced in birds whose diets were supplemented with activated charcoal (Neuvoneu *et al.*, 1989; Shabani *et al.*, 2010; Dim *et al.*, 2018).

These reductions in the serum cholesterol level could make activated charcoal a product of choice in the management of hypercholesterolemia (Joseph *et al.*, 2015; Roosdiana *et al.*, 2019). It becomes imperative to explore AC as an alternative medication for the treatment and management of hypercholesterolemia in both man and animals. The high availability and low price of AC produced in this study makes it affordable and within the reach of patients with this medical conditions. This coupled with the fact that activated charcoal is non-toxic and safe to be used in oral administration since it is neither absorbed nor metabolized in the GIT (Davis, 2005) becomes an added advantage. Therefore, the activated charcoal produced from co-pyrolysis of precursors (pig dung, oil palm fiber and palm kernel shell) in this study with high surface area and adsorption capacity is suitable for use in cases of hypercholesterolemia.

Conclusion and recommendations

Most of the hematological and biochemical parameters examined were within the patterns often found in avian species meaning that AC is non-toxic and safe to be used in oral administration. Its availability and low price makes it affordable to be used as feed additive for improvement of hematological indices and for patients with hypercholesterolemia to use and keep their serum cholesterol level under control at best inclusion level of 1.0kg/100kg of feeds. Therefore, agricultural waste-derived activated charcoal used in this study is suitable for use in cases of hypercholesterolemia to bind cholesterol and cholesterol-containing bile acids in the gut. It could also serve as a replacement for the synthetic drug used for this condition which is currently very expensive coupled with their long standing side effects which have generated a lot of complaints from patients.

REFERENCES

AAFCO (2012). Official publication. Association of American Feed Control Officials. USA.

Abdel-Hameed, S.A.A., Negin, S.S., Ismael, N.E.M., Soliman, M.M., Shukry, M. & Abdel-Latif, H.M.R. (2021). Effect of activated charcoal on growth, immunity, oxidative stress markers and physiological responses of Nile Tilapia exposed to sub-lethal imidacloprid toxicity. *Animals*, 11 (1357): 1-15.

- Agba, M. M., Ushie, E. F., Abam, I., Agba, M.S. and Okoro, J. (2010). Developing the biofuel industry for effective rural transportation. *European Journal of Scientific Research*, 40: 441-449.
- Akorede, M.F., Ibrahim, O., Amuda, S.A., Otuoze, A.O. and Olufeagba, B. J. (2017). Current status and outlook of renewable energy development in Nigeria. *Nigeria Journal of Technology*, 36(6): 196-212.
- Ambrosy, A.P., Dunn, T.P and Heidenreich, P.A. (2015).Effect of minor liver function test abnormalities and values within the normal range of survival in heart failure.*The American Journal of Cardiology*, 115(7): 938-941.
- Bawala, T. O., Akpan, U., Ogunnowo, A.O., Fusae, O. A. and Sogunle, O. M. (2007). The influence of magnesium supplementation on the hematological profile of young West Africa dwarf goats. *Proceedings of the 32nd conference of the Nigerian Society for Animal Production*, 76 -78.
- Bhatti, S.A., Khan, M.Z., Hassan, Z.U.,Saleemi, M.K., Saquib, M., Khatoon, A. and Akhter, M. (2018). Comparative efficacy of bentonite clay, activated charcoal and Trichosporon mycotoxinovorans in regulating the feed to tissue transfer of mycotoxin. *Journal of the Food Science and Agriculture* 98(3): 884-890
- Boonanuntanasarn, S., Khaomek, P., Pitasong, T. and Hua, Y. (2014). The effects of the supplementation of activated charcoal on growth, health status and fillet composition-odor of Nile Tilapia (*Oreochromis Niloticus*) before and after harvesting. *Aquaculture International*, 22: 1417 – 1436.
- Bounous, D and Stedman, N. (2000). Normal avian hematology: chicken and turkey. In: felman B.F, Zinkl J.G, Jain N.C, editors. Schalm's Veterinary Hematology. New York: Wiley. Pp. 1147 – 1154.
- Campbell, T.W and Coles, E.H. (1986). Avian Clinical Pathology, Fourth edition. W.B Sanders Company Philadelphia. Pp. 279-296.
- Chernecky, C and Berger, B. (2008).Laboratory tests and diagnostic procedures. St. Louis: Saunders Elsevier. *Brazilian Journal of Poultry Science*.
- Coles, E.H. (1986). Avian clinica pathology. W.B Sanders Company, Philadelphia.
- Cooney, D. D. (1980). Activated charcoal: Antidotal and other medical uses. Marcel Dekker Inc. New York, USA.
- Davis. (2005). Atherosclerosis an inflammatory process. *Journal of Insur Medicine*, (37): 72.

- Dawood, M.A.O., Metawally, A.E.S., El-Sharawy, M.E., Ghozlan, A.M., Abdel-Latiff, H.M.R., Van-Doan, H & Ali, M.A.M. (2020). The influences of ferulic acid on the growth performance, haemato-immunological responses, and immune-related genes of Nile Tilapia exposed to heat stress. *Aquaculture*, 525, 735320.
- Dim, C. E., Akuru, E. A., Egom, M. A., Nnajofor, N. W., Ossai, O. K., Ukaigwe, C. G. and Onyimonyi, A. E. (2018). Effect of dietary inclusion of biochar on growth performance, haematology and serum lipid profile of broiler birds. *Agro Science*, 17(2): 8-16.
- Esubonteng O.K.A. (2011). An assessment of the effect of Moringaolifera leaf powder as a nutritional supplement in the diet. *Kumasi Kwame Nkrumah University of Science and Technology*.
- Fischbatch, F.T and Dunning, M.B. (2009). A manual of laboratory and diagnostic tests. Philadelphia Lippincott Williams and Wilkins.
- Gunamantha, I.M. and Widana, G.A.B. (2018). Characterization of the potential of biochar from cow and pig manure for genecology application. *Conference Series: Earth Environmental Science*, 131: 12-55.
- Harr, K.E. (2002). Clinical chemistry of companion avian species: a review. *Veterinary Clinical Pathology*, 31(3): 140-151.
- Huwig, A., Freimund, S., Kappeli, O. and Dutler, H. (2001). Mycotoxin detoxification of animal feeds by different adsorptions. *Toxicology Letters*, 122: 179-188.
- Ibrahim, A., Aliyu, J., Abdu, M and Hassan, A. (2012). Effects of age and sex on serum biochemistry values of turkeys (*Meleagrisgallopardo*) reared in the semi-arid environment of Nigeria. *World Applied Science Journal*, 16(3): 433 – 436.
- Igwe, A.O., Eze, D.C and Nwakuda O.N. (2017). Hematological changes in Isa-brown laying chickens (*Gallus gullusdomesticus*) experimentally infected with velogenic New Castle diseases virus. *Sokoto Journal of Veterinary Sciences*, 15(1):27-35.
- Ileke, K.D., Odeyemi, O.O., Ashamo, M.O and Oboh, G. (2014). Toxicological and histopathological effects of cheese wood, alstoniaboonei de wild stem bark powder used as cowpea protectant against cowpea bruchid, *callosobruchus maculate* (fab.)(Coleoptera: Chrysomelidae) on albino rats. *Ife Journal of Science*, 16(1): 23 – 33.
- Iregbu, G. U., Kubkomawa, I. H., Okoli, C. G., Ogundu, E. C., Uchegbu, M. C. and Okoli, I. C. (2014). Environmental concerns of pig waste production and its potentials as bio-fuel Source. *Journal of Animal and Veterinary Sciences*, (3): 17-24.

- Irizarry-Rovira, A.R. (2004). Avian Hematology. In: Veterinary Clinical Pathology Secrets, Cowell, Rick L, (editor). Saunders Elsevier, Missouri. Pp. 282-305.
- Jiya, E. Z., Anyanwale, B. A., Ijaiya, A. T., Ugochukwu, A. and Tsado, D. (2013). Effect of activated coconut shell charcoal meal on growth performance and nutrient digestibility of broiler chicken. *British Journal of Applied Science and Technology*, 3(2): 268 – 276.
- Joseph, V., Christopher, A.K., James, V.D., Ekta, P and Richard, F.C. (2015). A retrospective review of the prehospital use of activated charcoal. *AM Journal of Emergency Medicine*. 33(1)56 – 59.
- Juul-Madsen, H.R., Viertlboeck, B., Smith, A.L and Gobel T.W.F. (2008). Avian Immunology, Saunders Elsevier, Missouri. Pp. 129-158.
- Kana, J. R., Ngoula, F., Tchoffo, H., Tadondjou, C. D., Sadjo, Y. R., Tegua, A. and Gnonlonfingbemenou, J. B. (2014). Effect of biocharcoals on hematological, serum biochemical and histological parameters in broiler chickens fed aflatoxin B1-contaminated diets. *Journal of Animal Science Advances*, 4(7): 939-948.
- Latimer, K.S and Bienzie, D. (2000). Determination and interpretation of the Avian Leukogram. In: Schalm's Veterinary Hematology. (BF Feldman, JG Zinkl, NC Jain, editors). Lippincott William and Wilkins, Philadelphia, USA. Pp 417-432.
- Majewska, T., Mikulski, D. and Siwik, T. (2009). Silica grit, charcoal and hardwood ash in turkeys. *Journal of Feed Science*, 11: 135 – 141.
- Makama, R.S., Onimisi, P.A. and Afolayan, M. (2021). Apparent nutrient digestibility, liver function indices and lipid profile of broiler chickens fed raw and boiled sickle pod (*Senna obtusifolia*) seed meal. *Nigerian Journal of Animal Science*, 23(2): 207-212.
- Maklad, A. I., Ashraf, M. E., Enas, I. E. and Mennat-Allah, A. M. E. (2012). Pediatric poisoning in Egypt. *Journal of Poultry Elem*, 14: 489-500.
- Meluzzi, A., Primiceri, G., Giordani, R and Fabris, G. (1992). Determination of blood constituents reference values in broilers. *Poultry Science*, 71(2): 337 – 345.
- Mohammed, Y. S., Mustafa, M.W., Bashir, N. and Mokhtar A. S. (2013). Renewable energy resources for distributed power generation in Nigeria: A review of the potential. *Renewable and Sustainable Energy Reviews*, 22: 257-268.
- Moyes, C.D and Schute, P.M (2008). Principles of Animal Physiology. 2nd Edition. Pearson International Edition. New York.

- Neuvonen, P.J., Kuusisto, P., Vapaatalo, H. and Manninen, V. (1989). Activated charcoal in the treatment of hypercholerolaemia: Dose- response relationships and comparison with cholestyramine. *European Journal of Chemical Pharmacology*, 37: 225-230.
- Oyedepo, S. O. (2012). Energy and sustainable development in Nigeria: The way forward. *Energy, Sustainability and Society*, 2: 14.
- Roosdiana, A., Vidiastuti, D and Herenda, H. (2019).The preventive effect of activated charcoal on HDL levels and aorta histopathological profiles in hypecholesterol rat models. *Journal of Physics: Conference Series*, (1374) 012029.
- Samour, J. (2009). Diagnostic value of hematology. In: Clinical Avian Medicine: (GJ Harrison, TL Lightfoot, (editors). www.clinicalavainmedicine.com.
- Schalm, O.W., Jain, N.C and Carrol, E.J. (1975). Veterinary Haematology, 3rd Edition, Lea and Febiger, Philadelphia, Pp. 197-199.
- Schmidt, H.P., Hagemann, N., Draper, K. and Kammann, C. (2019).The of biochar in animal feeding. *Peer Journal*, 7: e7373.
- Shabani, A., Dastar, B., Khomeiri, M., Shabanpour, B. and Hassani, S. (2010). Response of broiler chickens to different levels of nanozeolite during experimental aflatoxicosis. *Journal of Biological Science*, 10(4): 362-367.
- Sharma, S.L., Singh, P., Patil, A.K and Sharma, J (2015). Effect of feeding compressed complete feed block containing sugar meal on blood biochemical profile of crossbred calves. *Journal of Animal Research*, 5(3): 575.
- Sufiriyanto, N.I and Emmy, S. (2018). Hematological profiles and performance of broiler chickens fed on commercial feed. *Journal of Animal Production*, 20(3): 183 – 190.
- Ugbogo, A.E., Okezie, E and Ijioma S.N. (2017). Introducing Biochemistry practical by Justman Publishers International, Pp. 143 – 202.
- Ugochukwu, E.I (2001). Laboratoryexamination in veterinary clinical diagnosis. In: Ugochukwu, E.I (ed). Manual of Veterinary Clinical Practice.Amazing Grace Printing and publishing Co. Nigeria. Pp. 174 – 194.