

Original Research Article

Severe malaria in adults at the Yalgado Ouédraogo University Hospital of Ouagadougou, Burkina Faso: epidemiological, clinical, therapeutic and evolutionary aspects

Abstract

Aims: To describe the epidemiological, clinical, therapeutic and evolutionary characteristics of severe malaria in adults at the Yalgado Ouédraogo University Hospital from 2004 to 2021.

Methodology: This was a cross-sectional study, with retrospective data collection covering the period from January 1, 2004 to December 31, 2021. Patients with severe malaria diagnosed on the basis of clinical and biological signs, with a positive thick blood drop were included.

Results: During the study period, a total of 217 cases of severe malaria were recorded, representing a hospital prevalence of 2.1%. The mean age of the patients was 40 years \pm 20 with extremes ranging from 18 years to 92 years. Females predominated (52.8%) compared to 47.2% for males, i.e. a sex ratio of 0.89. Thirty percent of the patients lived in urban areas, 24% in semi-urban areas and 46% in rural areas. They were mostly housewives, farmers, pupils or students. Clinically, the functional signs of our patients were dominated by neurological signs, digestive signs and respiratory signs. The physical examination noted a systemic inflammatory response syndrome and multivisceral failure in the majority of patients. Biological disturbances were dominated by renal failure, severe anemia, thrombocytopenia and metabolic acidosis. Comorbidities were dominated by arterial hypertension, HIV infection and diabetes. Under treatment, the evolution was marked by a case fatality of 29.5%.

Conclusion: In clinical practice, severe malaria in adults is rare but potentially serious. It is frequently responsible for sepsis associated with multivisceral failure. Its prognosis depends on the rapidity of the diagnosis and the initiation of appropriate treatment.

Key words: severe malaria, adult, sepsis, multivisceral failure.

1 Introduction.

Malaria is a vector-borne parasitic infection that has been prevalent since ancient times [1]. Nearly half of the world's population is at risk of malaria [2]. Despite a global funding effort through the Global Fund, the US President's Malaria Initiative, the Roll Back Malaria program, and the Abuja Declaration [2,3], malaria remains the most common parasitic endemic in the world. It is a priority public health problem, particularly in sub-Saharan Africa [4,5]. In Burkina Faso, malaria remains the leading cause of consultations in health facilities and the World Health Organization (WHO) estimates that the country recorded 3.2% of global cases in 2020 [6,7,8]. This is why Burkina Faso has subscribed to the global strategy to fight this scourge [2,4]. The COVID-19 pandemic that occurred in 2019 has had a negative impact on the fight against this disease [8]. Indeed, WHO estimates that in 2020, there would have been 14 million cases of malaria with 69,000 more deaths compared to 2019 [8]. Ninety-three percent (93%) of malaria cases and 96% of deaths occurred in the WHO African Region [8,9]. Most deaths were due to severe forms of *Plasmodium falciparum*. However, severe forms due to *P vivax* and *P Knowlesi* have also been reported to a lesser extent [10]. Several authors have reported the high lethality of severe forms of malaria, particularly in children, who pay a heavy price for this disease [11,12,13]. The development of the antimalarial vaccine, RTS, S/AS01, is a ray of hope since it would reduce severe forms of malaria in children by 30% [9]. Although children are more exposed to this disease, adult cases are also encountered in routine practice. The objective of this study is to describe the epidemiological, clinical, therapeutic and evolutionary characteristics of severe malaria in adults at the Yalgado Ouédraogo University Hospital from 2004 to 2021.

2 Patients and method.

This study involved all adult patients hospitalized for severe malaria in the medical emergency services and in the infectious diseases service of the CHU-YO from January 1, 2004 to December 31, 2021.

This was a cross-sectional study, with retrospective collection from January 1, 2004 to December 2021. Patients with severe malaria diagnosed on the basis of clinical and biological signs with a positive thick blood drop were included. The definition of severe malaria was the WHO definition[24].

Patients with severe malaria but whose records were not usable were excluded. For each patient included, socio-demographic, clinical, therapeutic and evolutionary variables were collected.

Data were entered, processed, and analyzed on a microcomputer using EPI info software version 7.2.2.6. Chi-square tests were used for statistical comparisons.

From an ethical point of view, the confidentiality and anonymity of the patients were respected in the processing and analysis of the data.

No information about test of biological signs, please include it with material and method

3 RESULTS AND DISCUSSION

3 1 RESULTS

Epidemiological aspects

During the study period, a total of 217 cases of severe adult malaria were collected, representing a hospital prevalence of 2.1%. The mean age of the patients was 40 years \pm 20 with extremes ranging from 18 years to 92 years. Females predominated (52.8%) compared to 47.2% for males, i.e. a sex ratio of 0.89. Thirty percent (30%) of the patients lived in urban areas, 24% in semi-urban areas, while the majority (46%) lived in rural areas. Table I shows the distribution of patients by occupation.

Table 1: Distribution of patients by occupation

Occupations	Number of patients	Percentage rate%
Housewives	64	29,5
Farmers	53	24,5
Pupils / Students	52	23,9
Workers	20	9,2
Shopkeepers	17	7,8
Civil servants	11	5,1
Total	217	2.1%.

Clinical aspects

The average time to consultation for our patients was 5 days \pm 3. The medical history is presented in Table II. (2)

Table 2: Distribution of patients by medical history

Background	Number of patients	% percentage rate
HTA	23	10,5
Diabetes	13	5,9
HIV	13	5,9
Sickle cell disease	6	2,7
STROKE	6	2,7
Total	217 ?	

Table III shows the distribution of patients according to functional signs.

Table 3: Distribution of patients according to functional signs

Functional signs	Number of patients	Percentage %
Headache	67	31,15
Vomiting	57	26,4
Disturbances of consciousness	53	24,4
Abdominal pain	41	19,05
Asthenia	37	17,1
Delirium	27	12,4
Anemia	25	11,7
Anorexia	21	9,5
Prostration	17	8,6
Coughing	8	3,8
Chills	8	3,8
Dizziness	8	3,8
Diffuse pains	8	3,8
Total	217 ?377 ?	?

NB: a patient could have several functional signs.

The clinical and biological signs presented by the patients are presented in Table IV.

Table 4: distribution of patients according to clinical and biological signs

Clinical signs	Number	Percentage %
Fever	155	71,4
Tachycardia	111	51,2
Coma	50	23,15
Hypotension	39	18
Respiratory distress	37	17,14
Altered general condition	35	16%
Hemoglobinuria	27	12,4
Septic shock	27	12,4
Delirium	27	12,4
Jaundice	25	11,66
Convulsions	21	9,7
Hemorrhage	20	9
Agitation	17	7,85
Biological signs		
Creatinemia greater than 265µmol/l	87	40,31
Severe anemia	85	39,25
Thrombocytopenia	84	38,5

Metabolic acidosis	70	32,36
Peripheral oxygen saturation<90%.	39	18
Hypoglycemia	18	8,15
Total	Its should be 217 ?	

Therapeutic aspects

Ninety-four percent (94%) of patients received artesunate injection with oral artemether + lumefantrine. Eight percent (8%) of patients had received quinine, and 1% of patients had received artemether. All patients had received adjuvant therapy: 180 patients (83%) had received analgesic and antipyretic treatment; 73 patients (33.6%) had been rehydrated; 41 patients (19%) had received oxygen therapy; 39 patients (18%) had received anticonvulsant treatment with injectable diazepam or phenobarbital; 74 patients (34%) had received electrolyte supplementation; 116 patients (53.4%) had received antibiotic therapy; six patients (2.8%) had received resuscitation; 21 patients (9.7%) had received 10% hypertonic glucose serum; 70 patients (32.3%) had received a blood transfusion based on red blood cell concentrate.

The duration of hospitalization was less than seven days in 78% of patients. It was more than seven days in 22%. The evolution was marked by a lethality of 29.5%.

Please your total patients is 217

Diagnosis of malaria according to what you have mentioned in material and methods as well as in results(methods either serological, blood smear,,,,,,,,,,,,,?????)

Species of malaria are same in all studied area?????

3 2 Discussion

In our series, as in that of Yameogo, there was a predominance of the female sex, contrary to most African series [5,11,12,14,26]. The mean age of our patients was higher than in most African series [5,11,13,14,26]. As in the Eholié and Soumaré series [12,26], the functional signs of our patients were dominated by headache. On physical examination of our patients, neurological signs predominated, while biologically, disturbances of renal function were frequently associated in our patients. The same observation had been made by Senegalese and Ivorian authors [5,12,15]. The mechanism of renal failure would be related to vascular obstruction by parasitized erythrocytes responsible for tubular necrosis [21]. The clinical and biological signs observed in our patients often evolved in a context of sepsis with multivisceral failure, severe anemia, thrombocytopenia and acidosis, all of which are factors that worsen the prognosis of the patients [6, 16, 17, 18]. Bruneel made the same observation in patients with severe imported malaria [19]. And he suggested that these patients should benefit from special and urgent management in intensive care [19,20]. Therapeutically, severe malaria is treated with intravenous artesunate according to WHO recommendations [22]. As soon as the patient is clinically stable and able to swallow, oral treatment with artemisinin derivatives should be administered [22]. The management of severe malaria must necessarily take into account the different disturbances that are associated [23,24,25]. The evolution was marked by a high case fatality in our series compared to most African series [5,11,12,15,26]. We agree with Eholié that despite the good efficacy of the antimalarial drugs used, the prognosis of severe malaria in adults remains severe. However, the prognosis could be improved by early diagnosis and appropriate management. To achieve this, adequate equipment of health care facilities is necessary [12].

There is variation between clinical signs of your study compared to others why?

The physical examination of patients, neurological signs predominated, while biologically, disturbances of renal function were frequently associated in patients is varied to other study ?

What about male and female compared it with other study?

What about age in your study compared to other clarified it please also??

Conclusion

Severe malaria in adults is rare in routine practice. It should be considered and a thick blood sample should be requested for any adult patient presenting with sepsis. The relative frequency of septic shock states during severe malaria requires the setting up and consequent equipment of intensive care and resuscitation units in our hospitals. Particular emphasis should be placed on prevention through vector control, the use of insecticide-treated mosquito nets, rapid diagnosis and treatment of simple forms of malaria, both in health facilities and in the community, and chemoprophylaxis in subjects at risk.

Références

- 1) Argy N., Houzé S.. Severe malaria: from pathophysiology to therapeutic novelties. *Journal of Anti-infectives* 2014, 16(1): 13-17
- 2) WHO. World malaria report 2020. 20 years of global progress and challenges. *World-malaria-report-2020-briefing-kit-eng.pdf*
- 3) WHO calls for a renewed fight against malaria. WHO calls for renewed fight against malaria (who.int)
- 4) WHO 2015. Global Technical Strategy for Malaria Control 2016-2030.. <https://www.who.int/fr/publications-detail/9789240031357>
- 5) Diop SA, Ndour CT, Dia MN, Manga MN et al. Current aspects of severe malaria in adults at the infectious diseases clinic of the CHUN of Fann of Dakar(Senegal). *Med Afr Noire* 2010, 57, 193-197.
- 6) Bruno Pradines, Marie-Gladys Robert. Situation of malaria in the world. *La Revue du Praticien*, 2019 69(2) :146-9
- 7) Ministry of Health Burkina Faso. National directive for the management of malaria in Burkina Faso, National Malaria Control Program, 2017 p12-8.
- 8) WHO. World Malaria Report 2021. [WHO. World Malaria Report 2021 - Search \(bing.com\)](#)
- 9) WHO. Malaria Report 2020. www.who.int/fr/news-room/fact-sheets/detail/malaria
- 10) Adoumbaye Noubadoum, Diouf JB, Leye MY, Faye MA, Siby CT, Ndong SB, SENE PA, Kamwa DG, Lo G, Taha IA and Sani DRE. Severe oval Plasmodium malaria in adults: a case report and review of the literature. *J Biol Today's World* 2020; 9(5): 222
- 11) Soumaré M., Seydi M., Diop S.A., Diop B.M. & Sow P.S.. Adult neuromalaria at the infectious diseases clinic of the CHU de Fann in Dakar, Senegal. *Bull Soc Pathol Exot*, 2008, 101, 1, 20-21
- 12) Eholié SP, Ehui E, Adou-Bryn K, Kouamé Ke, Tanon A et al - Severe malaria in indigenous adults in Abidjan, Ivory Coast. *Bull Soc Pathol Exot*, 2004, 97, 340-344.
- 13) Diallo A. H., Guiguemde T. R. & G. Ki-Zerbo. *Bull Soc Pathol Exot*, 2003, 96, 2, 99-100
- 14) Yaméogo T. M., Kyelem C. G., Ouédraogo S. M., Diallo O. J., Moyenga L., Poda G. E. A., Guiguemdé T. R.. Diagnosis and management of severe malaria in adults: compliance with national guidelines in Burkina Faso. *Bull Soc Pathol Exot*, 104, 284-287.
Year ???
- 15) Lakhe N A, Sylla K, Diallo Mbaye K, Dia Badiane N M, Jaber C, Massaly A, Cisse Diallo V M P, Ka D, Fortes Deguenonvo L, Ndour C T, Soumare M et Seydi M. Aspect actuel du paludisme grave au service des maladies infectieuses et tropicales de Fann (Dakar). *Archives of Infect Diseases & Therapy*, 2018 ; 2 (2) :4 - 7.

- 16) Savadogo M., Sondo K A., Thiombiano R. Sangaré R., Drabo YJ. Ouédraogo A., clinical and therapeutic epidemiological aspects of malaria at the CHU Yalgado Ouédraogo of Ouagadougou, Burkina Faso. *Med d'Afr. Noire.* 2014, 61, 374-378.
- 17) Rabo A. Study of anemia in the infectious diseases department of the Yalgado Ouédraogo University Hospital (Burkina Faso). Med thesis n°65, Ouagadougou, 2021.
- 18) Ouedraogo G. B A. Comparative study of clinical and biological manifestations of severe malaria in children versus adults from July 1 to September 30, 2021 at CHU-YO. Med thesis n°407, Ouagadougou, 2021.
- 19) Bruneel F. Raffetin - A. - Roujansky A. - Corne P. - Tridon C. - Llitjos J.-F. - B. Mourvillier - V. Laurent -S. Jauréguiberry. Management of severe imported malaria in adults. *Méd. Intensive Réa* (2018) 27:228-238
- 20) Kagone A. Severe malaria in adults at the chu Yalgado Ouédraogo: epidemiological, clinical, therapeutic and evolutionary aspects from 1 November 2019 to 31 October 2020. Med thesis n° 452, Ouagadougou, 2020.
- 21) Kissou S A , R Cessouma, M Barro, H Traoré, B Nacro. Acute renal failure and Plasmodium falciparum malaria: a case report. *Arch Pediatr.* 2012 Jan;19(1):34-7.
- 22) Trampuz A, Jereb M, Muzlovic I, Prabhu R M. Clinical examination: severe malaria. *Care Crit.* 2003 Aug;7(4):315-23.
- 23) World Health Organization, Severe malaria. *Trop Med Int Health* (2014) 19: 7–131
- 24) WHO. MANAGEMENT OF SEVERE MALARIA, A Practical Guide to the Management of Severe Malaria 3rd Edition 2013; <https://www.who.int/fr/publications-detail/9789241548526>
- 25) Marques M , Un Gupta-Wright , J F Doherty , M Chanteur , D Walker. Management of malaria in the intensive care unit. *F J Anaesth* 2014 Dec;113(6):910-21.
- 26) Soumare M, Seydi M, Diop S A, Ba TA et al. Place of malaria in a department of infectious pathologies in Dakar (Senegal). *Med TROP* 2008; 68: 485-490.