

## Original Research Article

### EXPLORING EVIDENCE FOR DISRESPECT AND ABUSE IN FACILITY-BASED CHILDBIRTH AT A TERTIARY FACILITY IN GHANA

#### ABSTRACT

**Background:** Pregnant women's decisions to seek maternity care are influenced by the poor quality of health care they receive, as well as their fears of being disrespected and abused during childbirth, which are perpetuated by health workers. Physical abuse, non-consented care, non-confidential care, non-dignified care, discrimination, abandonment, and detention in facilities are some of the most common manifestations of disrespect and abuse. Maternal mortality interventions have primarily targeted facility-based delivery, but the proportion of births in facilities has remained persistently low in many low-resource settings, despite the fact that facility-based birth rates have increased in recent years.

**Purpose:** The purpose of this study was therefore to explore women's views on disrespect and abuse in childbirth in the Tamale Teaching Hospital.

**Methodology:** Analytical cross-sectional design was used to collect data for this study, and a purposive sampling technique was used to select the respondents from the study area. A total of two hundred and forty respondents were surveyed for the purpose of gathering information for the study. The majority of the primary data was gathered through the use of administered questionnaires. The information gathered was analyzed with the help of Microsoft Word Excel.

**Key findings:** Results from the study revealed that, the vast majority (23 percent) of respondents identified insulting as a form of abuse and disrespect during childbirth, and that the vast majority (91%) of respondents believed that abuse and disrespect during childbirth could never be stopped. A further finding was that the vast majority of respondents (86%) viewed abuse and disrespect during childbirth as negative, whereas 51.6% of respondents viewed the absence of abuse as a means of improving the overall quality of health care during childbirth.

**Recommendations:** On the basis of the study's findings, a number of recommendations were also made, including the intensification of interventions that could help prevent neglect or isolation among new mothers in the study area.

**Key Words:** *Disrespect, Abuse, Childbirth, Maternity Care, Facility-Based Birth, Interventions and Tamale Teaching Hospital*

## BACKGROUND

The relationship that a woman has with her maternity care providers is critical to her overall well-being (Berry, 2010). Furthermore, not only do these encounters serve as a conduit for essential lifesaving health services, but women's interactions with caregivers can either empower and comfort them or cause them long-term harm and emotional trauma (Kruk, 2009). However, women's memories of their childbearing experiences last a lifetime and are frequently shared with other women, contributing to a climate of confidence or doubt surrounding childbearing and health-care utilization (White Ribbon Alliance, Respectful Maternity Care, 2011). Obstetrics and obstetrical care continue to put women at risk for significant morbidity and mortality, particularly in sub-Saharan Africa (Baer, Singer & Johnsen, 2010).

Pregnant women require access to the infrastructure, equipment, and personnel necessary for routine and emergency obstetric and newborn care in order to exercise their right to readily available, easily accessible, acceptable, and of high-quality care (Thaddeus & Maine, 2009). Despite the advancements, many pregnant women, particularly in developing countries, continue to face challenges in obtaining high-quality services (Watson-Jones, 2012). Although services are available, care may be compromised by social, ethnic, and cultural barriers; a chilly reception at the health care facility; a lack of privacy and limited information sharing between the client and the provider; and, in some cases, outright neglect or abuse (Bowser & Hill, 2010). The right to health establishes the broadest definition of disrespect and abuse, which is the most common type of abuse.

Those interactions or facility conditions that are perceived as or intended to be humiliating or undignified by the local community, as well as those interactions or facility conditions that are experienced as or intended to be humiliating or undignified by the local community (WHO, 2008). A landmark review of evidence for disrespect and abuse in facility-based childbirth conducted by Bowser and Hill (2010) identified numerous examples, including physical abuse, non-dignified care, nonconsented care, non-confidential care, discrimination, abandonment of care, and detention in facilities. They observed that women avoid seeking care in health facilities due to mistreatment, which jeopardizes the achievement of **Millennium Development Goal (MDG) 5** - to improve maternal health - by the international community.

**Comment [MA1]:** The background section is too long, several phrases can be submitted in these details in the discussion section, please keep focusing to highlight the core objective and the target of the study!

**Comment [MA2]:** For the journal readers please submit a reference! To advance the health of women and children, gender equality, education, the environment and global partnerships. re write the full perspective of this goal, (MDG 8) !

According to international law, the right to health necessitates the provision of health services that are readily available, easily accessible, acceptable, and of high quality (Rosenfield, 2010). While numerous official interpretations and guidance documents have been issued in support of the application of this right to childbirth, reports of disrespectful or abusive treatment during labor and delivery continue to appear in many parts of the world (Rosenfield, 2010). Recently published data from low-resource settings revealed a widespread practice of disrespectful and abusive treatment by health-care workers when pregnant women seek childbirth, as well as the ability of such experiences to undermine access to childbirth care during subsequent deliveries (Bowser, 2010).

The World Health Organization (WHO) recently issued a call for action to put an end to disrespectful behavior during skilled birth, from policy to practice (d'Oliveira, 2010). According to the call, women from low socioeconomic status, migrant women, and women from ethnic minorities are particularly vulnerable to disrespect and abuse during pregnancy and childbirth (d'Oliveira, 2010). According to the growing body of anecdotal and research evidence collected in maternity care systems around the world - from the wealthiest to the poorest countries - it appears that disrespect and abuse of women seeking maternity care are becoming increasingly serious issues (Kruk, Mbaruku, de Pinho & Galea, 2009). Taking a stand against disrespect and abuse, the World Health Organization (WHO) states that every woman has the right to the highest attainable standard of health, including the right to dignified and respectful care during pregnancy and childbirth (Bowser & Hill, 2010).

## MATERIALS AND METHODS

Study design: The study employed a cross-sectional design that was primarily analytical in nature. This design was chosen due to the nature of the research topic, as well as the fact that it is relatively inexpensive and provides the opportunity to evaluate both exposures and outcomes at the same time. As described by Avoke (2005), analytical cross-sectional study designs are intended to accurately portray the characteristics of specific individuals, situations, or groups. In order to provide information on which to base sound decisions and to prepare the groundwork

**Comment [MA3]:** Please explain the study perspective in the discussion section! Why we applied this type of analysis!

for a more constructive program of educational research, it is used as a needs assessment tool (Avoke, 2005).

Setting: The study was conducted in the Tamale Teaching Hospital which is located in Tamale; the regional capital of the Northern region of Ghana. The Hospital was established to serve as a medical referral center for the Northern, Upper East and Upper West Regions, the northern parts of the Brong-Ahafo Region, Oti Region and the neighbouring countries of Ivory Coast, Burkina Faso and Togo (Annual Report of TTH, 2018).

Target Population: The study participants were primarily post-partum women who had been in the Postnatal Ward of the Tamale Teaching Hospital for no more than three days at the time of the study's implementation. The study population consisted of women of reproductive age (15 to 49 years) who had given birth in the hospital within the previous (0-3) days and who had been attended by a skilled birth attendant during their delivery.

Inclusion Criteria: Participants were included in the study if they were 15-49 years, who had given birth.

Exclusion Criteria: Participants who does not fall within the reproductive age.

Sampling Technique and Size: The respondents for the study were selected using a technique known as purposive sampling. The Fisher's method was used to narrow down the pool of potential respondents. The numbers used as identification were given to post-partum women who had delivered between (0-3) days prior and were written on pieces of paper that were then folded in half. Using a clean rubber bucket, the folded papers were thoroughly mixed and kept together. To obtain the required sample size of 240 folded papers, the technique of sampling without replacement was used to pick the folded papers out of the rubber bucket one by one until the required sample size of 240 was obtained.

Data Collection Instrument: For the purpose of gathering information, a structured questionnaire consisting of both closed- and open-ended questions was used as the primary tool. Primary and secondary data sources were used to gather the information. When collecting primary data, the researchers used structured questionnaires to gather as much information as they possibly could. Individually administered questionnaires were given out by the researchers at the Tamale

Comment [MA4]: No need to submit such details !

Comment [MA5]: You mentioned that previously , please remove!

Comment [MA6]: This is the target of the study you already mentioned! No need to re submit that again!

Comment [MA7]: This section should be before the sample technique and size!

Teaching Hospital's Post-Natal Ward. Obtaining secondary information was accomplished through a review of related literature, which included books, journals, and internet articles.

**Data Analysis:** The completed questionnaires were crossed checked for completeness and accuracy. The questions were coded and entered into the excel computer software for analysis using Microsoft Excel Word 2018. The descriptive tools were tables, pie chart, graph and bar charts together with relevant and narrative expression.

**Comment [MA8]:** As I see the authors re write these phrase with an acceptable repetitive way to the journal reader, please you can prescribe that under the Sampling Technique and Size:

The researchers conducted the data collection in order to ensure that the information gathered was trustworthy and valid. This ensured that the questions were consistent throughout. The questionnaires were also pre-tested on a total of fifteen post-partum women before being distributed. The pre-testing was carried out in order to assist in the restructuring of the questionnaire. After double-entry of data was completed, the two data sets were compared at the conclusion of the analysis.

**Ethical Consideration:** Before beginning the research, permission was sought from the hospital administration at the Tamale Teaching Hospital. This was granted. Before they were administered, the questionnaires were also reviewed by the appropriate authorities. We sought respondents' informed consent because the issues of disrespect and abuse during childbirth are extremely sensitive. The objectives and significance of the study were clearly stated and explained to the potential respondents in the consent form, which they signed. Individuals who chose not to participate in the study were given the opportunity to do so at their discretion. In order to maintain the anonymity and confidentiality of the actual source (s) of information obtained from the study, the names of those who took part in the study were withheld from publication. Because no names were provided on the data collection tools, there were no clues left for someone to follow in order to determine the source of the information.

**Comment [MA9]:** Please remove!

**Comment [MA10]:** Again no need to re write the name of the institute!

**Comment [MA11]:** Written informed consent was obtained from the participants ( respondents), it is to long for the journal audiences !

## **STUDY FINDINGS**

This chapter presents the results from the data that was collected from the respondents. The data was entered and analyzed by Microsoft word excel 2018. Data is presented according to sections.

**Comment [MA12]:** Results !

The information is displayed in the form of tables and charts for easy description and interpretations.


Comment [MA13]: The authors should mention that in the methods section!

## SOCIO-DEMOGRAPHIC CHARACTERISTICS OF RESPONDENTS

Figure 1.1 below shows the age distributions of respondents who formed part of the study. Most of the respondents representing 52% were within the age of 31-35 years, 19% respondents were within 26-30 years, 8% respondents were within 20-25 years whiles 21% respondents were 36 and above years.

Comment [MA14]: The authors should highlight the number or percentage either in the results section or just to keep the figures represent the ideas, no place to submit double same data !

Figure 1.1: Age of respondents in years




From Figure 1.2 shown below majority of the respondents (90%) were married whiles (10%) respondents were single or in consented relationship none of the respondents however stated divorce, separated or widowed.

Comment [MA15]: The authors should consider our previous comment!

Figure 1.2: Marital status of respondents


### Marital status of respondents



From Figure 1.3 shown below, most of the respondents (62.5%) were practicing Islam with 37.5% respondents practicing Christianity. None of the respondent was practicing other religions at the time of the research

**Comment [MA16]:** The authors should consider our previous comment!

**Figure 1.3: Religious status of respondents**



From Table 1. indicated below, majority of the respondents (66.7%) were self-employed, 14.2% respondents said they were students whiles 8.3% respondents stated they were unemployment. 10.8% respondents were gainfully employed hence salaried workers.


**Table 1: Occupations of respondents**

VARIABLE	FREQUENCY	PERCENT
Students	34	14.2
Self-employed	160	66.7
Gainfully employed	26	10.8
Unemployed	20	8.3
<b>TOTAL</b>	<b>240</b>	<b>100.0</b>

From Figure 1.4 shown below, most of the respondents 144 (60%) were Dagombas, 42 (17%) respondents were Gonja, 28 (12%) respondents were Dagarti, 8 (3%) respondents were Akans and 8 (18%) of the respondents were Bimoba

Comment [MA17]: The authors should consider our previous comment!


Figure 1.4: Ethnicity of respondents




From Figure 1.5 indicated below, majority (118) of the respondents had no formal education with 28 and 34 respondents having primary and secondary education respectively. 40 respondents had tertiary education whiles 29 said they had vocational education.

Comment [MA18]: The authors should consider our previous comment!

**Figure 1.5: Educational status of respondents**



**Figure 1.6: Parity of respondents**



From Figure 1.6 shown below, majority of the respondents (128) had 3-4 children, 62 respondents had 5 and above children while 50 respondents had 1-2 children at the time of conducting the research at the study area.

**Comment [MA19]:** The authors should consider our previous comment!


From Table 2 presented below, 52.5% respondents had conceived for three and above times, 39.2% respondents conceived for the second time while 8.3% respondents however were primigravidae

**Table 2: Number of times of pregnancy of respondents**

VARIABLE	FREQUENCY	PERCENT
First	20	8.3
Two	94	39.2
3 and above	126	52.5
<b>TOTAL</b>	<b>240</b>	<b>100.0</b>

From Figure 1.7 illustrated below, 91% respondents' mode of delivery was Spontaneous Vagina Delivery (SVD) while 9% respondents said by Caesarian Section (C/S).

**Figure 1.7: Mode of delivery of respondents**



#### **FORMS OF ABUSE AND DISRESPECT BY RESPONDENTS**

Conversely, a definition of disrespect and abuse based during childbirth on the actual experience of violations from the perspectives of post-partum women and health care providers is difficult to coin, especially when aspects of disrespect and abuse during child birth are so common among providers or so expected by post-partum women as to be normalized in the health system. The study sought to find a definition for disrespect and abuse or what respondents termed as disrespect and abuse in their desire to deliver in the hospital.

From the responses gathered from the respondents, it was evident that there is no single definite definition for disrespect and abuse by post-partum women. Some direct quotes from respondents on what abuse and disrespect in child birth are is; "shouting at pregnant women by midwives and nurses during labour", "maltreating a pregnant women during childbirth", "is the way a pregnant woman is received and handle during childbirth", "insulting and shouting at a pregnant woman


whiles in labour”, “it is making a pregnant women in labour uncomfortable”, “giving pregnant women emotional pain during labour by making unnecessary comments likes; shout up and push”.

**Comment [MA20]:** No place for such comments in the result section!

From the Figure 1.8 indicated below, majority of the respondents representing 43% said they heard of abuse and disrespect during childbirth from the radio while 21% respondents mentioned others like; family members, friends, workshops attended and personal experiences. From the Figure also, 6% respondents said they heard of abuse and disrespect during child birth from the social media and 25% respondents stated clinic with 5% respondents identifying School.

**Comment [MA21]:** The authors should highlight the number or percentage either in the results section or just to keep the figures represent the ideas, no place to submit double same data!

**Figure 1.8 : Sources of respondent’s knowledge of abuse and disrespect**



Majority (23%) of the respondents identified insulting as a form of abuse and disrespect during childbirth, 21% respondents mentioned non-dignified care, 8% respondents stated non-


confidential care with 6% respondents stating physical beating. 2% respondents and 40% respondents respectively identified buying of things and shouting as abuse and disrespect during childbirth.

All the respondents (100%) identified the midwives and nurses as the most frequent health workers who abused pregnant women during child birth probably because they are directly involved in the art and process of child birth. All the respondents (100%) stated that the negative effect of abuse and disrespect during childbirth is the emotional instability they usually experienced

From Figure 1.9 illustrated below, majority of the respondents (91%) were of the opinion that abuse and disrespect during childbirth can never be stopped whiles 9% respondents expressed a contrary view.

**Comment [MA22]:** Again either comments or let the figure present itself, we found several repetitive phrases may make an illusion for the journal readers!

**Figure 1. 9: Respondents opinion on abuse and disrespect**



## PERCEPTION AND VIEWS ON ABUSE AND DISRESPECT IN CHILDBIRTH


All pregnant women who report negative experiences in the health facilities either during pregnancy or during previous births often decide to give birth at home for fear of humiliation, bad treatment and for not being respected by health care providers.

**Comment [MA23]:** These phrases already submitted in the background section and even in methods section, again no need for the repetitive phrases please!

From Figure 1.10 presented below, majority of the respondents (86%) perceived abuse and disrespect during childbirth as bad while 14% respondents perceived it not to be bad. Among the latter, they mentioned factors such as; some pregnant women are always too reluctant to comply to basic issues during labour and during labour most pregnant women become emotional unstable and unnecessary angrily and need to be controlled.

**Figure 1.10: Respondent's perception of abuse and disrespect during childbirth**


**Comment [MA24]:** What sort of the ideas the authors highlighted in this figure and the significance of finding from the previous one!



Source: Field Survey, 2015

From the responses, most of the respondents have had one form of abuse or disrespect before or otherwise. As indicated in Figure 1.11 below, majority (94.1%) of the respondents said they have ever been abused before while 5.9% said they have never been abused before.

Figure 1.11: Respondents and abuse during childbirth



Comment [MA25]: What sort of the ideas the authors highlighted in this figure and the significance of finding from the previous one!

Source: Field Survey, 2015

From Table 3 illustrated below, 51.6% respondents perceived absent of abuse as a way of enhancing quality of health care during child birth, 11.3% respondents mentioned friendly care while 37.1% respondents stated proper communication by health care providers.

Table3: Respondents perception of quality of health care

VARIABLE	FREQUENCY	PERCENT
proper communication	89	37.1

Comment [MA26]: I think the tables can play a good roles and sure the authors can presented that with core ideas to the journal readers, the results section submitted with a very long phrases and it may reflect un illusive images for the audiences!


Friendly care	27	11.3
Absent of abuse	124	51.6
<b>TOTAL</b>	<b>240</b>	<b>100</b>

**Comment [MA27]:** The authors did not submitted the classical significance of the results by the statistical point (P value)! Despite you mentioned the fisher tests!

**Source: Field Survey, 2015**

From Figure 1.12 presented below, 72.3% respondents said educational level of pregnant women determines whether the person was going to be abused or disrespected while 27.7% respondents however, stated that educational level was not linked to abuse and disrespect during childbirth

**Figure 1.12: Education and abuse during child birth**




**Source: Field Survey, 2015**

From Figure 1.13 indicated below, majority (84.4%) of the respondents believed that the study will improve child birth in the Hospital while 15.6% respondents expressed a different view. 56% respondents said financial status of pregnant women is a contributory factor to abuse and disrespect during child birth as poor pregnant women were more likely to suffer abuse than the rich while 44% respondents said there was no relationship between abuse and financial status of pregnant women during child birth.

76% respondents strongly agreed that lack of policies that support respectful and non-abusive maternity care are contributory factors to abuse and disrespect during child birth while 15% respondents agree to that assertion 9% respondents however disagree. Eight-seven percent respondents strongly agree that lack of standard and leadership for respect and non-abusive in child birth are contributory factors to abuse and disrespect while 13% respondents disagree.

**Figure 1.13: Respondents assessment of the study on the impact of childbirth**

**Comment [MA28]:** The study itself can build a programme to improve the child birth when the authors can analyse the data in a precise qualification and usage of the appropriate statistics to pick up the final target and claim the objectives.



Source: Field Survey, 2015

## RESULT AND DISCUSSION

### RELATIONSHIP BETWEEN WOMEN DEMOGRAPHIC CHARACTERISTICS AND ABUSE AND DISRESPECT DURING CHILDBIRTH

The majority of those who answered the survey questions (52 percent) were between the ages of 31 and 35. According to the findings of the study, the vast majority of respondents (118) had no formal education, 40 respondents had tertiary education, and 29 respondents had vocational education. According to the findings of the study, there was no statistically significant relationship between post-partum women's education, religion, or ethnicity and any form of abuse or disrespect they experienced. According to the findings of the study, 72.3 percent of respondents stated that the educational level of a pregnant woman determines whether or not the person will be abused or disrespected during childbirth, while 27.7 percent of respondents stated that educational level was not associated with abuse and disrespect during childbirth. This finding of the study is in contrast to the findings of Warren et al. (2013) studies conducted in Kenya, which found that there was no significant relationship between education and any form of abuse or disrespect among post-partum women. According to the findings of the study, the vast majority of those who took part in it stated that education and abuse during child labor were inversely related.

**Comment [MA29]:** Please submit the results and the discussion parts in a separate section and re write all the related comments accordingly!

**Comment [MA30]:** With such preliminary educational background, it is quite difficult to announced their opinions whether this study is beneficial to improve or do not the child birth!

The findings of the study are not surprising, given that the majority of the respondents lacked formal education and had previously experienced abuse during childbirth in one form or another, depending on their own understanding of the term. It is important to remember that these respondents took into consideration not only physical abuse during childbirth, but also non-verbal abuse during childbirth. It is possible to engage in nonverbal abuse in a variety of ways, including puffing, ignoring during labour and shouting at the woman to push. The vast majority of respondents (90 percent) were married, with the majority (62.5 percent) identifying as Muslims.

The Dagomba ethnic group was the most prevalent, with an overwhelming majority of the respondents (128) having three or four children at the time of the study's conduct. According to the study's findings, 56 percent of respondents believe that a pregnant woman's financial situation is a contributing factor to abuse and disrespect during childbirth, and that poor pregnant women are more likely than rich pregnant women to suffer abuse and disrespect. 44 percent of those who answered the survey said there was no link between abuse and the financial situation of pregnant women during pregnancy and childbirth. This finding from the study is consistent with the finding made by Haud (2012), who found that pregnant women with high household income are more than six times more likely than pregnant women with low household income to access birth care, with fewer abuses and disrespect at health facilities, than pregnant women with low household income in Addis Ababa, according to the study.

#### **FORMS OF ABUSE AND DISRESPECT**

Motherhood mortality has proven to be an intractable problem despite several decades of global health initiatives aimed at improving the health of women giving birth (Mselle et al., 2013). The maternal mortality ratio, which is a key indicator of maternal health for the Millennium

**Comment [MA31]:** It is quite long and repetitive phrases, the core ideas of the manuscript with such re submission may withdraws the audiences to the black road far away from the main objectives!

Development Goals, has remained essentially unchanged over the past 15 years, with an estimated mean annual decline of 0.4 percent since 1990—a far cry from the progress required to meet the Millennium Development Goal target of 75 percent reduction by 2015. (Mselle et al., 2013).

The risk of dying during pregnancy or shortly after delivery is measured by the maternal mortality ratio, which is primarily used to assess the safety of childbirth in the United States. Most maternal deaths occur in developing countries, with sub-Saharan Africa having the highest maternal mortality ratio in the world (900 deaths per 100 000 live births), according to the World Health Organization (Radoff et al., 2012).

According to the findings of the study, all of the respondents have ever heard of abuse or disrespect during childbirth, which lends support to the findings of Mrisho et al. (2010), who found that abuse and disrespect in facility-based care are extremely common among post-partum women. According to the responses gathered from the respondents, it is clear that there is no single, definitive definition for disrespect and abuse perpetrated against post-partum mothers.

Once again, this finding supports the findings of Ratcliffe (2013), who found that disrespect or abuse toward pregnant women during childbirth in health-care facilities is multidimensional and difficult to define. The results, on the other hand, indicate that some respondents attempted to define abuse or disrespect. Maltreating a pregnant woman during childbirth, the way a pregnant woman is received and handled during childbirth, and is the way a pregnant woman is received and handled during childbirth are some of the highlights. According to Bowser and Hill (2010), yelling during labor, insulting, beating, and providing less care during labor are some of the measures that have been used to describe what pregnant women have actually labeled as abuse

and disrespect during childbirth, and this range and variety of explanations is consistent with their findings.

According to the findings of the study, the majority of respondents (23 percent) identified insulting as a form of abuse and disrespect during childbirth, while 21 percent mentioned non-dignified care, 8 percent stated non-confidential care, and 6 percent stated physical beating. During childbirth, 2 percent of respondents and 40 percent of respondents, respectively, identified the purchasing of items and shouting as forms of abuse and disrespect. In accordance with Ratcliffe (2013), the findings of this study confirm the findings of a previous study conducted in Boston, which revealed that the most common abuse experience described across respondents was that of feeling ignored or neglected. In addition, verbal abuse was common, but it appeared to be less disconcerting among those who responded. As previously stated by Watson-Jones, Balira, and Ross (2012), 79 percent of postpartum women identified insulting as a common form of abuse and disrespect, 16 percent reported being shouted at during labor, and 5 percent of postpartum women reported receiving non-dignified care during the delivery process. Even in a normal situation, insults are the most common thing that causes a pregnant woman to become enraged while in labor. A person has a tendency to retaliate in response to what they perceive as insults. As a result, these respondents were subjected to verbal or nonverbal abuse during their time at work.

### **PERCEPTION AND VIEWS ON ABUSE AND DISRESPECT IN CHILDBIRTH**

Currently, there is broad agreement that the most effective intervention for lowering maternal mortality is universal employment of professional birth attendants based in functional health care facilities who are trained to diagnose obstetric problems and manage or refer them. Most poor countries' primary care strategies include it as a crucial module, as well as a critical component

of the essential package of health services, is the provision of medically supervised childbirth. Despite this, facility delivery rates in Sub-Saharan Africa are among the lowest in the world, with only 47 percent of women in 28 sub-Saharan African countries giving birth in a health facility, according to recent surveys. It was discovered through the study's findings that the vast majority of respondents (86 percent) believed abuse and disrespect during childbirth were harmful, while only 14 percent did not believe these things to be harmful. That this study's findings support previous research conducted by Watson-Jones et al. (2012), who found that post-partum women admitted that abuse and disrespect during childbirth are harmful.

It is only natural that human beings, such as these respondents, would consider abuse of the anuform, and especially abuse during childbirth, to be detrimental to their wellbeing. As a result, it should come as no surprise that the vast majority of respondents stated that they considered abuse to be harmful. The study's findings also revealed that the vast majority (94.1 percent) of respondents had experienced abuse in the past, with only 5.9 percent reporting that they had never been abused before. This finding from the study confirms the findings of Ross (2012), who found that in Kenya, all post-partum women interviewed admitted to having been abused during childbirth in the previous year or two. Even when pregnant women are in severe pain, the instructions given to them by nurses and midwives to follow are often interpreted as abuse by the majority of pregnant women who are in labor. It should be noted that the overwhelming findings of the study, which indicated the majority of respondents had experienced some form of abuse during childbirth, were due to the various forms of abuse identified as a result of the respondents' diverse cultural and ethnic backgrounds, as abuse is always relative. A situation that one person may consider abusive may not even be considered abusive in other contexts, and vice versa.

## CONCLUSION

Many of the post-partum women who took part in this study reported that they were subjected to unfavorable conditions while giving birth in the hospital. One approach to addressing this issue could be to improve the status of women in the community. Some of the structural barriers that contribute to abuse and disrespect, as well as poor maternal care services for these populations during childbirth, are discussed below. Respondents provided a variety of explanations for abuse

**Comment [MA32]:** Please re write the discussion section trying to imagine the ideal frame or the target of the study, very important to the authors to announce and focusing on the aim and all the results should conclude why we are going to investigate such a problems in this area with a limited resources! Inconclusive, to long results and discussion put the image and target of the study in obscure view!

**Comment [MA33]:** It is the site of conclusion, no place to put a further discussions below!

and disrespect, making it difficult to define. The findings revealed that the most common forms of abuse and disrespect in the hospital are: being yelled at, being insulted, receiving substandard care, and purchasing items. Eighty-six percent of respondents considered abuse and disrespect during childbirth to be bad, with 72.3 percent of respondents believing that low education was associated with more abuse and disrespect during childbirth, according to the survey.

**Comment [MA34]:** For me this is the best conclusion, no needs to make the journal readers engaged in a very long section! The authors just need to highlight the exploring of the evidence of abuse and disrespect as the title said!

## References

- Awuah, T. Farouk, G. & Thomas, O. (2010). Assessing evidence of disrespect in public hospitals in Ghana, 45.50
- Baer. H. Singer, M. Johnsen, J. (2010). Toward a critical medical anthropology. *SocSci Med*, Vol. 23:95-98.
- Bartu, D. (2009). Client Satisfaction: Why bother? *Journal of Substance Abuse*, Vol. (1): pp20 – 26.
- Berry, N.S. (2010). Who's judging the quality of care? Indigenous Maya and the problem of "not being attended". *Med Anthropol*, Vol.27:164-189
- Bowser, D. Hill, K. (2014). Exploring evidence for disrespect and abuse in facility-based childbirth: report of a landscape analysis. USAID-TRAction Project, Harvard School of

Public Health and University Research Co., September 20, 2010. Available at [www.traction.org](http://www.traction.org). Accessed June 3, 2015.

Bowser, D. Hill, K. (2010). Exploring evidence for disrespect and abuse in facility-based childbirth. USAID-Traction Project, Harvard School of Public Health.

d'Oliveira, A. Diniz, SG. Schraiber, LB. (2010). Violence against women in health-care institutions: an emerging problem. *Lancet*, Vol. 359:1681-1685.

D'Ambruoso, L., Abbey, M. Hussein, J. (2009). Please understand when I cry out in pain: women's accounts of maternity services during labour and delivery in Ghana. *BMC Public Health*, Vol. 5:140.

Eric Amakwah (2007). Assessing patients' satisfaction level in Ghana: A case study of KomfoAnokye Teaching Hospital, pp45.

- Frisch, A. (2010). Principles of post-operative patients care Nursing Standards. Vol. (5): pp 43-52.
- Jewkes, R. Abrahams, N. Mvo, Z. (2010). Why do nurses abuse patients? Reflections from South African obstetric services. *SocSciMed* , Vol. 47:1781-1795.
- Filippi, V. Ronsmans, C. Campbell, O. Graham, W., Mills, A. Borghi, J. Koblinsky, M. Osrin, D. (2011). Maternal health in poor countries: the broader context and a call for action. *Lancet*, Vol. 368:1535-1541.
- Kruk, M.E. Paczkowski, M. Mbaruku, G. de Pinho, H. Galea, S (2009). Women's preferences for place of delivery in rural Tanzania: a population-based discrete choice experiment. *Am Public Health*, Vol. 99:1666-1672.
- Kruk, M. Kujawski, S. Mbaruku, G. Ramsey, K. Moyo, W. Freedman, L. (2014). Disrespectful and abusive treatment during facility delivery in Tanzania: a facility and community survey. Health Policy Plan. doi:10.1093/heapol/czu079
- Kruger, L. Schoombee, C. (2010). The other side of caring: abuse in a South African maternity ward. *J Reprod Infant Psychol*, Vol.28:84-101
- Kumbani, L.C. Chirwa, E. Odland, J. Bjune, G. (2012). Do Malawian women critically assess the quality of care? A qualitative study on women's perceptions of perinatal care at a district hospital in Malawi. *ReprodHealth*, Vol. 9:30.
- Mamdani, M. Bangser, M. (2010). Poor people's experiences of health services in Tanzania: a literature review. *Reprod Health Matters*, Vol. 12:138-153.

- Moyer, C. Adongo, P. Aborigo, R.A. Hodgson, A. Engmann, CM. (2014). 'They treat you like you are not a human being': maltreatment during labour and delivery in rural northern Ghana. *Midwifery*, Vol. 30(2):262-268.
- Mrisho, M. Schellenberg, J. Mushi, A.K. Obrist, B. Mshinda, H. Tanner, M. Schellenberg, D. (2010). Factors affecting home delivery in rural Tanzania. *Trop Med Int Health*, Vol. 12:862-872
- Mselle, LT. Moland, KM. Mvungi, A. Evjen-Olsen, B. Kohi, TW. (2013). Why give birth in health facility? Users' and providers' accounts of poor quality of birth care in Tanzania. *BMC Health Serv Res*, Vol. 13:174
- Radoff, K.A. Levi, A. Thompson, L.M. (2012). From home to hospital: mistreatment of childbearing women and barriers to facility-based birth in Nicaragua. *Int J Childbirth*, Vol. 2:40-50.
- Ratcliffe, H. (2013). *Creating an Evidence Base for the Promotion of Respectful Maternity Care*. Boston, MA: Harvard School of Public Health
- Reichenheim, M. Zylbersztajn, F. Moraes, C., Lobato, G. (2009). Severe acute obstetric morbidity (near-miss): a review of the relative use of its diagnostic indicators. *ArchGynecolObstet*, Vol. 280:337-343.
- Rosenfield, A. Maine, D. (2010). Maternal mortality—a neglected tragedy. Where is the M in MCH? *Lancet*, Vol.2:83-85.

- Schroll, A.M. Kjærgaard, H. Midtgaard, J. (2013). Encountering abuse in health care; lifetime experiences in postnatal women-a qualitative study. *BMC Pregnancy Childbirth*, Vol.13:74.
- Shimpuku, Y. Patil, CL. Norr, KF. Hill, PD. (2013). Women's perceptions of childbirth experience at a hospital in rural Tanzania. *Health Care Women Int* , Vol. 34:461-481.
- Souza, J. Gülmezoglu, A. Vogel, J. Carroli, G. Lumbiganon, P. Qureshi, Z.(2013). Moving beyond essential interventions for reduction of maternal mortality (the WHO Multicountry Survey on Maternal and Newborn Health): a cross-sectional study. *Lancet*; Vol. 381(9879):1747–55. [http://dx.doi.org/10.1016/S0140-6736\(13\)60686-8](http://dx.doi.org/10.1016/S0140-6736(13)60686-8) pmid: [23683641](https://pubmed.ncbi.nlm.nih.gov/23683641/)
- Thaddeus, S. Maine, D. (2009). Too far to walk: maternal mortality in context. *SocSci Med*, Vol. 38:1091-1110.
- Turan, J. Nyblade, L. (2013). HIV-related stigma as a barrier to achievement of global PMTCT and maternal health goals: a review of the evidence. *AIDS Behav*;17:2528–2539.
- UNICEF (2013). *The State of the World's Children*. New York, NY: United Nations Publications.
- United Nations High Commissioner for Human Rights (2012). Technical guidance on the application of a human rights-based approach to the implementation of policies and programmes to reduce preventable maternal morbidity and mortality [A/HRC/21/22]. New York: United Nations.

Warren, C. Njuki, R. Abuya, T. Ndwiga, C. Maingi, G. Serwanga, J. Mbehero, F. Muteti, L. Njeru, A. Karanja, J. (2013). Study protocol for promoting respectful maternity care initiative to assess, measure and design interventions to reduce disrespect and abuse during childbirth in Kenya. *BMC Pregnancy Childbirth* , Vol. 13:21.

Watson-Jones, D. Balira, R. Ross, DA. (2012). Missed opportunities: poor linkage to ongoing care for HIV-positive pregnant women in Mwanza, Tanzania. *PLoS One*;7:e40091.

UNDER PEER REVIEW