

# Original Research Article

## Antidiabetic and Toxicity Studies of the Extract of Four Nigerian Medicinal Plants

### ABSTRACT

**Aims:** To evaluate the anti-hyperglycaemic efficacy and safety of the methanol extract of the combination of *Senecio bialfrae* leaf, *Xylopia aethiopica* fruit, *Carica papaya* seed and *Spondias mombin* stem bark mixed together in ratio 1:1:1:1

**Study Design:** Extract of medicinal plants was assayed using glucose and streptozotocin-induced hyperglycaemic rats model.

**Place and Duration of Study:** Department of Pharmacognosy, Obafemi Awolowo University, IleIfe, Nigeria, between May, 2019 and January, 2022.

**Methodology:** The extract of the combined plant parts was tested for toxicity in rats while its effects on glucose level, blood and biochemical components were also assessed. Its *in-vitro* anti-hyperglycaemic activity was assayed in  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitory models while its *in-vivo* effects were tested in glucose and streptozotocin-induced hyperglycaemic rats. The antioxidant activity of the extract was also carried out.

**Results:** The extract did not show any adverse effects on blood sugar levels, haematological and biochemical parameters in normal rats in sub acute toxicity tests. The extract gave comparable ( $p > 0.05$ )  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitory effects to acarbose. In glucose-induced hyperglycaemic rats, its 100 mg/kg was the most effective dose with 19, 40, 43, and 57 % activity that was significantly higher ( $p < 0.05$ ) than the 10, 18, 24, and 40 % activity given by glibenclamide (5 mg/kg) at the same time points. In streptozotocin-induced diabetic assay, its 50 mg/kg showed 31, 85, 85 and 82 % effects on days 4, 7, 10 and 14, respectively that was significantly higher than its 100 mg/kg and glibenclamide on days 7 and 10. The extract also elicited high free radical scavenging effects in all the antioxidant assays.

**Conclusion:** The extract of the combination of four Nigerian antidiabetic plants mixed together in equal ratio gave significantly better antidiabetic activity at low doses than the individual plants without toxic effects.

**Keywords:** Diabetes mellitus, anti-hyperglycaemic effect, plant combination, antioxidant activity.

### 1.0. INTRODUCTION

Diabetes mellitus is a collection of severe metabolic dysfunction marked by hyperglycaemia as a result of absence of secretion of insulin by pancreatic  $\beta$ -cells, inefficient insulin usage by the body, or both [1-3]. Globally, diabetes has been reported to affect estimates of 415 million in 2015, 425 million in 2017 and 463 million persons in 2019. This number is expected to surge to 578 and 700 million in 2030 and 2045, respectively with prevalence in low and middle-income

countries than in high-income countries [3,4]. Diabetes mellitus can only be managed, not cured and majority of the available synthetic antidiabetics are accompanied with serious side effects hence, the need for investigation of new drugs from natural sources is highly essential [5-7].

Traditional medicine often offers polyherbal therapy with individual component plants managing different symptoms and exerting its effect on different organs. This concept is similar to the theory of poly-pharmacy [8]. The use of herbs as combination therapy in the management of several disease conditions such as cancer, acquired immunodeficiency syndrome (AIDS), malaria, pulmonary, tuberculosis, diabetes mellitus, etc has resulted in many outcomes. These include: optimum activity at lower dose than individual plants, additional pharmacological effect exerted by the individual components of the extract, synergistic effect of the different components in the extract and ease of administration resulting in a better adherence to herbal drug use to achieve enhanced therapeutic effects [9]. Although, herb-herb combinations have been in use in traditional medicine practice for several years, scientific understanding of various herb-herb interactions or proof of clinical benefits have been slow and weak resulting in the need for further research in herbal combination therapy [8]. Poly-herbal formulations in the management of diabetes mellitus have been variously reported [10-15].

*Senecio biafrae* Oliv. & Hiern (Asteraceae), known as 'worowo' in Yoruba (Nigeria) and 'Gnanvule' (Coted'Ivoire) [16] is used ethnomedically as galactagogue [17] and management of diabetes in combination with the seeds of *Aframomum melegueta* and *Carica papaya* [18]. Its antihyperglycaemic, hypolipidemic and antioxidant activities have been reported [19,20,21]. *Xylopiya aethiopica* (Dunal) A. Rich (Annonaceae) is used in Nigeria to manage diabetes, hypertension, hyperlipidaemia and obesity [22]. It has been reported for its antihyperglycaemic [23], antiproliferative [24] and hypolipidemic and antioxidant potentials [25]. Different parts of *Carica papaya* L. (Caricaceae) are used in the management of diabetes, hypertension and stroke [22]. It has been reported for its antihyperglycemic, anti-inflammatory and antioxidant activities [26], larvicidal [27], antisickling [28] activities. *Spondias mombin* L. (Anacardiaceae) is used in Nigeria and African countries in the management of diabetes [18,22,29,30].

These plants have been individually studied for their antidiabetic activities in our research laboratories and the findings have been reported [21,23,26,31]. This work was therefore designed to investigate the antidiabetic activities of the combined extract of the active plant parts in equal ratio and its safety in order to discover the possible advantage(s) of the combined plant extract over the individual plants.

## **2.0. MATERIALS AND METHODS**

### **2.1. Chemical Equipment and Instrumentation**

Rotary evaporator (RE301/601/801 model, Yamato Scientific America, Inc., U.S.A), chiller (Churchill, Instrument Co. Ltd, U.K), vacuum pump (MB 338618 model, Edwards High Vacuum Int., England), oven (Hearson & Co. Ltd, London), Mettler electronic weighing balance (AB 54 model, Mettler Toledo, U.S.A), Ultra-violet (UV) lamp (254 and 366 nm) (Grant Instrument, U.K), Oral cannula, ACCU-CHEK Glucometer (model GB 11558973, Roche, Germany) with ACCU-CHECK test strips (Roche, Germany), UV spectrophotometer, Dutrao (Model SM 600, Shang Yhai Yong Chuang Medical Instrument Co. Ltd) spectrophotometric microplate reader, Automated haematology analyzer, Centrifuge, Semi-automated biochemistry analyzer.

## 2.2. Plant Materials and Extraction

*Senecio biafrae* leaf, *Carica papaya* seed and *Spondias mombin* stem bark were collected on Obafemi Awolowo University campus while *Xylopia aethiopica* fruit was purchased from Oja Oba market, Modakeke, Osun State. They were authenticated and their voucher specimens with herbarium specimen numbers, FPI 2324, 2323, 2325 and 2326, respectively were deposited at the Faculty of Pharmacy Herbarium. The various morphological plant parts were separately air-dried, powdered and mixed together in ratio 1:1:1:1. A total of 2600 g of plant material (containing 650 g each of the individual plant part) was extracted by maceration in methanol with mechanical agitation. The marc was re-extracted three times and concentrated *in-vacuo* to obtain a yield of 20.2 % w/w.

## 2.3. Animals

Healthy Wistar rats (120-180 g) of both sexes were bred under standard conditions (temperature 27-30°C, relative humidity 65 %) in the animal house, Department of Pharmacology, Faculty of Pharmacy, O.A.U., Ile-Ife, Nigeria. They were given regular pellets diet (Bendel Feeds, Nigeria) and water *ad libitum*. Ethical clearance number IPH/OAU/12/1758 was issued by the Health Research Ethics Committee, Institute of Public Health, Obafemi Awolowo University on the use of animals for this work.

## 2.4. Acute Toxicity Test

Acute toxicity test was carried out according to the modified OECD Test Guideline 423 Annex 3 model [32]. Two groups of 8 animals each were administered distilled water and single oral administration of 5000 mg/kg extract, respectively. They were observed for signs of gross toxicity, behavioural changes and mortality, one hour after administration and daily for 14 days [32].

## 2.5. Subacute Toxicity Test

Sub acute toxicity test was carried out following the modified OECD Test Guideline 407 [33]. The extract was solubilised in 1 % Tween 80 in distilled water and administered to groups of 8 rats daily for 28 days at graded doses of 250, 500, and 1000 mg/kg. The blood glucose levels in rats were monitored on days 7, 14, 21 and 28. The animals were anaesthetized using chloroform and blood sample (5 mL) collected by cardiac puncture after the 28<sup>th</sup> day [33].

## 2.6. Haematological Analysis

About 50 µL of blood was aspirated into the automated haematology analyzer for haematological analysis [34].

## 2.7. Biochemical Assays

In the assay of transaminases (Aspartate Transaminase (AST), Alanine Transaminase (ALT)), serum (100 µL) was added to 1000 µL of working solution of SGOT R1 and R2 (4:1) and the mixture was aspirated into the analyser to obtain the absorbance for the AST. While 100 µL of serum was added to 1000 µL of working solution of SGPT R1 and R2 (4:1) and the mixture was aspirated into the analyser to obtain the absorbance for the ALT Serum (20 µL) was added to

1000 µL of working solution containing one tablet of ALP substrate and 10 mL of ALP diluents and the solutions were aspirated into the analyser to obtain the absorbance of the blank and test. To 1000 µL of working solution containing equal volume of creatinine picrate and diluents was placed inside three test tubes for blank, standard and test for the Creatinine test. Serum sample (50 µL) was added into the tube for test while creatinine standard (50 µL) was added into the tube for standard. The solutions were aspirated into the analyser to obtain the absorbance of the blank, standard and test. Serum (10 µL) and urea standard (10 µL) were added to working solution containing urea R1 and R2 (4:1) placed inside two test tubes for the standard and test. The solutions were aspirated into the analyser to obtain the absorbance of the standard and test. Serum (10 µL), cholesterol standard (10 µL) were added into the tubes for test and sample containing 1000 µL working solution of cholesterol reagent. The mixtures were incubated at 37°C for 10 minutes after which they were aspirated into the analyser and the absorbance for the blank, standard and test were obtained.

## 2.8. Antidiabetic Studies

### 2.8.1. *In-vitro* α-amylase inhibitory activity of the extract

The assay mixture consists of 1.0 mL of 0.020 M solution of sodium phosphate buffer (pH 6.90 with 0.0060M NaCl), 1 mL alpha amylase solution (from *Aspergillus oryzae*) and 0.4 mL extract at different concentration (0.05, 0.1, 0.25, 0.5, 1.0 mg/mL). Pre-incubation was done at 37 °C for 10 minutes, followed by the addition of 1 mL of 1 % solution of boiled potato starch into the tubes. The reaction mixture was kept at 37°C for another 15 minutes before being stopped with 1 mL of 3,5-dinitrosalicylic acid (DNSA) (containing 1.00 g of 3,5-dinitrosalicylic acid, 20.0 mL of 2.0 M NaOH and 30.0 g of sodium potassium tartarate in 100 mL distilled water). After 5 minutes in a boiling water bath, the tubes were cooled to room temperature. The mixture was diluted with 5 mL distilled water, and the absorbance was measured with a spectrophotometer at 540 nm. With the exception of the extract, a control indicating 100 % enzyme activity was carried out in a similar manner. The synthesis of reducing sugar occurs when 3,5-dinitrosalicylic acid is reduced to 3-amino-5-nitrosalicylic acid. Acarbose was used as the positive control. The following was used to calculate the alpha amylase inhibition and represented as a % of inhibition:

$$\text{Inhibition (\%)} = \frac{[(A_{c+}) - (A_{c-})] - [A_s - A_b]}{\{(A_{c+}) - (A_{c-})\}} * 100$$

Where:

$A_{c+}$  = absorbance of 100 % enzyme activity,

$A_{c-}$  = 0 % enzyme activity (only solvent without enzyme),

$A_s$  = test sample (with enzyme) and

$A_b$  = blank (a test sample without enzyme) [35,36].

### 2.8.2. *In-vitro* α-glucosidase inhibitory activity of the extract

The α-glucosidase inhibitory assay was carried out with a Dutrao spectrophotometric microplate reader (Model SM 600, Shangyhai Yong chuang Medical Instrument Co. Ltd). A total of 60.0 µL

of reaction mixture was added to each well, consisting of 20.0  $\mu$ L of 100.0 mM phosphate buffer (pH 6.80), 20.0  $\mu$ L of 2.50 mM (pNPG) 4-Nitrophenyl-D-glucopyranoside, and 20.0  $\mu$ L methanol dissolved sample, afterwards 20.0  $\mu$ L of 10.0 mM phosphate buffer (pH 6.80) containing 0.20 U/mL alpha glucosidase to the mixture in the wells. The reaction was incubated at 37°C and terminated with 80.0  $\mu$ L of 0.20 mM sodium carbonate 15 min later. The absorbance was estimated using a microplate reader at 405.00 nm. The above procedure was repeated using acarbose as the positive control [37].

### **2.8.3. Antihyperglycaemic effect of extract on glucose induced hyperglycaemic rats**

Groups of 5 rats each fasted for 18 hours and given 10 g/kg of glucose (*p.o.*) were used for the experiment. After 0.5 hour (time point 0), rats having blood glucose levels  $\geq 7.0$  mmol/L (126 mg/dL) were considered hyperglycaemic and given (*p.o.*) vehicle (Tween 80 (1 %) in distilled water), methanol and aqueous extracts separately (25, 50, 100 and 200 mg/kg), and 5.0 mg/kg glibenclamide. At 0.00, 0.50, 1.00, 2.00, and 4.00 hours, blood drop from each rat's caudal vein was placed on to a glucometer strip inserted into the glucometer. The percentage decrease in blood glucose level at these time points were calculated and compared to the negative and positive controls [38-41].

### **2.8.4. Antihyperglycaemic effect of extract on streptozotocin-induced diabetic rats**

Streptozotocin (65 mg/kg) in freshly prepared buffer solution (0.1M, pH 4.5) was intraperitoneally administered to overnight fasted rats to induce diabetes. After 72 hours of induction, the rats' blood glucose levels were measured, and they were then left for another 5 days. Rats with Fasting Blood Sugar  $\geq 11.0$  mMol/L were considered diabetic and divided into four groups of five rats: negative control, administered with 1 % Tween 80 in distilled water; test groups, 50 and 100 mg/kg and positive control, glibenclamide (5 mg/kg). On days 1, 4, 7, 10, and 14, blood glucose levels were measured and compared to the control group [5].

## **2.9. In vitro Antioxidant Studies**

### **2.9.1. 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity**

The DPPH or radical scavenging properties of the extract was determined using the stable radical 1,1-diphenyl-2-picrylhydrazyl (DPPH) according to [26,39,42]. The absorbance was then measured at 517 nm against a DPPH control containing 1 ml of methanol in place of the extract. The scavenging activity was then calculated using the formula below:

$$\text{Percentage scavenging activity} = \frac{\text{absorbance control} - \text{absorbance sample}}{\text{absorbance control}} \times 100$$

The  $IC_{50}$  was then obtained from a linear regression plot of percentage inhibition against concentration of extract.

### **2.9.2. Evaluation of Total Antioxidant Capacity (TAC)**

The total antioxidant assay was carried out based on the reduction of Mo (VI) to Mo (V) by the extract and subsequent formation of a green phosphate/Mo (v) complex at acidic pH [43].

### **2.9.3. Ferric Reducing Antioxidant Power (FRAP)**

The FRAP assay uses antioxidant as reductants in redox linked colorimetric method [44].

#### **2.9.4. Hydroxyl Radical Scavenging Activity (HRSA)**

The ability of the extracts to scavenge the hydroxyl radical generated by the Fenton reaction was measured according to the modified method of [45].

#### **2.9.5. Determination of Total Flavonoid Content**

The estimation of the Total Flavonoid Content of the extract was based on the Aluminium chloride colorimetric method according to the method of [46] and as described by [47]. The result was expressed as mg rutin equivalent (RE)/g of the extract. Analysis was done in triplicates.

#### **2.9.6. Total Phenolic Content (TPC)**

The total phenol content of the extracts was determined using Folin-Ciocalteu's method of [48] Singleton as described by Gulcin [49]. The principle is based on reduction of phosphomolybdic-phosphotungstic acid (Folin reagents) to a blue-coloured complex in an alkaline solution which occurs in the presence of phenolic compounds.

#### **2.10. Statistical analysis.**

Data were expressed as the mean  $\pm$  SEM for the number (n) of animals in the group. Analysis of variance (ANOVA) then Student Newman Keul's test was used to obtain the source of significant differences for all determination.  $P < 0.050$  was considered as statistically significant.

### **3.0. RESULTS AND DISCUSSION**

#### **3.1. Acute Toxicity of the Methanol Extract**

The result of the acute toxicity study of the methanol extract showed no observable changes in behaviour of the rats with respect to breathing, cutaneous effect, sensory and nervous system responses or gastrointestinal effect. No mortality or gross toxicity was also observed. Thus, the median lethal dose, LD<sub>50</sub> was estimated to be greater than 5000 mg/kg, suggesting that the extract is safe and possesses low risk of toxicity. It also indicated that the doses of 25-1000 mg/kg used in the study were experimentally safe. Extracts of *S. bialfrae*, *X. aethiopica*, *C. papaya* and *S. mombin* had previously been reported to possess LD<sub>50</sub> at  $> 5000$ ,  $3464$ , and  $> 2000$   $> 2000$  mg/kg, respectively [21,50-52].

#### **3.2. Sub acute Effect of the Extract on Blood Glucose Level in Normal Rats**

The negative control normoglycaemic rats that were given distilled water did not show any appreciable decrease in the level of blood glucose of the rats throughout the period of the study confirming that the water did not contain any bioactive constituents. The extract of the plant combination at the tested doses of 250, 500 and 1000 mg/kg and for all the days of the experiment had comparable ( $p > 0.05$ ) effect on the blood glucose level of the rats to the negative control (Table 1). This indicated safety of the extract and that it cannot precipitate hypoglycaemic coma when administered to non diabetic individuals. Similar effect has been reported for *Eugenia uniflora*, *Olex subscorpioidea*, *Entandrophragma cylindricun*, *Triclisia subcordata* [40,41,53,54].

#### **3.3. Effect of Extract on Haematological Parameters**

Investigation of haematological components of humans/animals, such as red blood cells, white blood cells or leucocytes, mean corpuscular volume, mean corpuscular haemoglobin and mean corpuscular haemoglobin concentration is valuable in monitoring toxicity as well as the health status of animals [55]. Administration of 250, 500, 1000 mg/kg of the extract to rats for 28 days in this study showed a significant increase in RBC level which suggested positive effect on the haemopoietic system of the test rats indicating possible anti-anaemic effect of the extract (Table 2). However, it gave no significant difference in other evaluated haematological parameters when compared to the control. This indicated that the extract had no adverse effect on blood components and further established its lack of toxicity. Extracts of *Mangifera indica* stem bark and *Telfaria occidentalis* had been reported to show similar effects on the haematological components in rats [56,57].

### **3.4. Effect of Extract on Biochemical Parameters**

The results of the biochemical analysis of the blood samples after treatment for 28 days showed comparable activity ( $p > 0.050$ ) of the extract on aspartate transaminase (AST), alanine transaminase (ALT), creatinine and urea level to distilled water (negative control) (Table 3). This result indicated the extract does not have any toxic effect on the heart, liver and kidney of the rats. High AST and ALT levels have been associated with liver diseases or hepatotoxicity [58,59]. The extract however consistently elicited a reduction in cholesterol level at all the tested doses compared to negative control showing possible antihyperlipidemic effect of the extract. Hypolipidaemic activity has been reported for *Senecio bialfrae*, *Carica papaya* and *Xylopi aethiopia* [19,20,25] which confirmed the findings of this study.

### **3.5. *In vitro* $\alpha$ -amylase Inhibitory Effect of the Extract**

Pancreatic  $\alpha$ -amylase and  $\alpha$ -glucosidase play a critical role in carbohydrate digestion and effective inhibitors of these enzymes such as acarbose, miglitol and voglibose are being used in the management of diabetes [60-62]. However, their side effects necessitated a search for new  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitors of natural origin without side effects and to provide more candidates of drug choices. In the  $\alpha$ -amylase inhibitory assay in this study, the extract showed a significant concentration dependent antihyperglycaemic effect from 0.05-1.0 mg/mL similar to acarbose. The extract with  $IC_{50}$  of 0.06 mg/mL that was lower than 0.1 mg/mL of the positive control indicated better  $\alpha$ -amylase inhibitory antihyperglycaemic potential of the extract (Table 4).

### **3.6. *In vitro* $\alpha$ -glucosidase Inhibitory Effect of Extract**

Similar to the  $\alpha$ -amylase inhibitory antihyperglycaemic result (Table 4), the extract exhibited a concentration dependent antihyperglycaemic activity that was comparable to the positive control (Table 5). The  $IC_{50}$  values of  $0.17 \pm 0.02$  and  $0.19 \pm 0.01$  for the extract and acarbose, respectively indicated the extract like acarbose effectively inhibited the breaking down of complex carbohydrates to glucose thereby preventing glucose assimilation from the intestine into the bloodstream [26,63]. This confirmed the extrapancreatic activity in addition to the insulin stimulation effect that had been reported for *Senecio bialfrae* [21] and the  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitory activities reported for *Carica papaya* [26].

### **3.7. Anti-hyperglycaemic Activity of the Extract**

The results of glucose-loaded rat model or Oral Glucose Tolerance Test (OGTT) model with glibenclamide and other insulin-stimulating drugs as positive controls have been reported to mimic type 2 diabetes state in humans and reportedly used in anti-hyperglycaemic assays of medicinal plants [6,7,64,65]. The glucose loaded hyperglycaemic rats that received distilled water (negative control) gave a significant ( $p < 0.050$ ) time-dependent decrease in blood glucose level till the fourth hour due to homeostatic regulatory process in normal animals (Fig.1) [6,26,66]. Generally, the extract at the tested doses of 25, 50, 100 and 200 mg/kg elicited a time dependent antihyperglycaemic effect till the fourth hour similar to glibenclamide (5 mg/kg) which suggested that the extract may have early extrapancreatic and late insulin stimulating mechanism of action of glibenclamide [41,53,54,67]. The activity of the extract was dose dependent with maximum effect at 100 mg/kg that was significantly better ( $p < 0.050$ ) than glibenclamide (5 mg/kg) at all time points which indicated an additional extapancreatic effect of the extract at this dose (Fig.1). High blood glucose level reduction of 19, 40 and 43 % observed between 0.5-2 h demonstrated by 100 mg/kg of the extract which indicated better extrapancreatic effect was justified by the high  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitory activities exhibited by the extract (Tables 4 and 5). In the previous work done on the individual plants on this combination [21,23,26], highest antihyperglycaemic effect was obtained at 400 mg/kg. This suggested that the antihyperglycaemic activity of the component plants were highly enhanced in the combination and showed optimum activity at lower dose of 100 mg/kg indicating synergism (Fig.1).

### **3.8. Effect of the Extract on Streptozotocin-induced Diabetic Rats**

The most active dose in the glucose-induced antihyperglycaemic experiment and its lower dose, 100 and 50 mg/kg, respectively (Fig. 1) were used in this experiment. The non treated diabetic rats constititently maintained an hyperglycaemic state throughout the study period showing that the diabetes induced by the administered streptozotocin was permanent (Fig. 2). Glibenclamide (5 mg/kg), the positive control drug gave a time dependent antidiabetic activity of 13, 56, 75, 79 % on days 4, 7, 10 and 14, respectively that was due to insulin stimulating action of the drug on the remaining pancreatic  $\beta$ -cells of the diabetic rats (Fig. 2). The extract of the plant combination at 50 mg/kg, with 31, 85 and 85 % antidiabetic effect on days 4, 7 and 10, respectively were significantly higher than the 14, 56 and 75 % activity of glibenclamide on the same days and 71 and 72 % of 100 mg/kg on days 7 and 10. This indicated a better activity of the extract at 50 mg/kg (Fig. 2). This observation further confirmed that the antidiabetic effect of the combination plant extract was significantly better at lower doses than the individual plants, an advantage poly-herbal therapy [9,18,21,23,31].

### **3.9. Antioxidant Activity of the Extract**

The antioxidant activity of the extract was determined using multiple assays (DPPH, FRAP, TAC, HRSA, TPC and TFC) in order to determine the possible contribution of antioxidant effect of the extract to its antidiabetic activity [21]. In the DPPH assay (Table 6), the extract gave an  $IC_{50}$  value of 0.341 mg/ml which showed a better radical scavenging effect than that reported for *S. bialfrae* leaf and *C. papaya* seeds alone [21,26]. The combined plant extract with values 17.4 mgAAEq/g and  $IC_{50}$  of 0.368 mg/ml for FRAP and HRSA assays indicated better activity than that reported for *S. bialfrae* and *C. papaya* seeds alone [21,26]. The antioxidant activity of the extract was significantly higher in the TAC assay compared to that reported for *C. papaya*, *X. aethiopica*, *S. bialfrae* and *S. mombin* alone. The extract also showed high values for TFC and

TPC (Table 6), indicating that it was rich in phenolics and flavonoids which was confirmed for *S. bialafrae*, *C. papaya* and *S. mombin* [21,26,68].

**Table 1: Effect of extract on blood glucose level in normal rats**

Extract/Drug	Day 1	Day 7	Day14	Day 21	Day 28
(mg/kg)					
DW	100	95.36±3.37 <sup>a</sup>	94.26±4.45 <sup>a</sup>	99.34±3.00 <sup>a</sup>	100.26±2.81 <sup>a</sup>
SXCS (250)	100	102.31±7.00 <sup>a</sup> (-7.29%)	94.59±4.83 <sup>a</sup> (-0.35%)	107.05±9.00 <sup>a</sup> (-7.76%)	89.49±5.24 <sup>a</sup> (10.74%)
SXCS (500)	100	94.66±4.18 <sup>a</sup> (0.73%)	91.92±5.54 <sup>a</sup> (2.48%)	102.50±3.16 <sup>a</sup> (-3.18%)	85.57±3.61 <sup>a</sup> (14.65%)
SXCS (1000)	100	94.81±2.71 <sup>a</sup> (0.58%)	102.48±1.75 <sup>a</sup> (-8.72%)	104.88±6.18 <sup>a</sup> (-5.58%)	96.58±4.45 <sup>a</sup> (3.67%)

Data show the mean ± SEM blood glucose levels at the different time points (Tt) expressed as percentages of level at day 1, percentage reductions in the bgl's relative to negative control for each time point, N = 8. Values with similar superscript are comparable (p>0.05). One-way analysis of variance followed by the Student-Newman-Keuls' post-hoc test). **DW**: Distilled Water; **SXCS** (250, 500, 1000): *S. bialafrae*, *X. aethiopica*, *C. papaya*, *S. mombin* extract.

**Table 2: Effect of extract on haematological parameters in normal rats**

Blood Parameters	D W	SXCS(250 mg/kg)	SXCS(500 mg/kg)	SXCS(1000 mg/kg)
WBC (10 <sup>3</sup> /μL)	15.49±0.82 <sup>a</sup>	18.34±1.27 <sup>a</sup>	15.34±1.22 <sup>a</sup>	18.34±2.40 <sup>a</sup>
RBC (10 <sup>6</sup> /μL)	6.60±0.22 <sup>a</sup>	7.30±0.22 <sup>a,b</sup>	7.41±0.30 <sup>a,b</sup>	7.90±0.43 <sup>b</sup>
HGB (g/dL)	12.16±0.33 <sup>a</sup>	12.86±0.31 <sup>a</sup>	13.20±0.19 <sup>a</sup>	12.49±0.84 <sup>a</sup>
HCT (%)	46.66±1.91 <sup>a</sup>	48.79±1.80 <sup>a</sup>	46.17±1.84 <sup>a</sup>	50.68±2.62 <sup>a</sup>
MCV(fl)	70.66±1.12 <sup>b</sup>	66.73±1.26 <sup>a</sup>	62.39±0.79 <sup>a</sup>	64.33±1.92 <sup>a</sup>
MCH(pg)	18.51±0.49 <sup>a</sup>	17.64±0.41 <sup>a</sup>	18.01±0.85 <sup>a</sup>	16.08±1.30 <sup>a</sup>
MCHC(%)	26.24±0.79 <sup>a</sup>	26.34±0.54 <sup>a</sup>	28.89±1.26 <sup>a</sup>	24.89±1.81 <sup>a</sup>
Platelet(10 <sup>3</sup> /μL)	444.25±77.09 <sup>a</sup>	498.29±87.89 <sup>a</sup>	454.29±79.78 <sup>a</sup>	644.50±94.01 <sup>a</sup>

Data show the mean ± SEM haematological parameters at the different doses, n=8. Results having separate superscripts within row are significantly different (p < 0.050), while those that

are alike are comparable ( $p > 0.050$ ): One-way variance analysis (ANOVA) followed by Student-Newman-Keul's test. **DW**: Distilled Water; **SXCS** (250, 500, 1000): *S. bialafrae*, *X. aethiopica*, *C. papaya*, *S. mombin* extract. **WBC**: White Blood Corpuscles, **RBC**: Red Blood Corpuscles, **HGB**: Haemoglobin, **HCT**: Haematocrit, **MCV**: Mean Corpuscular Volume, **MCH**: Mean Corpuscular Haemoglobin, **MCHC**: Mean Corpuscular Haemoglobin Concentration.

**Table 3: Effect of extract on biochemical parameters in normal rats**

Biochemical Parameters	D W	SXCS (250 mg/kg)	SXCS (500 mg/kg)	SXCS (1000 mg/kg)
AST	351.95±14.08 <sup>a</sup>	339.00±29.92 <sup>a</sup>	287.15±17.26 <sup>a</sup>	367.85±20.62 <sup>a</sup>
ALT	65.87±3.26 <sup>a</sup>	57.17±5.33 <sup>a</sup>	53.98±3.36 <sup>a</sup>	60.82±4.10 <sup>a</sup>
CREA	6.77±0.10 <sup>a</sup>	6.47±0.19 <sup>a</sup>	5.65±0.41 <sup>a</sup>	6.20±0.42 <sup>a</sup>
CHOL	256.55±18.89 <sup>b</sup>	196.55±17.12 <sup>a</sup>	199.20±7.32 <sup>a</sup>	201.20±5.73 <sup>a</sup>
ALP	149.87±24.84 <sup>b</sup>	138.41±12.56 <sup>a,b</sup>	81.42±12.42 <sup>a</sup>	119.23±15.18 <sup>a,b</sup>
UREA	23.81±0.71 <sup>a</sup>	23.56±0.37 <sup>a</sup>	22.49±0.05 <sup>a</sup>	22.98±0.12 <sup>a</sup>

Data show the mean ± SEM biochemical parameters at different doses, n=8. Results having separate superscripts within row are significantly different ( $p < 0.050$ ), while those that are alike are comparable ( $p > 0.05$ ). **DW**: Distilled Water; **SXCS** (250, 500, 1000): Extract of *S. bialafrae*, *X. aethiopica*, *C. papaya*, *S. mombin*. **AST**: Aspartate Transaminase, **ALT**: Alanine Transaminase **CREA**: Creatinine, **CHOL**: Cholesterol, **ALP**: Alkaline phosphatase

**Table 4: *In vitro*  $\alpha$ -amylase inhibitory effect of extract**

Concentration of extract and Acarbose (mg/ml)	Average % Inhibition		IC <sub>50</sub> (mg/kg)	
	Extract	Acarbose	Extract	Acarbose
1	85.63 <sup>a</sup>	83.09 <sup>a</sup>	0.06±0.02 <sup>a</sup>	0.10±0.01 <sup>a</sup>
0.5	77.59 <sup>a</sup>	80.2 <sup>a</sup>		
0.25	68.73 <sup>b</sup>	63.03 <sup>a</sup>		
0.1	64.75 <sup>b</sup>	55.31 <sup>a</sup>		
0.05	35.45 <sup>a</sup>	46.60 <sup>b</sup>		

Data show the mean ± SEM  $\alpha$ -amylase inhibitory effect of extract at different concentrations, n=3. Values with different superscripts within rows are significantly different ( $p < 0.05$ , one-way


analysis of variance followed by the Student–Newman–Keuls' test); Results with separate superscripts within rows are significantly different ( $p < 0.050$ ), while those that are alike are comparable ( $p > 0.05$ ).

**Table 5: *In vitro*  $\alpha$ -glucosidase inhibitory effect of extract**

Concentration of extract and Acarbose (mg/mL)	Average % Inhibition		IC <sub>50</sub> (mg/mL)	
	Extract	Acarbose	Extract	Acarbose
0.5	68.21 <sup>a</sup>	89.52 <sup>b</sup>	0.17±0.02 <sup>a</sup>	0.19±0.01 <sup>a</sup>
0.25	58.24 <sup>a</sup>	59.43 <sup>a</sup>		
0.125	48.64 <sup>b</sup>	40.69 <sup>a</sup>		
0.0625	38.86 <sup>b</sup>	31.80 <sup>a</sup>		
0.03125	11.21 <sup>a</sup>	12.40 <sup>a</sup>		


Data show the mean  $\pm$  SEM  $\alpha$ -glucosidase inhibitory effect of extract at different concentrations, n=3. Values with different superscripts within rows are significantly different ( $p < 0.05$ , one-way analysis of variance followed by the Student–Newman–Keuls' test); Results with separate superscripts within rows are significantly different ( $p < 0.050$ ), while those that are alike are comparable ( $p > 0.05$ ).

**Figure 1. Dose related antihyperglycaemic effect of extract in rats**



Data shows the mean  $\pm$  SEM levels of blood glucose at the various period of time represented in percentage, n=5. Results having separate superscripts are significantly different ( $p < 0.050$ ), while those that are similar are comparable ( $p > 0.050$ ): One-way variance analysis (ANOVA) then Student-Newman-Keul's test. **GLU** (10 g/kg): Glucose 10 g/kg; **SXCS (25-200)**: *S. biafrae*, *X. aethiopica*, *C. papaya*, *S. mombin* extract **GLI**: Glibenclamide ( 5.0 mg/kg).

**Figure 2. Effect of the extract on streptozotocin-induced diabetic rats**



Data show the Mean  $\pm$  SEM level of blood glucose at the various period of time expressed as percentage, n=5. Results having separate superscripts are significantly different ( $p < 0.050$ ), while those that are alike are comparable ( $p > 0.050$ ): One-way variance analysis (ANOVA) then Student-Newman-Keul's test. **DW**: Distilled Water; **SXCS (50, 100)**: *S. bialfrae*, *X. aethiopica*, *C. papaya*, *S. mombin* extract, **GLI (5)**: Glibenclamide (5.0 mg/kg).

**Table 6: Antioxidant activity of the extract**

Extract/Drug	IC <sub>50</sub> (mg/mL)	(mg GAEq/g)	(mg REq/g)	(mgAAEq/g)		IC <sub>50</sub> (mg/mL)
	DPPH	TPC	TFC	TAC	FRAP	HRSA
<b>HRE</b>	0.341	54.7 $\pm$ 0.04	69.5 $\pm$ 1.4	30.1 $\pm$ 1.7	17.41 $\pm$ 0.7	0.368
<b>Ascorbic Acid</b>	0.113					0.168

Data show the mean  $\pm$  SEM (n=3). IC<sub>50</sub> concentration needed to give 50% activity; mgAAEq/g: mg Ascorbic acid equivalent per g; mgGAEq/g: mg Gallic acid equivalent per g; mg REq/g: mg Rutin equivalent per g. DPPH: (1,1-diphenyl-2-picrylhydrazyl) radical scavenging activity assay; FRAP: Ferric Reducing Antioxidant Power; TAC: Total Antioxidant Capacity; HRSA: Hydroxyl Radical Scavenging Activity; TPC: Total Phenolic Content; TFC: Total Flavonoid Content

The overall blood glucose level reduction elicited by the combination plant extract in this study could be adduced to both its  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitory (extra pancreatic) activities (Tables 4 and 5), insulin stimulation (Fig. 1 and 2) as well as free radical scavenging activity (Table 6). *Senecio biafrae*, *Carica papaya*, *Xylopi aethiopica* and *Spondias mombin* had been reported to exert their anti-hyperglycaemic effects through extra pancreatic and insulin stimulating mechanisms of action with additional anti-oxidant effects [18,21,23,25,31,69]. Furthermore, the anti-hyperglycaemic and antioxidant activities elicited by the combined plant extract were significantly higher than those given by each of the component plants.

### 3. CONCLUSION

The results of this study confirmed that the combination of *Senecio biafrae* leaf, *Xylopi aethiopica* fruit, *Carica papaya* seed and *Spondias mombin* stem bark mixed together in equal ratio had significant antidiabetic activity at lower doses than the individual plants without toxic effects on glucose levels, haematological and biochemical components of animal blood samples. It also confirmed that the combined plant extract will be more beneficial in the management of diabetes mellitus than the component plants and could therefore be used as an antidiabetic recipe.

### CONSENT

#### ETHICAL APPROVAL

Ethical clearance number IPH/OAU/12/1758 was issued by the Health Research Ethics Committee, Institute of Public Health, Obafemi Awolowo University on the use of animals for this work.

This is not applicable to this study.

### REFERENCES

1. World Health Organization(WHO). (2016). *Global Report on Diabetes*, p 88.
2. American Diabetes Association (ADA) (2021). Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes. *Diabetes Care*, 44(Suppl. 1), S15-S33.
3. World Health Organization. (2021b). The Global Diabetes Compact. pp. 1-7.
4. International Diabetes Federation. *Diabetes Atlas*. 3rd ed. Belgium: B-1000 Brussels, 19 Avenue Emile de Mot; 2006;387.
5. Faloye KO, Ayoola MD, Amos-Tautua BM, and Famuyiwa SO. Anti-diabetic Activity of Convallatoxin Isolated from the Root Bark of *Parquetina nigrescens* (Afzel.) Bullock (Asclepiadaceae). *European Journal of Medicinal Plants*, 2018; 25(4): 1-9.
6. Ayoola MD, Oriola AO, Faloye KO, and Aladesanmi AJ. Justifying the Antidiabetic Ethnomedicinal Claim of *Massularia Acuminata* Through Its Antihyperglycaemic Activity. *Am J Biomed Sci & Res*,2020; 8(2): 76-258.
7. Ajileye AJ, Ayoola MD, Elujoba AA, Akinwunmi KF. Antihyperglycaemic and Antioxidant

Activities of *Sansevieria liberica* as justification for its antidiabetic claims. *Afr. J. Pharm. Pharmacol*, 2020; 14(3): 59-66.

8. Che CT, Wang ZJ, Chow MS, Lam CW. Herb-herb combination for therapeutic enhancement and advancement: theory, practice and future perspectives. *Molecules*, 2013; 18(5): 5125-5141.
9. Kong YC. Huangdi Neijing, A Synopsis with Commentaries; *The Chinese University Press: Hong Kong, China*, 2010.
10. Ogbonnia SO, Mbaka GO, Adekunle A, Anyika EN, Gbolade OE, Nwakakwa N. Effect of a poly-herbal formulation, Okudiabet, on alloxan-induced diabetic rats. *Agric Biol J North Am*, 2010; 1:139-145.
11. Akinola OB, Omotoso GO, Akinola OS, Dosumu OO, Adewoye ET. Effects of combined leaf extract of *Vernonia amygdalina* and *Azadirachta indica* on hepatic morphology and hepatotoxicity markers in streptozotocin-induced diabetic rats. *PubMed*, 2011; 9: 1373-1379.
12. Ejike CE, Awazie SO, Nwangozi PA, Godwin CD. Synergistic postprandial blood glucose modulatory properties of *Vernonia amygdalina* (Del.), *Gongronema latifolium* (Benth.) and *Occimum gratissimum* (Linn.) aqueous decoctions. *J. Ethnopharmacology*, 2013; 149: 111-116.
13. Tiwari BK, Kumar, D, Abidi AB, Rizvi SI. Efficacy of composite extract from leaves and fruits of medicinal plants used in traditional diabetic therapy against oxidative stress in alloxan-induced diabetic rats. *ISRN Pharmacol*, 2014: 608590.
14. Etuk EU, Mohammed BJ. Informant consensus selection method: A reliability assessment on medicinal plants used in north western Nigeria for the treatment of diabetes mellitus. *African Journal of Pharmacy and Pharmacology*, 2019; 3(10): 496-500.
15. Shokoohi R, Kianbakht S, Faramarzi M, Rahmanian, M, Nabati, F, Mehrzadi, S. et al. Effects of an Herbal Combination on Glycemic Control and Lipid Profile in Diabetic Women: A Randomized, Double-Blind, Placebo-Controlled Clinical Trial. *Journal of Evidence-Based Complementary & Alternative Medicine*, 2017; 22(4): 798-804.
16. Schippers RR. African indigenous vegetables. An overview of the cultivated species. Chatham, United Kingdom: Natural Resources Institute/ACP-EU Technical Centre for Agricultural and Rural Cooperation. 2000.
17. Gbadamosi IT, and Okolosi O. Botanical galactogogues: Nutritional values and therapeutic potentials. *J. Appl Biosci*, 2013; 61: 4460- 4469.
18. Gbolade AA. Inventory of antidiabetic plants in selected districts of Lagos State, Nigeria. *J. Ethnopharmacol*, 2008; 121(1): 135-139.
19. Ajiboye BO, Ojo OA, Okesola MA, Akinyemi AJ, Talabi JY, Idowu OT, et al. *In-vitro* antioxidant activities and inhibitory effects of phenolic extract of *Senecio bialfrae* (Oliv and Hiern) against key enzymes linked with type II diabetes mellitus and Alzheimer's disease. *Food Sci Nutr*, 2018; 6: 1803–1810.
20. Ajiboye BO, Edobor G, Ojo AO, Onikanni S. A., Olanwaju, O. I., and Muhammad, N. O. (2014). Effect of aqueous leaf extract of *Senecio bialfrae* on hyperglycaemic and serum lipid profile of alloxan-induced diabetic rats. *International Journal of Diseases and Disorders*, 2(1), 59-64.

21. Ayoola MD, Adebajo AC, Zotor FB, Pinkoane MG. Justifying antidiabetic ethnomedicinal claim of *Senecio bialfrae* through its antihyperglycemic and antioxidant activities. *Ann Complement Altern Med*. 2019;1(2):1006.
22. Sidiq LO, Segun PA, Ogbole OO. Medicinal Plants Used in Four Local Government Areas of Southwestern Nigeria for the Management of Diabetes and Its Comorbidities: An Ethnobotanical Survey. *Research Square*, 2020; 1-17.
23. Famuyiwa FG, Ayoola, MD, Famuyiwa SO, Aladesanmi AJ. (2018). Hyperglycaemia Lowering Effect of Kaurane Diterpenoids from the Fruits of *Xylopiya aethiopyca* (A. Dunal) Rich. *International Journal of Medicinal Plants and Natural Products (IJMPNP)*, 4(3), 2018; 11-19.
24. Choumessi AT, Danel M, Chassaing S, Truchet I, Penlap VB, Pieme AC. et al. Characterization of the antiproliferative activity of *Xylopiya aethiopyca*. *Cell Div*, 2012; 7(1), 8.
25. Nwozo SO, Orojobi BF, Adaramoye OA. Hypolipidemic and antioxidant potentials of *Xylopiya aethiopyca* seed extract in hypercholesterolemic rats. *J Med Food*, 2011; 14(1-2): 114-149.
26. Akinwunmi KF, and Ayoola, MD. Anti-hyperglycaemic, anti-inflammatory and anti-oxidant activities of *Carica papaya* and *Citrullus lanatus* seeds. *Ife Journal of Science*, 2018; 20(2), 207-218.
27. Malathi P, Vasugi SR. Evaluation of mosquito larvicidal effect of *Carica papaya* against *Aedes aegypti*. *Int. J. Mosq. Res*, 2015; 2(3), 21-24.
28. Oduola T, Idowu TO, Bello IS, Adeniyi F, Ogunyemi E. Haematological response to intake of unripe *Carica papaya* fruit extract and the isolation and characterization of caricapinoside: A new antisickling agent from the extract. *Asian J. Pharm. Clin. Res*, 2012; 5(Suppl 3), 77-81.
29. Ogbole OO, and Ajaiyeoba EO. Traditional Management of Tuberculosis in Ogun State of Nigeria: The Practice and Ethnobotanical Survey. *Afr. J. Trad. CAM*, 2010; 7(1), 79 - 84.
30. Mohammed A, Islam MS. Antioxidant potential of *Xylopiya aethiopyca* fruit acetone fraction in a type 2 diabetes model of rats. *Biomed Pharmacother*, 2017; 96; 30-36.
31. Gbolade A, Ekor M, Akinlolu A, and Ayoola, M.D (2009). Antidiabetic Activity of Ethanolic extracts of *Spondias mombin* (Anacardiaceae) stem bark on Alloxan-induced Diabetic Rats. *African Journal of Traditional, Complementary and Alternative Medicines*, 2009; 6: 437.
32. OECD. (2001). The OECD guidelines for the testing of chemicals: 423. Acute Oral Toxicity – Acute Toxic Class Method
33. OECD. (2008). The OECD guidelines for the testing of chemicals: 407. Repeated dose 28-day oral toxicity study in rodents.
34. Sharma H, Tun-Abraham ME, and McAlister V. 'Bidirectional graft-host hematological traffic in liver transplantation', *Hepatobiliary Surgery and Nutrition*, 2019; 8: 253.
35. Bahman N, Leyla A, and Hamidreza I. Alpha-amylase inhibitory activities of six *Salvia spp.* *Iranian J. Pharm. Res*, 2008; 7: 297-303.

36. Jain C, Kumar P, Singh A, and Jindal A. *In vitro* comparisons of anti-diabetic activity of drugs used for the treatment of diabetes. *Journal Clinical Biochemistry Nutrition*, 2013; 40: 163–167.
37. Li T, Zhang XD, and Song YW. A microplate-based screening method for alpha-glucosidase inhibitors. *Chinese Chin J Clin Pharmacol Ther*, 2005; 10: 1128-1134.
38. Adebajo AC, Iwalewa EO, Obuotor EM, Ibikunle GF, Omisore NO, Adewunmi CO, et al. Pharmacological properties of the extract and some isolated compounds of *Clausena lansium* stem bark: Antitrichomonal, anti-diabetic, antiinflammatory, hepatoprotective and antioxidant effects. *J Ethnopharmacol*. 2009;122(1): 10-19.
39. Adebajo AC, Ayoola MD, Odediran SA, Aladesanmi AJ, Schmidt TJ, Verspohl EJ. Evaluation of ethnomedical claims III: Antihyperglycaemic activities of *Gongronema latifolium* root and stem. *J Diabetes*. 2013a;5(3):336-343.
40. Adebajo AC, Ayoola MD, Obagbemi OR, Obuotor EM, Ogunsina MO, Verspohl EJ. Antihyperglycaemic and antioxidant activities of *Eugenia uniflora* leaf: evaluation of ethnomedical claims IV. *Ife J Sci Technol*. 2013b;1(1):1-18.
41. Ayoola MD, Adebajo AC, Obuotor EM, Oladapo TO, Fleischer TC. Antihyperglycaemic and anti-oxidant activities of five Nigerian antidiabetic plants. *J Sci Technol. (Ghana)*. 2017;37(2): 71-84.
42. Brand-Williams W, Cuvelier, ME, Berset, C. Use of a free radical method to evaluate antioxidant activity. *LWT- Food Science and Technology*, 1995; 28, 25-30.
43. Prieto P, Pineda M, Aguilar M. Spectrophotometric quantification of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. *Anal Biochemistry*, 1999; 269(2):337-41.
44. Benzie IFF, and Strain JJ. Ferric reducing (antioxidant) power as a measure of antioxidant capacity: the FRAP assay. *Methods Enzymol.*, 1999; 299:15-36.
45. Ferrer-Sueta G, and Radi R. Chemical biology of peroxynitrite: kinetics, diffusion, and radicals. *ACS Chemical Biology*, 2009; 4(3):161-77.
46. Zhilen J, Mengcheng T, Jianming W. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chemistry*, 1999; 64: 555-559.
47. Miliauskas G, Venskutonis PR, Van Beek TA. Screening of radical scavenging activity of some medicinal and aromatic plant extract. *Food Chemistry*, 2004; 88: 231-237.
48. Singleton VL, Rudolf O, Lamuela-Raventos RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*. 1999; 299:152-178.
49. Gulcin I, Sat, GI, Beydemir S, Elemastas M, Kfrevioglu OI. Comparison of antioxidant activity of clove (*Eugenia caryophyllata* Thumb) buds and lavender (Lavender stoechas L.). *Food Chemistry*, 2003; 87: 393-400.
50. Halim SZ, Abdullah NR, Afzan A, Abdul Rashid BA, Jantan I, and Ismail Z. Acute toxicity study of *Carica papaya* leaf extract in Sprague Dawley rats. *Journal of Medicinal Plants Research*, 2011; 5(20): 1867-1872.
51. Ayodele PF, Ore A, Akinloye OA. Median Lethality Dose of *Xylopiya aethiopicum* Fruit Ethanol Extract. *J Anal Tech Res*, 2019; 1(1): 33-36.

52. Powal M, Gautam SK, Khan NA, and Maheshwari KK. Evaluation of Toxicity and Antihyperlipidemic Activity of *Spodias mombin* Leaves Methanolic Extract in Laboratory Rats. *Cardiovasc Hematol Disord Drug Targets*, 2020;20(4): 289-296.
53. Ayoola MD, Akinwunmi KF, and Agboola OB. Anti-diabetic and antioxidant activities of *Entandrophragma cylindricum* and *Triclisia subcordata*. *Nigerian Journal of Natural Products and Medicine*, 2017b; 21: 24-31.
54. Ayoola MD, Balogun, JO, Famuyiwa FG, Yeboah SO, Famuyiwa SO. Isolation and characterization of 2-hydroxy-3-[4-hydroxyphenyl]-2-propenoic acid and 4-bromophenol from anti-diabetic extract of the root bark of *Uvaria afzelii*. *South African Journal of Botany*, 2017c; 112:527–532.
55. Sunmonu TO, and Oloyede OB. 'Performance and haematological indices in rats exposed to monocrotophos contamination'. *Human & experimental toxicology*, 2010;29: 845-50.
56. Nwinuka NM, Michael OM, and Nwiloh BI. 'Effects of aqueous extract of *Mangifera indica* L.(Mango) stem bark on haematological parameters of normal albino rats', *Pakistan Journal of Nutrition*, 2008; 7: 663-66.
57. Bashir L, Shittu OK, Rotimi AA, Olalekan IA, Kamooru AA, and Ossai PC. Effect of Methanol Extract of *Telfairia occidentalis* on Haematological Parameters in Wister Rats. *J. Med. Sci.*, 2015: 15 (5):246-250.
58. Brautbar N, and Williams J. 'Industrial solvents and liver toxicity: risk assessment, risk factors and mechanisms'. *International Journal of Hygiene and Environmental Health*, 2002; 205: 479-91.
59. Desai SN, Patel DK, Devkar RV, Patel PV, and Ramachandran AV. Hepatoprotective potential of polyphenol rich extract of *Murraya koenigii* L.: An *in vivo* study. *Food and chemical toxicology*, 2012; 50(2): 310-314.
60. Lee SS, Yu S, and Withers SG. Detailed Dissection of a New Mechanism for Glycoside Cleavage:  $\alpha$ -1,4-Glucan Lyase. *Biochemistry*, 2003; 42(44): 13081–13090.
61. Ngugi M, Njagi J, Kibiti C, and Mwenda M. Pharmacological Management of Diabetes Mellitus. *Asian Journal of Biochemical and Pharmaceutical Research*, 2012; 2: 2012.
62. Derosa G., and Maffioli P. Mini-special issue paper management of diabetes patients with hypoglycaemic agents  $\alpha$ -Glucosidase inhibitors and their use in clinical practice. *Arch. Med. Sci.*, 2012; 5: 899-906.
63. Rosak C, and Mertes. Critical evaluation of the role of acarbose in the treatment of diabetes: patient considerations. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 2012; 1, 357.
64. Verspohl EJ, Bauer K, Neddermann E. Antidiabetic effect of *Cinnamomum cassia* and *Cinnamomum zeylanicum* *in vivo* and *in vitro*. *Phytother Res.* 2005;19(3):203- 206.
65. Verspohl EJ. Recommended testing in diabetes research. *Planta Med.* 2002; 68(7):581-590.
66. Kar A, Choudhary BK, Bandyopadhyay NG. Preliminary studies on the inorganic constituents of some indigenous hypoglycaemic herbs on oral glucose tolerance test. *J Ethnopharmacol.* 1999; 64(2):179-184.
67. Luzi L, Pozza G. Glibenclamide: an old drug with a novel mechanism of action? *Acta Diabetol.* 1997;34(4):239- 244.

68. Ishola IO, Ikuomola BO, and Adeyemi OO. Protective role of *Spondias mombin* leaf and *Cola acuminata* seed extracts against scopolamine-induced cognitive dysfunction. *Alexandria Journal of Medicine*, 2018; 54(1):27–39.
69. Cabral B, Siqueira EMS, Bitencourt MAO, Lima MC, Lima AK, Ortmann CF et al., “Phytochemical study and anti-inflammatory and antioxidant potential of *Spondias mombin* leaves.” *Revista Brasileira de Farmacognosia*, 2016; 26(3): 304–311.

UNDER PEER REVIEW