

Association of morphological and biochemical traits by different treatments of beetroot (*Beta vulgaris* L.) cv. crimson globe under alkaline conditions of Telangana, India.

ABSTRACT

Beet root is one of the most important root crops with health promoting substances like vitamins, minerals and betalains, grown in northern and southern parts of India. The present study was carried to assess the morphological and biochemical characters by using principal component analysis over eight treatments and twenty-one parameters under alkaline conditions where the pH of soil is around 9.5 in mojerla, Telangana, India. Principal component analysis accounted 81.57% variation among eight treatments of beet root grown under alkaline conditions. Root yield was observed positive significant with height of the plant, number of leaves/plant, leaf area chlorophyll content index, length of the root, root diameter, root yield/plant, root yield per hectare, root to shoot ratio, harvest index, TSS, ascorbic acid content, betanine content, nitrogen, phosphorus and potassium content (in the soil following crop harvest) and organic carbon (in the soil following crop harvest) whereas negative significant correlation with number of days required for 80 % germination, pH and EC. From this experiment it is concluded that increasing in pH and Ec decreasing the yield and yield attributing traits under alkaline conditions of mojerla, Telangana.

Key words: Correlation, Principal component analysis, traits, pH and Electrical Conductivity.

Introduction

Alkaline soil are those soils having high pH and a wide range of electrical conductivity congruity with (39). In addition to salt toxicity, these soils also contain high levels of CO_3^{2-} and HCO_3^- all of which are detrimental to plant development (24) these salt-affected soils cover around 7% of the earth's land surface, and sodium-affected soils are even more common. Salt components of alkaline soils include cations and anions (36). SO_4^{2-} , Cl^- and Na^+ are main ions present naturally in alkaline soils (34). With increase in pH, the availability of plant nutrients like iron, copper, boron, manganese decreased and molybdenum increased (22). Reduced ability of root to absorb nutrients and photosynthetic activity of plant is decreased are the symptoms observed in beet root under alkaline conditions. Present experiment shows the relationship between yield and yield attributing traits under alkaline soils of mojerla, Telangana, India.

Beet root belongs to the family Chenopodiaceae (14). It is one of the important root crops originated in Mediterranean region cultivated throughout Europe. It is mainly used as food coloring agent (6). Beet root is having betacyanin which is composed of betanin's and isobetanins (11) which impart reddish to purple pigment having the properties to obviate cancer and cardiovascular diseases by increasing the low-density lipoproteins to oxidation by reducing the oxidative effect of lipids (30, 35, 8). Sucrose is the primary sugar in beetroot, with minor amounts of glucose and fructose. (3) Juice of beetroot has capacity to reduce blood pressure and protect from liver diseases (7, 19) and it is also best-known anti-inflammatory properties (39, 15). In India major beet root crop growing states are Haryana, Uttar Pradesh, Himachal Pradesh, West Bengal and Maharashtra. Beetroot is grown on an area of 0.079 lakh hectares in India, with an annual production of 1.51 lakh million tonnes, and Telangana has 0.04 lakh hectares with an annual production of 0.11 lakh million tonnes. (13).

Day by day world population is increasing in log face. To feed the growing population we have to cultivate the crops according to the soil and climatic conditions. In these experiment association of different traits of beetroot under alkaline condition is conducted. The most significant characteristic, yield, is controlled by a variety of traits and its interaction. The link between the variables is understood through correlation analysis (4). Without mentioning cause and effect, the correlation coefficient provides information about the relationship among the independent and dependent variables (1). A. From these experiment relationship for the yield and yield attributing traits can be analyzed. B. Variability among the treatments is ascertained using principal component analysis (13).

MATERIALS AND METHODS

The current study was carried out at the College of Horticulture, Mojerla, SKLTSH University, Mulugu, Telangana State, India, between the last week of October 2019 and the second week of January 2020. The farm has a semi-arid climate and is located at an elevation of 401 meters above mean sea level on 770.96' East longitude and 160.36' North latitude. pH of soil is 9.5, EC is 0.18, Organic carbon, Available Nitrogen, Phosphorus and potassium are 0.35 %, 252.11, 29.2 and 262.3 kg/ha respectively before conducting the experiment.

All agronomic practices are done and plant protection measures were followed to raise a healthy crop.

Fertilizer Application

Organic manures viz., well decomposed farmyard manure @ 4.8 kg/plot. Vermicompost @ 1.2 kg per plot were incorporated in to the respective experimental plots uniformly, before sowing as basal

application. Similarly, N, P and K @ 28:44:28kg/ha were applied in the form of urea, single super phosphate and murate of potash respectively. Urea was applied in two splits, the first dose as basal application and another dose at 25 days after sowing. The entire dose of single super phosphate and murate of potash were applied at the time of sowing as basal dose. Before sowing, the seeds were treated with each of Azotobacter and PSB @ 10kg/ha.

During the experiment, correlation and PCA analysis was carried for the following parameters to fulfill the objectives under the study of southern Telangana, India. For that mean values are taken for each parameter under different treatments mentioned in the below table. By using Statistical Package for the Social Science (SPSS) correlation coefficient analysis is done. The statistical tools used WINDOSTAT 9.1, created by INDOSTAT Services Ltd., Hyderabad, India.

These are the parameters taken into consideration Number of days required for 80 % germination, height of the plant, number of leaves/plants, leaf area chlorophyll index, root length, root diameter, root yield/plant, root yield /ha, root to shoot ratio, harvest index (HI), TSS, ascorbic acid content, betanine content, nitrogen, phosphorus, potassium content, pH, EC and organic carbon in the soil following the crop harvest and five plants were randomly selected and data is recorded. The degree of association of character root yield and among themselves are calculated by Correlation coefficients.

Treatment details
75 % RDNPK + Farm Yard Manure (12 t/ha) + Azotobacter (10 kg/ha) + PSB (10 kg/ha)
75 % RDNPK + Vermicompost C (3 t/ha) + Azotobacter (10 kg/ha) + PSB (10kg/ha)
75 % RDNPK + Farm Yard Manure (6 t/ha) + Vermicompost (1.5 t/ha) + Azotobacter (10 kg/ha) + PSB (10 kg/ha)
50 % RDNPK + Farm Yard Manure (12 t/ha) + Azotobacter (10 kg/ha) + PSB (10 kg/ha)
50 % RDNPK + Vermicompost (3 t/ha) + Azotobacter (10 kg/ha) + PSB (10 kg/ha)
50 % RDNPK+ Farm Yard Manure (6 t/ha) + Vermicompost (1.5 t/ha) + Azotobacter (10 kg/ha) + PSB (10 kg/ha)
75 % RDNPK + Farm Yard Manure (12 t/ha)
50 % RDNPK + Farm Yard Manure (12 t/ha)

RESULTS AND DISCUSSION

Correlation Analysis

In order to understand how numerous characteristics interacted and were connected to the root yield, correlation analysis is used. Table 2 shows that all correlations are significant at the 1% level.

In present experiment a significant positive correlation between root yield (kg plot^{-1}) with plant height (0.9090 cm). (37) in garlic who reported that improvement in plant height and leaf number as they have physiological capacity to mobilize and translocate photosynthates to organ of economic value which in turn might have increased the yield. congruity with (28).

Number of leaves per plant (0.6619). According to (32) in garlic it might be due to increase in photosynthate assimilates with increasing the leaf number, led to higher total yield, leaf area (0.9448 cm^2) (12). (5) reported that similar association of characters between plant height, number of leaves, leaf area, bulb diameter, bulb yield and TSS were observed in garlic.

Chlorophyll content index at (0.9679). Similar results were also reported by (16) who reported that chlorophyll content index showed positive association with plant height at 30 and 60 days, foliage yield, leaf area and ascorbic acid content, root length (0.9265 cm), root diameter (0.9640 cm), root yield/plant (0.9528 g), root yield/hectare (0.9973 t). Similar association is seen according to (31, 30, 26). Inference of investigation revealed that, the capacity of the plant to produce more photosynthates, which are mobilized to the organ of economic value, could be improved by the timely application of Nitrogen fertilizer this would increase root weight and root diameter, which would ultimately result in a higher yield per unit area, as reported by (12).

Root to shoot ratio (0.9198), harvest index (0.8594). According to (27) in beet root there was similar results of direct association of root to shoot ratio with harvest index and TSS, while indirect association with root diameter and root yield, ascorbic acid content mg/100g (0.9186) similar results are reported by (2, 21),

Betanine, content mg/100g (0.5753) TSS (0.9934 $^{\circ}\text{Brix}$) (5) reported similar positive association of TSS with plant height, number of leaves per plant, leaf area, bulb diameter and bulb yield in garlic. Nitrogen, phosphorus and potassium content kg/ha in the soil following the crop harvest (0.9858, 0.4708, 0.8469 respectively) and organic carbon (%) in the soil following the crop harvest (0.9493).

Root yield was negatively correlation with number of days required for 80 % germination (-0.7287 days), pH in the soil after the harvest of the crop (-0.9660). With increase in pH the nutrients present in the soil are unavailable to roots and it also led to deficiency of iron resulting in chlorosis. (10) shown similar results of negative association of pH with chlorophyll content in tomato and Electric conductivity (dS mol^{-1}) in the soil after the harvest of the crop (-0.9429 these results are in congruity with (20)

The characters like root yield were observed positively significant with height of the plant, number of leaves/plants, leaf area, chlorophyll content index, root length, root diameter, root yield/plant, root yield/hectare, root to shoot ratio, harvest index (HI), TSS, ascorbic acid content, betanine content nitrogen, phosphorus and potassium content and organic carbon in the soil following the crop harvest. Selection above traits is desirable in selection among beet root genotypes in further breeding programmes. Similar observations are recorded by (17,18). 27 reported in beet root genotypes, positive correlation was observed between root yield with shoot length, shoot weight, root girth, root volume.

Principal component analysis (PCA)

Table 1 and figure 1 presents the principal component analysis (PCA) findings. The results of the principal component analysis on the correlation matrix for the eight treatments and 21 parameters used to study morphological and biochemical characters include the eigenvalues, percent of variability, cumulative percent of variability, and factor loading values for the various traits for the respective principal components shown in table 1 and fig 1. According to principal component analysis, significant principal components had eigen values greater than one, whereas important principal components had eigen values less than one. During analysis two principal components (PCs) exhibited more than 1.00 Eigen value, and showed about 81.57% variability among the trait studied. These results are according with the (23).

As per the analysis, the first components accounted for most of the variation. The first component F1 explained 81.57% of total variation and more divergent with leaf area, chlorophyll, root length, root diameter, root yield (kg/plot), Root yield (t/ha), Nitrogen and Total soluble solids. Similar results were obtained by (9) reported results may help in technical improvement of the genotype characteristics which might be used for further selection in breeding program. From the results it is stated that traits which are showing significant contribution to the observed variation is used for future breeding program. The treatments which are showing variation among the traits should be given most importance. to increase the quality of future segregant selection regarding parent selection during hybridization programmes. Results are congruity with (25) in tomato (33) in mango. It is necessary to know the best treatment and traits whose contribution is the most for

the variation so as to improve that character in the further breeding programs

Conclusion

Experimental result shows that root yield was observed in positive association with height of the plant, number of leaves/plants, chlorophyll content index, leaf area, length of root, root diameter of root, root yield/plant, root yield/hectare, root to shoot ratio, harvest index (HI), TSS, ascorbic acid content, betanine content nitrogen, phosphorus and potassium content in the soil following the crop harvest and organic carbon in the soil following the crop harvest. In future for breeding purpose and or increasing the yield, the traits which are directly associated can be selected.

UNDER PEER REVIEW

References

1. Akintunde A. Path analysis step by step using excel. *Journal of Technical science and Technologies*. 2012 May 30;1(1): 09-15.
2. Ashish K. Variability and Correlation Study in Tomato (*Solanum lycopersicum* L.). *International Journal of Agriculture Sciences*, ISSN. 2017:0975-3710.
3. Bavec M, Turinek M, Grobelnik-Mlakar S, Slatnar A, Bavec F. Influence of industrial and alternative farming systems on contents of sugars, organic acids, total phenolic content, and the antioxidant activity of red beet (*Beta vulgaris* L. ssp. *vulgaris* Rote Kugel). *Journal of Agricultural and Food Chemistry*. 2010 Nov 24;58(22):11825-31.
4. Bizeti HS, Carvalho CG, Souza JR, Destro D. Path analysis under multicollinearity in soybean. *Brazilian archives of biology and technology*. 2004; 47:669-76.
5. Chotaliya P, Kulkarni GU. Character association and path analysis for quantitative traits in garlic (*Allium sativum* L.). *International Journal of Current Microbiology and Applied Sciences*. 2017;6(8):175-84.
6. Clifford T, Howatson G, West DJ, Stevenson EJ. The potential benefits of red beetroot supplementation in health and disease. *Nutrients*. 2015 Apr;7(4):2801-22.
7. Coles LT, Clifton PM. Effect of beetroot juice on lowering blood pressure in free-living, disease-free adults: a randomized, placebo-controlled trial. *Nutrition journal*. 2012 Dec;11(1):1-6.
8. Delgado-Vargas F, Jimenez AR, Paredes-Lopez O. Natural pigments: carotenoids, anthocyanins, and betalains—characteristics, biosynthesis, processing, and stability. *Critical reviews in food science and nutrition*. 2000 May 1;40(3):173-289.
9. Diel MI, Lucio AD, Tartaglia FD, Tischler AL, Lambrecht DM, Zemolin JA, Marques LE. New insights on the influence of the quality of tomato seedlings on production of fruits cultivated in substrates. *Ciencia Rural*. 2022;52.
10. Etissa, E., Dechassa, N., Alamirew, T., Alemayehu, Y. and Desalegn. (2014). Growth and physiological response of tomato to various irrigation regimes and integrated nutrient management practices. *African journal of agricultural research*. 9(19), 1484-1494.
11. Frank T, Stintzing FC, Carle R, Bitsch I, Quaas D, Straß G, Bitsch R, Netzel M. Urinary pharmacokinetics of betalains following consumption of red beet juice in healthy humans. *Pharmacological Research*. 2005 Oct 1;52(4):290-7.
12. Gateri M, Nyankanga R, Ambuko J, Muriuki A. Growth, yield and quality of onion (*Allium cepa* L.) as influenced by nitrogen and time of topdressing. *International Journal of Plant & Soil Science*. 2018; 23(3):1-3.

13. HAPIS (Horticulture Area and Production Information System), Final advanced estimate of area and production of horticultural crops. 2019-2020. <https://aps.dac.gov.in>.
14. Jeffrey C. An introduction to plant taxonomy. CUP Archive; 1982 Aug 19.
15. Jurgen, W., Gundula, W., Stefan, H., Pinar, U., Peter, L. and Ulrike M. (2015). Compositional characteristics of commercial beetroot products and beetroot juice prepared from seven beetroot varieties grown in Upper Austria. *Journal of Food Composition and Analysis*. 42:46-55.
16. Kassambara A. Practical guide to principal component methods in R: PCA, M (CA), FAMD, MFA, HCPC, factoextra. Sthda; 2017 Aug 23.
17. Kathayat K, Kushwaha ML, Rawat M. Assessment of Genetic Variability, Correlation and Path Coefficient Analysis In Taro (*Colocasia Esculenta* L. Schott. Var. Antiquorum) Germplasm. *Think India Journal*. 2019 Aug 22;22(16):1068-81.
18. Katiyar SK, Singh HC, Gangwar RS, Verma MC. and Singh RK. (2014). Genetic variability and selection parameters in pea (*Pisum sativum* L.). *Bioscience*. 2014;7 (9): 720-723.
19. Katoch V, Singh P, Devi MB, Sharma A, Sharma GD, Sharma JK. Study of genetic variability, character association, path analysis and selection parameters for heterotic recombinant inbred lines of garden peas (*Pisum sativum* var. hortense L.) under mid-hill conditions of Himachal Pradesh, India. *Legume Research*. 2016 Apr 1;39 (2):163-9
20. Krajka-Kuzniak V, Szaefer H, Ignatowicz E, Adamska T, Baer-Dubowska W. Beetroot juice protects against N-nitrosodiethylamine-induced liver injury in rats. *Food and chemical toxicology*. 2012 Jun 1;50(6):2027-33.
21. Kumar SR, Arumugam T, Premalakshmi V. Evaluation and variability studies in local types of brinjal for yield and quality (*Solanum melongena* L.). *Electronic Journal of Plant Breeding*. 2012 Dec 31;3(4):977-82.
22. Lavanya KS, Srinivasa V, Ali S, Lakshmana D, Kadian MS. Correlation and path analysis for yield and yield-related traits of potato (*Solanum tuberosum* L.) in Karnataka. *National Academy Science Letters*. 2020 Apr;43(2):137-40.
23. Nkebiwe PM, Weinmann M, Bar-Tal A, Müller T. Fertilizer placement to improve crop nutrient acquisition and yield: A review and meta-analysis. *Field crops research*. 2016 Sep 1;196:389-401.
24. Paliwal S, Sharma S, Pathak N. Principal component analysis in mungbean (*Vigna radiata* L. Wilczek) genotypes under two seasons. *The Pharma Innovation Journal*.

- (2022). 11(3):1500-1505.
25. Piovan MJ, Zapperi GM, Pratolongo PD. Seed germination of *Atriplex undulata* under saline and alkaline conditions. *Seed Science and Technology*. 2014 Aug 1;42(2):286-92.
 26. Prakash MO, Vijay B. Principal component and cluster analysis of indigenous tomato genotypes based on morphological indicators. *Res J Biotechnol*. 2017 Jul 1;12(7):50-8.
 27. Sahu K, Sharma PK, Dixit A, Nair SK. Correlation and path coefficient analysis in kharif onion (*Allium cepa* L.) genotypes for Chhattisgarh plains. *Int. J. Curr. Microbiol. App. Sci*. 2018;6:256-63.
 28. Sanghera GS, Singh RP, Kashyap L, Tyagi V, Sharma B. Evaluation of sugar beet genotypes (*Beta Vulgaris* L.) for root yield and quality traits under subtropical conditions. *Journal of Krishi Vigyan*. 2016;5(1):67-73.
 29. Sharma VR, Omotayo K, Malik S, Kumar M. and Sirohi A. Character Association and Path Analysis in Garlic (*Allium Sativum* L.). *International Quarterly Journal of Life science*. 2016. 11(3):1931-1935
 30. Singh B. and Hathan BS. Chemical composition, functional properties and processing of Beetroot. *International Journal of Scientific Engineering and Research*. 2014. 5, 679–684.
 31. Singh DP, Deo C, Singh A, Gautam DK. and Kumar A. Studies on correlation and path coefficient analysis in sweet potato (*Ipomoea batatas* (L) Lam.) germplasm. *Research in Environment and Life Sciences*. 2015. 8(2): 343-346.
 32. Singh M, Yadav GC. Correlation and path coefficient analysis for yield and horticulture traits in different genotypes of *Colocasia* (*Colocasia esculenta* var. *antiquorum* (L.) Schott). *J Pharmac Phytochem*. 2018;3:288-92.
 33. Singh SR, Ahmed NA, Lal S, Amin A, Amin M, Ganie SA, Jan N. Character association and path analysis in garlic (*Allium sativum* L) for yield and its attributes. *SAARC Journal of Agriculture*. 2013;11(1):45-52.
 34. Soujanya B, Kumar AK, Vanisri S, Sreedhar M. Assessment of Genetic Divergence by Using Multivariate Analysis for Physico Chemical Characters of Mango Table and Juicy Cultivars Grown in Telangana Region.
 35. Tavakkoli E, Fatehi F, Coventry S, Rengasamy P, McDonald GK. Additive effects of Na⁺ and Cl⁻ ions on barley growth under salinity stress. *Journal of Experimental Botany*. 2011 Mar 1;62(6):2189-203.
 36. Tesoriere L, Allegra M, Butera D, Livrea MA. Absorption, excretion, and distribution of dietary antioxidant betalains in LDLs: potential health effects of betalains in humans. *The*

- American journal of clinical nutrition. 2004 Oct 1;80(4):941-5.
37. Tobe K, Li X, Omasa K. Effects of five different salts on seed germination and seedling growth of *Haloxylon ammodendron* (Chenopodiaceae). *Seed Science Research*. 2004 Dec;14(4):345-53.
 38. Tsega K, Tiwari A, Woldetsadik K. Genetic variability, correlation and path coefficient among bulb yield and yield traits in Ethiopian garlic germplasm. *Indian Journal of Horticulture*. 2010;64(4b):489-99.
 39. Wruss J, Waldenberger G, Huemer S, Uygun P, Lanzerstorfer P, Müller U, Höglinger O, Weghuber J. Compositional characteristics of commercial beetroot products and beetroot juice prepared from seven beetroot varieties grown in Upper Austria. *Journal of Food Composition and Analysis*. 2015 Sep 1;42:46-55.
 40. Yu J, Wang Z, Meixner FX, Yang F, Wu H, Chen X. Biogeochemical characterizations and reclamation strategies of saline sodic soil in northeastern China. *CLEAN–Soil, Air, Water*. 2010 Nov;38(11):1010-6.




Figure 1: Biplot graphic with two principal components (PC1 and PC2: 88.28 %) for 21 parameters in beet root grown under alkaline conditions.

Table 1: Principal component loadings of 21 parameters in beet root grown under alkaline conditions.

	F1	F2	F3	F4	F5	F6	F7
Eigenvalue	17.1304	1.4081	0.9546	0.6918	0.4032	0.2673	0.1445
Variability (%)	81.5734	6.7052	4.5459	3.2942	1.9200	1.2730	0.6883
Cumulative %	81.5734	88.2786	92.824	96.1187	98.0387	99.3117	100.000
Germination	-0.1635	0.5254	0.1315	-0.2421	-0.3242	0.4202	0.2190
Plant height	0.2128	-0.2971	-0.0300	0.0025	-0.0340	-0.2521	0.7512
Leaves	0.1660	0.1277	-0.4755	0.5933	-0.0629	0.3870	0.1676
leaf area	0.2349	0.1599	0.0016	-0.1225	-0.0706	-0.1160	0.1405
Chlorophyll	0.2331	-0.1038	0.1208	-0.1974	-0.1518	0.0975	0.0833
Rootlength	0.2341	0.1178	0.1471	0.0716	0.1711	0.0742	0.1718
Root diameter	0.2375	0.0355	0.0314	-0.0805	-0.0350	-0.2322	-0.2803
Root yield g/plant	0.2283	-0.1738	0.1635	-0.1617	-0.0479	0.2225	-0.2186
Root yield (KG/Plot)	0.2382	-0.0920	-0.0879	-0.1033	0.0369	0.0381	0.0246

Root Yield(t/HA)	0.2367	-0.1080	-0.0808	-0.1217	-0.0073	0.1554	0.0669
ROOT/SHOOT T	0.2286	0.0975	0.0653	-0.1633	-0.4069	-0.0821	0.0559
Harvest Index	0.2215	0.0493	0.3847	0.0566	-0.1648	0.0614	-0.0502
VITC	0.2252	0.1871	-0.2323	-0.1247	-0.1541	-0.1575	-0.1605
BETANINE	0.1629	0.0298	0.6156	0.5047	0.0567	0.1373	-0.0118
TSS	0.2367	-0.0900	-0.1163	-0.0889	-0.1161	0.1397	-0.0047
Nitrogen	0.2408	0.0169	-0.0035	-0.0436	0.1012	0.0004	0.0651
Phosphorus	0.1372	0.6666	-0.0320	0.0652	0.1780	-0.3594	0.0722
Potassium	0.2006	0.1070	-0.0631	-0.3222	0.6683	0.3825	0.0077
PH	-0.2383	0.0310	-0.0724	-0.0743	-0.1669	0.1147	0.1302
EC	-0.2339	-0.0412	0.1189	-0.1947	-0.0477	0.2326	0.1948
ORGANIC CARBON	0.2273	-0.0545	-0.2332	0.1034	-0.2751	0.1902	-0.2758




Figure 2: No. of days taken for germination, Ph and electrical conductivity is negatively

correlated with all other parameter indicated by blue color.

UNDER PEER REVIEW

Table 2: Correlation coefficient analysis showing relationship of different Characters on root yield of beet root Cv. Crimson Globe.

Variables	germination	Plant height	leaves	leaf area	Chlorophyll	Root length	Root diameter	Root Yield(g/plant)	Root yield (KG/PT)	Root yield (t/ha)	ROOT/SHOOT	Harvest Index
Germination	1	-0.8200	-0.4724	-0.5181	-0.6480	-0.5705	-0.6518	-0.6959	-0.7287	-0.7120	-0.4867	-0.5182
Plant height	-0.8200	1	0.5593	0.8132	0.8941	0.8113	0.8357	0.8619	0.9090	0.9072	0.8076	0.7688
Leaves	-0.4724	0.5593	1	0.6388	0.5243	0.6569	0.6046	0.4962	0.6619	0.6583	0.5740	0.4967
leaf area	-0.5181	0.8132	0.6388	1	0.9346	0.9589	0.9732	0.8834	0.9448	0.9352	0.9709	0.9001
Chlorophyll	-0.6480	0.8941	0.5243	0.9346	1	0.9184	0.9508	0.9842	0.9679	0.9738	0.9519	0.9253
Rootlength	-0.5705	0.8113	0.6569	0.9589	0.9184	1	0.9449	0.8973	0.9265	0.9183	0.9056	0.9420
Root diameter	-0.6518	0.8357	0.6046	0.9732	0.9508	0.9449	1	0.9299	0.9640	0.9500	0.9546	0.9128
Rootyield g/plant	-0.6959	0.8619	0.4962	0.8834	0.9842	0.8973	0.9299	1	0.9528	0.9605	0.8997	0.9164
Rootyield(KG/Plot)	-0.7287	0.9090	0.6619	0.9448	0.9679	0.9265	0.9640	0.9528	1	0.9973	0.9198	0.8594
Root Yield(t/HA)	-0.7120	0.9072	0.6583	0.9352	0.9738	0.9183	0.9500	0.9605	0.9973	1	0.9192	0.8590
ROOT/SHOOT	-0.4867	0.8076	0.5740	0.9709	0.9519	0.9056	0.9546	0.8997	0.9198	0.9192	1	0.9171
Harvest Index	-0.5182	0.7688	0.4967	0.9001	0.9253	0.9420	0.9128	0.9164	0.8594	0.8590	0.9171	1
VITC	-0.5031	0.7443	0.7119	0.9644	0.8655	0.8774	0.9440	0.8112	0.9186	0.9054	0.9344	0.7861
BETANINE	-0.4336	0.5532	0.4087	0.6141	0.6481	0.7759	0.6457	0.6767	0.5753	0.5714	0.6108	0.8646
TSS	-0.6983	0.8953	0.6904	0.9383	0.9678	0.9081	0.9532	0.9501	0.9934	0.9960	0.9331	0.8558
Nitrogen	-0.6661	0.8766	0.6706	0.9750	0.9595	0.9745	0.9790	0.9380	0.9858	0.9783	0.9340	0.9049
Phosphorus	0.0326	0.2519	0.5113	0.7041	0.4185	0.6664	0.6039	0.3340	0.4708	0.4372	0.5986	0.5395
Potassium	-0.4806	0.6536	0.5087	0.8276	0.7913	0.8511	0.8040	0.7940	0.8469	0.8431	0.7145	0.6946
PH	0.7323	-0.8710	-0.6503	-0.9418	-0.9396	-0.9702	-0.9761	-0.9365	-0.9660	-0.9527	-0.8990	-0.9196
EC	0.7105	-0.8330	-0.7763	-0.9356	-0.8764	-0.9315	-0.9610	-0.8556	-0.9429	-0.9233	-0.8877	-0.8488
ORGANIC CARBON	-0.6749	0.8194	0.8049	0.8897	0.8933	0.8529	0.9128	0.8798	0.9439	0.9455	0.8952	0.8007

Table 3: Correlation coefficient analysis showing relationship of different Characters on different nutrients and organic carbon present in beet root

Variables	VITC	BETANINE	TSS	N	P	K	PH	EC	ORGANIC CARBON
Germination	-0.5031	-0.4336	-0.6983	-0.6661	0.0326	-0.4806	0.7323	0.7105	-0.6749
Plant height	0.7443	0.5532	0.8953	0.8766	0.2519	0.6536	-0.8710	-0.8330	0.8194
Leaves	0.7119	0.4087	0.6904	0.6706	0.5113	0.5087	-0.6503	-0.7763	0.8049
leaf area	0.9644	0.6141	0.9383	0.9750	0.7041	0.8276	-0.9418	-0.9356	0.8897
Chlorophyll	0.8655	0.6481	0.9678	0.9595	0.4185	0.7913	-0.9396	-0.8764	0.8933
Root length	0.8774	0.7759	0.9081	0.9745	0.6664	0.8511	-0.9702	-0.9315	0.8529
Root diameter	0.9440	0.6457	0.9532	0.9790	0.6039	0.8040	-0.9761	-0.9610	0.9128
Root yield g/plant	0.8112	0.6767	0.9501	0.9380	0.3340	0.7940	-0.9365	-0.8556	0.8798
Root yield (KG/Plot)	0.9186	0.5753	0.9934	0.9858	0.4708	0.8469	-0.9660	-0.9429	0.9439
Root Yield(t/HA)	0.9054	0.5714	0.9960	0.9783	0.4372	0.8431	-0.9527	-0.9233	0.9455
ROOT/SHOOT	0.9344	0.6108	0.9331	0.9340	0.5986	0.7145	-0.8990	-0.8877	0.8952
Harvest Index	0.7861	0.8646	0.8558	0.9049	0.5395	0.6946	-0.9196	-0.8488	0.8007
VITC	1	0.4470	0.9241	0.9301	0.7086	0.7858	-0.8860	-0.9339	0.9208
BETANINE	0.4470	1	0.5597	0.6576	0.4055	0.4438	-0.7318	-0.6453	0.5322
TSS	0.9241	0.5597	1	0.9726	0.4495	0.8094	-0.9454	-0.9334	0.9683
Nitrogen	0.9301	0.6576	0.9726	1	0.5879	0.8673	-0.9854	-0.9605	0.9204
Phosphorus	0.7086	0.4055	0.4495	0.5879	1	0.5705	-0.5537	-0.6245	0.4539
potassium	0.7858	0.4438	0.8094	0.8673	0.5705	1	-0.8262	-0.7624	0.7089
PH	-0.8860	-0.7318	-0.9454	-0.9854	-0.5537	-0.8262	1	0.9687	-0.9004
EC	-0.9339	-0.6453	-0.9334	-0.9605	-0.6245	-0.7624	0.9687	1	-0.9387

Organic carbon	0.9208	0.5322	0.9683	0.9204	0.4539	0.7089	-0.9004	-0.9387	1
-----------------------	--------	--------	--------	--------	--------	--------	---------	---------	----------

UNDER PEER REVIEW