

Original Research Article

Identification of Garlic Genotypes for Resistant Sources against Purple Blotch Disease

ABSTRACT

Garlic is an herbaceous annual bulbous plant in the family Amaryllidaceae grown for its pungent and edible bulb. Among the various biotic factors hindering the production and productivity, purple blotch play a predominant role in changing the crops morpho-physiological characters thus deteriorating the metabolic activity in turn affecting the yield considerably. Fungicides have been recommended for the control of the disease however, use of fungicides is expensive and not eco-friendly. The best approach to control the disease is by using resistant genotypes. Hence, 37 genotypes were field screened at AINRPOG (All India Network Research Project on Onion and Garlic), MARS (Main Agricultural Research Station), Dharwad against purple blotch disease under artificial epiphytotic conditions during *Kharif*, in year 2019 to identify the resistant sources. The results revealed that twenty genotypes found to be resistant (1-10%) and ten genotypes were immune(0%) however, five genotypes were found moderately resistant (Same here) in reaction. The remaining two genotypes viz., HRG Local and DOGR409 were found moderately susceptible with maximum grade of 3 in (0-5) scale. None of the genotypes showed susceptible or highly susceptible reaction to purple blotch. Breeders can make use of these identified resistant lines in developing high yield purple blotch disease resistant varieties.

Keywords: Genotypes, garlic, artificial screening, Alternaria porri, disease reaction

1. INTRODUCTION

Garlic (*Allium sativum* L.) is one of the most popular spice crops having higher nutritive value than other bulb crops (cite source). All parts of the garlic plant are edible but the fresh leaves and the dried cloves are the parts mostly used in cooking consumed [1]. It is vegetatively propagated by cloves or bulbils. Some cultivars are reported to produce flowers but with no seed set there is no seed setting. Garlic cultivars differ in maturity, bulb size, clove size and number, scale colour, bolting and flowering habits (cite source). However, it encounters a number of diseases which that attack the crop during its growth and development resulting in low yield thus causing economic loss. Among several diseases that affect garlic, purple blotch caused by *Alternaria porri* (Ellis) Ciferri has been emerging as a serious disease affecting both foliage and bulb development. Since, it is a foliar pathogen it alters the metabolic processes by decreasing the photosynthetic activity. A foliar infection of upto 90 per cent-% has been reported in susceptible cultivars of garlic are grown (Source). Purple blotch initially starts as

Formatted: Highlight

numerous tiny, white, circular or irregular spots. These spots gradually increase in size, become oval-shaped or irregular and white coloured spots eventually changing to violet colour. Later stages of development show the central portion of the spots changing to purple, surrounded by a pale yellow-orange to salmon band beyond which is a pale green zone. The dark purple colour is the most distinctive symptom of the disease. A distinct yellowing usually extends from both ends of the spots, often reaching the tips and bases of the leaves [10]. Complete damage to the leaf tissues is observed at the time of bulb maturity followed by blackening of scales resulting in reduced shelf life of the bulb [2]. Significant-A significant reduction in bulb yield (25-60 %) due to drying of leaves has been observed reported in garlic[3].

Formatted: Highlight

Overall the purple blotch in garlic has been reported as economically important disease in Gujarat [4], Maharashtra [5], Andhra Pradesh [6], Haryana, Karnataka [7], Odisha [8] and Himachal Pradesh [9]. Purple blotch initially starts as numerous tiny, white, circular or irregular spots. These spots gradually increase in size, become oval-shaped or irregular and white coloured spot eventually changing to violet colour. Later stages of development show the central portion of the spots changing to purple, surrounded by a pale yellow-orange to salmon band beyond which is a pale green zone. The dark purple colour is the most distinctive symptom of the disease. A distinct yellowing usually extends from both ends of the spots, often reaching the tips and bases of the leaves [10]. Host plant resistance is an effective, economic and environmentally safe component in an integrated approach. To overcome the breakdown of resistance, an alternate option is to cultivate varieties with durable resistance. Therefore, an attempt was made to identify resistant sources with a view to mitigate loss in farmers' field.

2. MATERIAL AND METHODS

A Totally of thirty seven 37 garlic genotypes collected from ICAR-DOGR (Directorate of Onion and Garlic), Rajgurunagar, Pune and ICAR-IARI, New Delhi along with local genotypes were subjected for to a field screening during Kharif 2019 against purple blotch at AINRPOG (All India Network Research Project on Onion and Garlic), MARS (Main Agricultural Research Station), Dharwad. All the genotypes were sown in two rows of two meter length. All the recommended agronomic practices were followed, except disease management practices. Inoculum of *Alternariaporri* (2×10^6 spores/ml) was uniformly sprayed during evening hours at 30 days after sowing to create uniform disease pressure throughout the field. The disease severity was recorded at 90 days after sowing using 0-5 scale a scale of 0-5 [7] by randomly selecting ten plants in each genotype. Based on their reaction disease severity, genotypes were categorized into immune (0 %), resistant (1-10 %), moderately resistant (11-20 %), moderately susceptible (21-40 %), susceptible (41-75 %) and highly susceptible (75-100 %).

Chart 1: Disease scoring scale (0-5 grade) for purple blotch of garlic

Disease score	Per cent leaf area infected	Disease reaction
0	No disease symptoms	Immune

1	A few spots towards tip covering 10 per-cent% leaf area	Resistant
2	Several dark purplish brown patches covering 11 to 20 per-cent% leaf area	Moderately resistant
3	Several patches with pale outer zone covering 21 to 40 per-cent% leaf area	Moderately susceptible
4	Yellow streaks covering 41 to 75 per-cent% leaf area	Susceptible
5	Complete drying of the leaves or breaking of leaves from centre	Highly susceptible

3. RESULTS AND DISCUSSION

~~Thirty-Thirty~~seven genotypes were grouped into different reaction types based on the results obtained from the study (Table 1). Among 37 genotypes, five genotypes (DOGR81W, DOGR150, DOGR517, DOGR534 and DOGR543) showed immune reaction whereas, 20 genotypes (DOGR51, DOGR75, DOGR102, DOGR113, DOGR119, DOGR181, DOGR325, DOGR353, DOGR389, DOGR426, DOGR548, DOGR569, DOGR604 and DOGR744) showed resistant reaction and 10 genotypes (Godavari, GG2, GG4, G41, DWG1, DOGR185, DOGR228, DOGR329, DOGR440 and DOGR756) showed moderately resistant reaction ~~while, the~~ remaining two genotypes (DOGR409 and HRG Local) were found moderately susceptible to purple blotch of garlic. However, none of the genotypes showed susceptible or highly susceptible reaction to purple blotch (Table 2). (Indicate which were your checks/commercial varieties in the study, which was the most resistant and what was the severity score)

Management of the disease through host plant resistance has been the best choice in all crop improvement programmes. Utilization of resistant cultivars in farming system is the most simple, effective and economical method in the management of plant diseases. Besides this, the resistant cultivars conserve the natural resource and reduce the cost, time and energy when compared to the other methods of disease management [11].

The results are in agreement with previous reports which demonstrated that, only a few lines are immune while the majority have resistance to purple blotch under natural infestation in open field condition [12]. Sughaet *et al.* [13] evaluated 94 onion genotypes under natural conditions and designated just two varieties, IC39178 and IC49371 as resistant to purple blotch. In the same context, Behera *et al.* [14] observed VG-18 cultivar as resistant and another 12 lines as moderately resistant to purple blotch. The present findings are in line with Agarwal and Tiwari [15], Nandini *et al.* [16] and Kowser *et al.* [17] who screened different garlic genotypes against purple blotch and they noted that ~~per-cent the percentage~~ disease index was negatively and significantly correlated with the bulb yield and keeping quality. ~~Although,~~ the field screening of different genotypes showed variable responses to purple blotch, ~~itis required to be~~ further ~~confirmed throughfield~~ evaluation~~ng~~ in multi-location hotspot areas for purple blotch ~~is required~~ in order to determine their efficiency as pre-breeding lines.

Table 1: Reaction of garlic genotypes to purple blotch caused by *Alternaria porriduring* Kharif 2019

Sl. No.	Genotypes	Disease severity grade	Sl. No.	Genotypes	Disease severity
---------	-----------	------------------------	---------	-----------	------------------

Formatted: Highlight

					grade
1	G41	2	20	DOGR181	1
2	G50	1	21	DOGR185	2
3	G282	1	22	DOGR228	2
4	GG2	2	23	DOGR325	1
5	GG4	2	24	DOGR329	2
6	Bhima Omkar	1	25	DOGR353	1
7	Bhima Purple	1	26	DOGR389	1
8	Godavari	2	27	DOGR409	3
9	Phule Baswant	1	28	DOGR426	1
10	Gadag Local	1	29	DOGR440	2
11	HRG Local	3	30	DOGR517	0
12	DWG1	2	31	DOGR534	0
13	DOGR51	1	32	DOGR543	0
14	DOGR75	1	33	DOGR548	1
15	DOGR81W	0	34	DOGR569	1
16	DOGR102	1	35	DOGR604	1
17	DOGR113	1	36	DOGR744	1
18	DOGR119	1	37	DOGR756	2
19	DOGR150	0			

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Table 2: Grouping of garlic genotypes based on reaction to purple blotch caused by *Alternaria porri*

Disease grade	Per cent per cent leaf infection	Genotypes	Disease reaction	No. of Genotypes
0	0	DOGR81W, DOGR150, DOGR517, DOGR534 and DOGR543 G50, G282, Bhima Omkar, Bhima Purple, Gadag Local, Phule Baswant, DOGR51, DOGR75, DOGR102,	Immune	5
1	1-10	DOGR113, DOGR119, DOGR181, DOGR325, DOGR353, DOGR389, DOGR426, DOGR548, DOGR569, DOGR604 and DOGR744	Resistant	20
2	10-20	G41, GG2, GG4, Godavari, DWG1, DOGR185, DOGR228, DOGR329, DOGR440 and DOGR756	Moderately Resistant	10
3	21-40	HRG Local and DOGR409	Moderately susceptible	2
4	40-75	-	Susceptible	Nil
5	More than 75	-	Highly susceptible	Nil

I think table two is enough to avoid repetition,

Provide some important agronomic data (duration to maturity, plant height, bulb size etc) as well for each genotype

It would improve your paper if you perform cluster analysis to group your genotype and and correlation to understand how disease severity affects plant growth.

4. CONCLUSION

The results of screening of garlic genotypes against purple blotch reveal that, out of 37 genotypes screened under artificial epiphytotic condition, five genotypes showed immune reaction and five genotypes showed resistant reaction whereas, ten genotypes showed moderately resistant and the remaining two genotypes showed moderately susceptible reaction while, none of the genotypes showed a susceptible and highly susceptible reaction. So, farmers can cultivate those immune and resistant genotypes in place of local cultivars in the management of purple blotch disease to get higher bulb yield. [\(Can you compare the resistant of the commercial varieties and the genotypes evaluated?\)](#)

REFERENCES [\(Make sure you follow the journals guideline and include DOI where possible\)](#)

1. Vijaykumar KN, Shripad Kulkarni, Patil PV, Kambrekar DN, Shashidhar TR, Eco-friendly management of purple blotch of garlic caused by *Alternaria porri* (Ellis) Ciferri. Biol Forum Int J. 2022;14(2):1-7.
2. Vijaykumar KN, Shripad Kulkarni, Patil PV, Kambrekar DN, Shashidhar TR, *In vitro* evaluation of new molecules of fungicides against purple blotch *Alternaria porri* (Ellis) Ciferri of garlic (*Allium sativum* L.). Pharma Innov J. 2021;10(12):1048-1054.
3. Bisht IS, Agarwal RC, Susceptibility to purple blotch (*Alternaria porri*) in garlic. Ann App Biol. 1993;122:31-38.
4. Raju KS, Studies on purple blotch of onion (*Allium cepa* L.) caused by *Alternaria porri*. M. Sc. (Agri.) Thesis, Saurashtra Univ. Junagadh, Gujarat; 1970.
5. Bhangale GT, Joi MB, Role of thrips in development of purple blotch of onion. J Maharashtra Agric Univ. 1983;8(3):299-300.
6. Gupta RP, Srivastava PK, Pandey UB, Control of purple blotch disease of onion seed crop. Indian Phytopath. 1986;39(1):303-304.
7. Sharma SR, Effect of fungicidal sprays on purple blotch and bulb yield of onion. Indian Phytopath. 1986; 39:78-82.
8. Mishra D, Mahanta IC, Chhotaray PK, Chemical control of purple blotch of onion in Odisha. J Agric Res. 1989;2(1):25-28.
9. Sugha SK, Tyagi PD, Effectiveness of metalaxyl + mancozeb against purple blotch (*Alternaria porri*) of onion (*Allium cepa* L.) in Himachal Pradesh. Indian J Agric Sci. 1994;64(2):135-136.
10. Aveling TAS, Purple blotch (*Alternaria porri*) of onion. Recent Res Dev Plant Pathol. 1998;2:63-76.
11. Roopadevi, Patil PV, In vitro bioassay of fungicides, bioagents, botanicals and itk's against *Pyricularia grisea* (Cooke) Sacc. incitant of pearl millet blast. Int J Pure App Biosci. 2017;5(5):1457-1463.
12. Pathak DP, Singh A, Despande A, Sridar TT, Source of resistance to purple blotch in onion. Veg Sci. 1986;13:300-303.

13. Sugha SK, Develash RK, Tyagi PD, Performance of onion genotypes against purple blotch pathogen. South Indian Hort.1992;40:297.
14. Behera S, Santra P, Chattopadhyay S, Das S, Maity TK, Variation in onion varieties for reaction to natural infection of *Alternaria porri* (Ellis) Cif. and *Stemphylium vesicarium* (Wallr.). Bioscan.2013;8(3):759-761.
15. Agarwal A,Tiwari RS, Evaluation of garlic (*Allium sativum*) genotypes for yield and susceptibility to purple blotch. JHortic For.2013;5(4):48-52.
16. Nandini KS, Umamaheswarappa P, Srinivasa V, Abhishek KN, Sindhu K, Lavanya KS, Performance of garlic (*Allium sativum* L.) genotypes for yield and quality attributes under central dry zone of Karnataka. J PharmacognPhytochem.2018;3:329-332.
17. Kowser AR, Amarananjundeswara R, Doddabasappa H, Aravinda Kumar B, Pannure A, Reaction of garlic genotypes to thrips and purple blotch in South India. JEntomolZool Stud.2019;7(1):63-66.

UNDER PEER REVIEW