

Original Research Article

Studies on nutrient and weed dynamics on growth and yield attributes of direct seeded rice

ABSTRACT

Field experiment was carried out during *kharif* 2020 and 2021 at wetland farm of S. V. agricultural College, Tirupati, Andhra Pradesh. The experiment was laid out in a split plot design with three replications. The treatments consisted of three main plots *viz.*, 100 % RDF (N₁), 125 % RDF (N₂) and 150 % RDF (N₃) and five sub plots *viz.*, Pre emergence (PE) application of pyrazosulfuron-ethyl 25 g ha⁻¹ *fb* penoxsulam + cyhalofop-p-butyl 125 g ha⁻¹ at 20 DAS (W₁), Pre emergence (PE) application of pyrazosulfuron-ethyl 25 g ha⁻¹ *fb* florypyrauxifenbenzyl + cyhalofop-p-butyl 150 g ha⁻¹ at 20 DAS (W₂), Pre emergence (PE) application of pyrazosulfuron-ethyl 25 g ha⁻¹ *fb* bispyribac-sodium 20 g ha⁻¹ at 20 DAS (W₃), hand weeding twice at 20 and 40 DAS (W₄) and unweeded check (W₅). Nutrient levels and weed management practices significantly influenced the growth parameters *viz.*, plant height, number of tillers m⁻² and dry matter production and yield attributes *viz.*, number of panicles m⁻², total number of grains panicle⁻¹ and number of filled grains panicle⁻¹ of direct seeded rice during both the years of investigation as well as in pooled mean. Interaction effect due to nutrient levels and weed management practices on growth parameters and yield attributes was found to be significant during both the years of study and in the pooled mean.

Keywords: Direct seeded rice, growth parameters, yield attributes

1. INTRODUCTION

Rice has been traditionally cultivated by transplanting seedlings into puddled soils and submerging the area. Standing water in the field causes macropore clogging, soil aggregate breakup in puddled soils, and the creation of hard pans beneath the subsurface layers of soils, as well as effective weed control. But, the sustainability of transplanted rice system is under

jeopardy due to several production-related constraints, viz., water, labour and energy scarcity, during peak period of demand and growing environmental concerns.

Transplanting of rice can be delayed due to late start of monsoon rainfall and the discharge of water into canals, as well as drudgery of puddling and physical transplanting. Under traditional system of rice cultivation, it takes around 5000 litres of water in order to produce 1 kg of rice (IRRI, 2001). A Review Pyrolysis: Different Agricultural Residues and Their Bio Char Characteristics (Makavana et al., 2021). As a result, rice production techniques that are not only able to cope with existing limits but also inexpensive, economically feasible, and capable of fulfilling future food demand must be investigated. In this context, direct seeded rice (DSR) is being assessed as viable alternative resource conservation method to address the problems interrelated with traditional rice farming in the face of looming agricultural resource shortages (Pandey and Velasco, 2002).

Direct seeded rice is subjected to greater weed competition than transplanted rice because both weed and crop seeds emerge at the same time and compete with each other right from its germination time resulting huge loss in grain yield (Rao *et al.*, 2007). Diverse and intensified weed flora can be controlled by pre emergence herbicides in combination with post emergence herbicides. In addition to weed management problems in DSR, the nutrient management is also very difficult. Various researches under direct seeded rice have been conducted on nutrient and weed management practices in alone. The combined effect of nutrient levels and weed management practices may have different impacts than their individual effects considering better plant growth under effective weed management practices thus the nutritional requirement of the crop may change.

2. MATERIAL AND METHODS

The experiment was conducted during *kharif*, 2020 & 2021 at wetland farm of S. V. Agricultural College, Tirupati, Andhra Pradesh geographically situated at 13.5°N latitude and 79.5°E longitude and at an altitude of 182.9 m above the mean sea level in the Southern Agro-Climatic Zone of Andhra Pradesh. The experimental soil was sandy clay loam in texture, slightly alkaline in reaction, low in organic carbon and available nitrogen and medium in available phosphorus and available potassium. The treatments in main plot consisted of three nutrient

levels *viz.*, 100 % RDF (N₁), 125 % RDF (N₂) and 150 % RDF (N₃) under main plots and five weed management practices comprised to sub plots *viz.*, .. Pre emergence (PE) application of pyrazosulfuron-ethyl 25 g ha⁻¹ *fb* penoxsulam + cyhalofop-p-butyl 125 g ha⁻¹ at 20 DAS (W₁), Pre emergence (PE) application of pyrazosulfuron-ethyl 25 g ha⁻¹ *fb* florasulfuron + cyhalofop-p-butyl 150 g ha⁻¹ at 20 DAS (W₂), Pre emergence (PE) application of pyrazosulfuron-ethyl 25 g ha⁻¹ *fb* bispyribac-sodium 20 g ha⁻¹ at 20 DAS (W₃), hand weeding twice at 20 and 40 DAS (W₄) and unweeded check (W₅). Pre emergence herbicide was applied immediately after sowing of rice crop and post emergence herbicides were imposed at 20 DAS. The rice variety chosen for the experiment was NLR-34449 and adopted a spacing of 20 cm x 10 cm. Nutrients *viz.*, nitrogen, phosphorus and potassium in the form of urea, single super phosphate and muriate of potash, respectively were applied as per the treatments . Nitrogen was applied in three splits at 15 DAS, tillering and at panicle initiation stages. Entire quantity of phosphorus was applied at the time of sowing and potassium was applied in two splits, ½ at the time of sowing and the remaining ½ at panicle initiation stage. Observations on growth and yield attributes of direct seeded rice was recorded at the time of harvest and the data was statistically analyzed by following the analysis of variance for split plot design as suggested by the Panse and Sukhatme (1985). Statistical significance was tested with F test at 5 per cent level of probability and compared the treatmental means with critical difference.

3. RESULTS AND DISCUSSION

A. Effect of nutrient levels on growth and yield attributes of direct seeded rice

Growth parameters like plant height, number of tillers m⁻² and dry matter production and yield attributes *viz.*, number of panicles m⁻², total number of grains panicle⁻¹ and number of filled grains panicle⁻¹ at harvest followed similar trend. Significantly higher values of above parameters *viz.*, plant height, number of tillers m⁻², dry matter production number of panicles m⁻², total number of grains panicle⁻¹ and number of filled grains panicle⁻¹ were obtained with 150 % RDF (N₃) than rest of the nutrient levels during both the years of study and in the pooled mean. Adequate supply of nutrients at different growth stages of crop might have enhanced the greater uptake of nutrients by the crop that lead to better crop growth. Similar results were reported by Kumar *et al.* (2022). The next best nutrient level in obtaining higher values was 125 % RDF (N₂)

which was significantly superior to 100 % RDF (N_1) during both the years of study and in the pooled mean.

B. Effect of weed management practices on growth and yield attributes of direct seeded rice

Among the different weed management practices, higher values of plant height, number of tillers m^{-2} and dry matter production and yield attributes *viz.*, number of panicles m^{-2} , total number of grains panicle⁻¹ and number of filled grains panicle⁻¹ at harvest were obtained with hand weeding twice at 20 and 40 DAS (W_4) which was statistically comparable with PE application of pyrazosulfuron-ethyl 25 g ha^{-1} *fb* floryprauxifenbenzyl + cyhalofop-p-butyl 150 g ha^{-1} at 20 DAS (W_2) during both the years of study and in the pooled mean. This might be due to reduced weed infestation as evident from lower weed density and biomass that facilitated for better utilization of growth resources which enhanced the vegetative and reproductive potential of the crop. The results are in accordance with those of Yadav *et al.* (2018). Advantage of Agrivoltaics across the Food-Energy-Water Connection makavana *et al.* (2020). The next best weed management practice was PE application of pyrazosulfuron-ethyl 25 g ha^{-1} *fb* penoxsulam + cyhalofop-p-butyl 125 g ha^{-1} at 20 DAS (W_1) followed by PE application of pyrazosulfuron-ethyl 25 g ha^{-1} *fb* bispyribac-sodium 20 g ha^{-1} at 20 DAS (W_3) with significant disparity among them. The lowest values of plant height, number of tillers m^{-2} and dry matter production and yield attributes *viz.*, number of panicles m^{-2} , total number of grains panicle⁻¹ and number of filled grains panicle⁻¹ at harvest were registered with unweeded check (W_5) during both the years of study and in the pooled mean. This was obviously due to severe competition offered by the weeds for growth resources right from the seedling emergence leading to poor crop growth and inefficient translocation of photosynthates from source to sink which lead to diminished yield attributes of rice.

Among the various combinations of nutrient levels and weed management practices, significantly higher plant height, number of tillers m^{-2} and dry matter production and yield attributes *viz.*, number of panicles m^{-2} , total number of grains panicle⁻¹ and number of filled. Development of Solar Operated Walking Type Power Weeder kachhot *et al.* (2020).

Table 1. Growth parameters of direct seeded rice as influenced by nutrient levels and weed management practices at harvest

Treatments	Plant height (cm)			Number of tillers m ⁻² (No. m ⁻²)			Dry matter production (kg ha ⁻¹)		
	2020	2021	Pooled	2020	2021	Pooled	2020	2021	Pooled
Nutrient levels									
N ₁ : 100 % RDF	57.3	53.1	55.2	181	152	166	5468	4811	5139
N ₂ : 125 % RDF	68.4	63.3	65.8	220	190	205	6769	5937	6353
N ₃ : 150 % RDF	74.6	69.0	71.8	240	211	225	7454	6560	7007
SEm ±	1.45	1.34	1.40	4.5	4.9	4.2	159.2	146.5	161.3
CD (P=0.05)	5.7	5.3	5.5	18	19	16	625	575	633
Weed management practices									
W ₁ : PE application of pyrazosulfuron-ethyl 25 g ha ⁻¹ <i>fb</i> penoxsulam + cyhalofop-p-butyl 125 g ha ⁻¹ at 20 DAS	63.9	59.1	61.5	201	169	185	6395	5628	6011
W ₂ : PE application of pyrazosulfuron-ethyl 25 g ha ⁻¹ <i>fb</i> florypyrauxifenbenzyle + cyhalofop-p-butyl 150 g ha ⁻¹ at 20 DAS	83.0	76.8	79.9	283	253	268	8572	7510	8041
W ₃ : PE application of pyrazosulfuron-ethyl 25 g ha ⁻¹ <i>fb</i> bispyribac-sodium 20 g ha ⁻¹ at 20 DAS	57.2	53.0	55.1	165	133	149	5522	4860	5191
W ₄ : Hand weeding twice at 20 and 40 DAS	86.6	80.2	83.4	299	269	284	8917	7847	8382
W ₅ : Unweeded check	43.0	39.9	41.5	120	97	108	3411	3002	3207
SEm ±	1.41	1.31	1.36	6.5	6.0	6.9	147.8	130.2	131.8
CD (P=0.05)	4.1	3.8	4.0	20	18	21	431	380	385
Interaction									
N at W									
SEm ±	3.39	3.13	3.26	12.9	13.7	13.0	360.1	321.8	335.9
CD (P=0.05)	10.9	10.1	10.5	41	43	41	1169	1051	1110
W at N									
SEm ±	2.44	2.26	2.35	9.5	10.5	9.8	256.0	225.5	228.3
CD (P=0.05)	7.1	6.6	6.9	28	30	29	747	658	666

Table 2. Interaction effect of nutrient levels and weed management practices on plant height (cm) of direct seeded rice at harvest

Treatments	2020-21				2021-22				Pooled			
	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)
W ₁	52.3	66.7	72.6	63.9	48.4	61.7	67.2	59.1	50.4	64.2	69.9	61.5
W ₂	72.0	85.4	91.6	83.0	66.7	79.1	84.8	76.8	69.4	82.2	88.2	79.9
W ₃	53.2	56.5	62.0	57.2	49.2	52.3	57.5	53.0	51.2	54.4	59.7	55.1
W ₄	72.5	92.2	95.2	86.6	67.1	85.3	88.1	80.2	69.8	88.8	91.7	83.4
W ₅	36.6	41.1	51.4	43.0	33.9	38.1	47.6	39.9	35.3	39.6	49.5	41.5
Mean (N)	57.3	68.4	74.6		53.1	63.3	69.0		55.2	65.8	71.8	

	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)
N	1.45	5.7	1.34	5.3	1.40	5.5
W	1.41	4.1	1.31	3.8	1.36	4.0
N at W	3.39	10.9	3.13	10.1	3.26	10.5
W at N	2.44	7.1	2.26	6.6	2.35	6.9

Table 3. Interaction effect of nutrient levels and weed management practices on number of tillers (No. m⁻²) of direct seeded rice at harvest

Treatments	2020-21				2021-22				Pooled			
	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)
W ₁	146	211	245	201	114	179	213	169	130	195	229	185
W ₂	249	292	309	283	218	263	277	253	234	278	293	268
W ₃	141	163	191	165	109	131	159	133	125	147	175	149
W ₄	259	311	326	299	229	280	299	269	244	295	313	284
W ₅	110	121	127	120	90	96	104	97	100	109	115	108
Mean (N)	181	220	240		152	190	211		166	205	225	

	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)
N	4.5	18	4.9	19	4.2	16
W	6.5	20	6.0	18	6.9	21
N at W	12.9	41	13.7	43	13.0	41
W at N	9.5	28	10.5	30	9.8	29

Comment [A1]: DATA check it in proper formating

Table 4. Interaction effect of nutrient levels and weed management practices on dry matter production (kg ha⁻¹) of direct seeded rice at harvest

Treatments	2020-21				2021-22				Pooled			
	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)
W ₁	4890	6903	7392	6395	4303	6074	6505	5628	4597	6488	6949	6011
W ₂	7443	8803	9469	8572	6550	7647	8332	7510	6997	8225	8901	8041
W ₃	4748	5607	6213	5522	4178	4934	5467	4860	4463	5270	5840	5191
W ₄	7492	9500	9761	8917	6593	8360	8590	7847	7042	8930	9175	8382
W ₅	2765	3034	4435	3411	2433	2670	3903	3002	2599	2852	4169	3207
Mean (N)	5468	6769	7454		4811	5937	6560		5139	6353	7007	

	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)
N	159.2	625	146.5	575	161.3	633
W	147.8	431	130.2	380	131.8	385
N at W	360.1	1169	321.8	1051	335.9	1110
W at N	256.0	747	225.5	658	228.3	666

Table 5. Yield attributes of direct seeded rice as influenced by nutrient levels and weed management practices

Treatments	Number of panicles (No. m ⁻²)			Total number of grains panicle ⁻¹			Number of filled grains panicle ⁻¹		
	2020	2021	Pooled	2020	2021	Pooled	2020	2021	Pooled
Nutrient levels									
N ₁ : 100 % RDF	131	110	121	102	99	100	88	85	87
N ₂ : 125 % RDF	159	138	149	116	113	115	100	98	99
N ₃ : 150 % RDF	174	153	163	128	125	126	110	108	109
SEm ±	3.0	3.5	3.2	2.5	2.7	2.6	2.1	2.2	2.0
CD (P=0.05)	12	14	13	10	11	10	8	9	8
Weed management practices									
W ₁ : PE application of pyrazosulfuron-ethyl 25 g ha ⁻¹ /b penoxsulam + cyhalofop-p-butyl 125 g ha ⁻¹ at 20 DAS	146	123	134	111	107	109	95	93	94
W ₂ : PE application of pyrazosulfuron-ethyl 25 g ha ⁻¹ /b floryprauxifenbenzyle + cyhalofop-p-butyl 150 g ha ⁻¹ at 20 DAS	206	183	195	141	138	139	121	119	120
W ₃ : PE application of pyrazosulfuron-ethyl 25 g ha ⁻¹ /b bispyribac-sodium 20 g ha ⁻¹ at 20 DAS	120	97	108	99	96	98	86	83	84
W ₄ : Hand weeding twice at 20 and 40 DAS	217	195	206	148	144	146	127	125	126
W ₅ : Unweeded check	87	70	78	79	76	77	68	65	67
SEm ±	3.7	4.4	4.0	2.3	2.1	2.3	2.0	1.9	2.0
CD (P=0.05)	11	13	12	7	6	7	6	6	6
Interaction									
N at W									
SEm ±	8.3	9.9	9.1	5.6	5.4	5.4	4.8	4.8	4.7
CD (P=0.05)	26	31	28	18	18	18	16	16	15
W at N									
SEm ±	6.4	7.6	7.0	4.0	3.6	3.7	3.4	3.3	3.4
CD (P=0.05)	19	22	20	12	10	11	10	10	10

Table 6. Interaction effect of nutrient levels and weed management practices on number of panicles (No. m⁻²) of direct seeded rice

Treatments	2020-21				2021-22				Pooled			
	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)
W ₁	106	153	178	146	83	130	155	123	95	142	166	134
W ₂	181	212	224	206	158	191	201	183	170	202	213	195
W ₃	102	118	139	120	79	95	116	97	91	107	127	108
W ₄	188	226	237	217	166	203	217	195	177	215	227	206
W ₅	80	88	92	87	65	70	75	70	73	79	84	78
Mean (N)	131	159	174		110	138	153		121	149	163	

	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)
N	3.0	12	3.5	14	3.2	13
W	3.7	11	4.4	13	4.0	12
N at W	8.3	26	9.9	31	9.1	28
W at N	6.4	19	7.6	22	7.0	20

Table 7. Interaction effect of nutrient levels and weed management practices on total number of grains panicle⁻¹ of direct seeded rice

Treatments	2020-21				2021-22				Pooled			
	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)
W ₁	91	114	127	111	88	111	123	107	90	112	125	109
W ₂	127	141	154	141	124	139	151	138	125	140	153	139
W ₃	91	95	112	99	88	92	109	96	89	94	110	98
W ₄	128	156	160	148	125	153	156	144	126	154	158	146
W ₅	74	75	88	79	70	72	85	76	72	73	87	77
Mean (N)	102	116	128		99	113	125		100	115	126	

	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)
N	2.5	10	2.7	11	2.6	10
W	2.3	7	2.1	6	2.3	7
N at W	5.6	18	5.4	18	5.4	18
W at N	4.0	12	3.6	10	3.7	11

Table 8. Interaction effect of nutrient levels and weed management practices on number of filled grains panicle⁻¹ of direct seeded rice

Treatments	2020-21				2021-22				Pooled			
	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)	N ₁	N ₂	N ₃	Mean (W)
W ₁	79	98	109	95	76	96	106	93	77	97	108	94
W ₂	109	122	133	121	107	119	130	119	108	120	132	120
W ₃	78	82	96	86	76	80	94	83	77	81	95	84
W ₄	110	134	138	127	108	132	134	125	109	133	136	126
W ₅	64	65	76	68	60	62	73	65	62	63	75	67
Mean (N)	88	100	110		85	98	108		87	99	109	

	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)	SEm±	CD (P=0.05)
N	2.1	8	2.2	9	2.0	8
W	2.0	6	1.9	6	2.0	6
N at W	4.8	16	4.8	16	4.7	15
W at N	3.4	10	3.3	10	3.4	10

grains panicle⁻¹ at harvest were observed with 150 % RDF coupled with hand weeding twice at 20 and 40 DAS (N₃W₄) which was statistically comparable with 125 % RDF coupled with hand weeding twice at 20 and 40 DAS (N₂W₄) and 150 % RDF coupled with PE application of pyrazosulfuron-ethyl 25 g ha⁻¹ *fb* florpyrauxifenbenzyl + cyhalofop-p-butyl 150 g ha⁻¹ at 20 DAS (N₃W₂). The lowest value of plant height, number of tillers m⁻² and dry matter production and yield attributes *viz.*, number of panicles m⁻², total number of grains panicle⁻¹ and number of filled grains panicle⁻¹ at harvest were noticed with 100 % RDF in combination with unweeded check (N₁W₅) during both the years of study and in the pooled mean.

4. CONCLUSION

From the present investigation it was concluded that among nutrient levels, 150 % RDF was found to be significant superior to other nutrient levels. Among various weed management practices, hand weeding twice at 20 and 40 DAS (W₄) which was statistically comparable with PE application of pyrazosulfuron-ethyl 25 g ha⁻¹ *fb* florpyrauxifenbenzyl + cyhalofop-p-butyl 150 g ha⁻¹ at 20 DAS (W₂).

REFERENCES

- IRRI (International Rice Research Institute). 2001. Annual Report 2000-01. Rice research: the way forward. Los Banos. International Rice Research Institute, Philippines.
- Makavana J.M., Kalaiya, S. V. Chauhan P. M. and Dulawat M. S. (2020). "Advantage of Agrivoltaics Across the Food-Energy-Water Connection". *Acta Scientific Agriculture* 4.8: 15-17.
- Kumar R., Singh DK, Singh VK, Bhatnagar A, Pareek N, Nanda, G Rawat A. Influence of different establishment methods and nutrient management practices on growth attributes of rice (*Oryza sativa* L.). *Indian Journal of Agricultural Research*. 2022;10: 1-7.
- Kachhot, A. R., Dulawat, M. S., Makavana, J. M., Dobariya, U. D., & Vadher, A. L. (2020). Development of Solar Operated Walking Type Power Weeder. *International Journal of Environment and Climate Change*, 10(12), 211-223. <https://doi.org/10.9734/ijecc/2020/v10i1230298>
- Pandey S, Velasco L. Economics of direct seeding in Asia: patterns of adoption and research priorities. In: *Direct Seeding: Research Strategies and Opportunities*. (eds. Pandey S, Mortimer M, Wade L, Tuong TP, Lopes K, Hardy B) International Rice Research Institute, Los Banos Philippines. 2002.
- Panse, VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*. Indian Council of Agricultural Research, New Delhi. 1985;100-174.
- Rao AN, Johnson DE, Sivaprasad B, Ladha JK, Mortimer AM. Weed management in direct-seeded rice. *Advances in Agronomy*. 2007;93: 153–155.

Yadav V, Tiwari RK, Tiwari P, Tiwari J. Integrated weed management in aerobic rice (*Oryza sativa* L.). International Journal of Current Microbiology and Applied Sciences. 2018; 7(1): 3099-3104.

Rao AN, Johnson DE, Sivaprasad B, Ladha JK and Mortimer AM. 2007. Weed management in direct-seeded rice. Advances in Agronomy 93: 153–155.

Makavana J. M., Sarsavadia P. N., Chauhan P. M., Dulawat M. S., Dobariya U. D. and Yadav R. (2021). A Review Pyrolysis: Different Agricultural Residues and Their Bio Char Characteristics, International Journal of Environment and Climate Change, 11(7): 80-88.

UNDER PEER REVIEW