

FRONT LINE DEMONSTRATION OF PADDY DRUM SEEDER (8 ROW) UNDER WELL IRRIGATED RICE IN NALGONDA DISTRICT, TELANGANA

ABSTRACT

Front line demonstration was conducted on direct seeded technology with drum seeder among the farmers, assessment of drum seeder implement was done with the assistance of Krishi Vigyan Kendra, Kampasagar, Nalgonda under Professor Jayashankar Telangana State Agricultural University, Telangana for consequently three years during *Kharif*, from 2018-19 to 2020-21. The comparison was made between direct seeded technology with drum seeder and manual transplanting with an objective to reduce the cost of production of paddy and subsequently to increase the net returns of farmers. Demonstrations revealed that, there were more number of tillers (395) and panicles (370)/m² in direct seeded technology with drum seeder compared to farmers practice (365 and 346/ m²). Drum seeder technology influence plant height (68.2cm), number of tillers/m² (395) and yield attributing parameters viz., No. panicles m⁻² (370), Panicle length (16.9 cm) and No. of grains / panicle (182). Direct sowing of paddy has recorded higher grain yield (6,512 kg ha⁻¹) which was 6.4 per cent over farmers practice (6,094 kg ha⁻¹) with net saving of Rs. 8,990/- on cost of cultivation ha⁻¹. The Gross returns and net returns (Rs.1, 20,795 and Rs. 69, 265/ha) by drum seeder method was more compared to farmers practice (Rs.1, 13,036/ha and Rs. 52, 516/ha). Drum seeder method saved the cost of cultivation by 14.8 per cent whereas, gross returns and net returns were increased by 6.4 and 24.2 per cent respectively over farmers practice. Seed rate also reduced to 30.0 kg ha⁻¹ as against farmers practice (64.5 kg ha⁻¹). Direct seeding technology with drum seeder helps in reducing the cost on nursery raising and transplanting overall cost of cultivation reduction by 14.8 per cent. The crop attained to harvest before 7-10 days duration with early crop harvest.

Keywords: Rice, Frontline Demonstrations, Drum Seeder

Introduction

Rice (*Oryza sativa* L.) is the most important staple food crop in Nalgonda district of Telangana cultivating in 87642 ha area during *Kharif* and 63661 ha during Rabi with an average productivity of 3450 kg/ha. The major rice area is under NSP left canal irrigated red soils/ Red soils with wells, tanks, kuntas as irrigation source. Present management recommendations for rice in Telangana include planting 20 to 25 day-old seedlings @ 2 per hill at 20 x 15 cm spacing with hand weeding and continuous flooding (Vyavasaya Panchangam, 2018). Majority of farmers use over aged seedlings due to delayed monsoon. Rice is being cultivated by raising nursery for about 30 days. Later seedlings are pulled and transplanted manually in zigzag manner without adopting proper spacing in the main field after puddling. Cost of cultivation is rising year by year due to many reasons i.e social reasons, situational factors and input cost. At present cost of cultivation per hectare is between Rs.51300/- to Rs 53000/-. This is mainly due escalation of labour wages and scarcity of labour in villages during agricultural season, labour requirement is very intense at the time of transplanting and increase in price of fertilizers. To overcome this, direct seeding of Rice with drum seeder can reduce the labour requirement during transplanting, technology is very simple and can be adopted by the farmers easily, cost of cultivation can be reduced and receive quality crop harvest at 7- 10 days earlier than normal transplanted field. Direct seeding of Rice with drum seeder holds special significance in the present day production system with regard to saving labour component by 30-50% and increases productivity by 20-30%. Transplanting is not profitable due to high labour wages and problem of non-availability of labour during peak period of operation (Singh et al., 2005). Due to non-availability of labour, hike in input cost and water shortage has lead to uneconomic rice cultivation. In the last five years the cost of production on different operations is increased by

33 per cent on seed, 45 per cent on chemical fertilizers, 100 per cent on labour cost, 35-40 per cent on tillage operations. Transplanting alone costs about 15 per cent of total rice production cost and delayed transplanting due to labour shortage causes sustainable loss in yield (Ponnuswamy et al., 1999). Because of uncertainty in rainfall and increase in cost of production the cultivation of rice has become unprofitable in Nalgonda and same is the case in other growing regions of Telangana (Visalakshi and Sireesha, 2013). The transplanting of rice seedlings which is a highly labour-intensive and expensive operation can be replaced by direct seeding that can reduce labour needs by more than 20 per cent in terms of working hours required (Santhi et al., 1998). The cost of operation of drum seeder is Rs. 40.50 per hour and Rs. 320 per hectare (Chavan and Palkar, 2010). So, direct seeding is much helpful as it requires less labour and time by skipping the nursery raising and transplanting to the field manually. Considering the above points, Front line demonstrations were conducted to popularize the drum seeder among the farmers, feasibility of paddy sowing by drum seeder was done under supervision of Krishi Vigyan Kendra (Extension unit of Professor Jayashankar Telangana State Agricultural University, Telangana.), Nalgonda for three years during *Kharif*, 2018-19 to 2020-21. The comparison was made between direct sowing of paddy using drum seeder and farmers practice with an objective to reduce the cost of production of paddy and subsequently improve the returns from unit in farmers' fields.

RESEARCH PROCEDURE

The Krishi Vigyan Kendra, Kammasagar, Telangana has demonstrated 8 row paddy drum seeder supplied by Farm Machinery Research Centre, Rajendranagar during *Kharif*, 2018. Specifications of drum seeder are given in Table A. A total of 30 demonstrations were conducted (Ten in 2018-19, ten in 2019-20 and ten in 2020-2021) during *Kharif* season in farmers' fields under puddle conditions. Demonstration details are given in Table B. Test variety was Ankur Puja of 130 days duration in all the locations. The demonstrations

comprised of two treatments viz., T₁-Direct sowing with drum seeder and T₂-Farmers practice (Traditional method of transplanting). Plot size for each treatment of Front line demonstration was 4000 m².

Direct sowing of paddy with drum seeder: Direct sowing with drum seeder, the paddy seeds were soaked in water for 24 hours followed by incubation in gunny bags for 24-48 hours. The field was well puddled and leveled after draining the standing water before sowing to avoid damage of sprouted seed and to enable water to spread uniformly over the field. After puddling, the field was left for 1-2 days for settling of the puddle soil. Care was taken to maintain thin film of water in the field at the time of sowing. Seed rate required for direct sowing of sprouted seed in puddled fields was quantified and pre germinated seeds were filled in all drums up to 2/3rd of its capacity at a time. The ground wheels make the impression to serve as a marker for next row and helps for good movement of drum seeder due to lugs provided on the periphery of the wheels. After turning the drum seeder for second row, care should taken that the first wheel should go through the same line of previous row in order to maintain the inter row distance of 20 cm, care should be taken to watch for any blocks of the drum. Refill the drums with seed when it reaches to the 1/4th of its capacity and continue the operation. Minimum two labour were required for completing the sowing operation i.e. one labour is for pulling the seeder and the other is for checking the drop of seeds from holes and filling the pre-germinated seeds in the drum. The field was kept moist without standing water in the field up to 20 days after sowing. Uniform dose of FYM @ 5.0 t ha⁻¹ and 120-60- 50 kg ha⁻¹ NPK were applied through urea, SSP, MOP. Entire P and K and 1/3 N was applied as basal, remaining N was applied in two equal splits at active tillering and panicle initiation stage. Weed growth was controlled by using pre emergence herbicide Pyrazosulfuron Ethyl 10% WP@ 80 gm/acre was applied at 1-3 days after sowing and post emergence application of bis pyribac sodium @ 80 ml/acre or herbicide mixture Cyhalofop-

Butyl 5.1% + Penoxsulam 1.02% @ 800- 1000 ml per acre was applied at 15-20 days after sowing.

Table A: Specifications of drum seeder

Power source	Hand operated
Row to row spacing	20 cm
Shape of the seed drum	Hyperboloid
Number of rows	8 rows
Diameter of the drum	20 cm
Diameter of the seed metering hole	9 mm
Number of seed metering hole	9 No
Weight of the unit	10 Kg
Type of ground wheel	
Diameter of the ground wheel	Lugged wheel
Operating speed	600 mm
Level of filling the seed drum	1 kmph / Walking speed
Weight of seed drum	Half volume
Weight of seed drum	600 g
Seed requirements	12 kg per acre

Table B: Details of Front Line demonstrations

Sr. No.	Year	No. of villages	No. of locations	Area (ha)
1	2018-19	5	10	4
2	2019-20	5	10	4
3	2020-21	5	10	4

Traditional method of nursery raising and transplanting:

In farmer's practice of traditional rice cultivation seedlings of 25-30 days old were pulled out manually from the nursery and transplanted in the main field at random @ 30-35 hills m⁻² using 4-6 seedlings per hill, maintaining 2 cm depth of water up to panicle initiation and 5-7 cm depth of water thereafter up to one week before harvest. The field was drained before application of fertilizers and one week before harvest. Manual weeding was done once at tillering to control weeds. Uniform dose of FYM @ 5.0 t ha⁻¹ and 120-60- 50 kg ha⁻¹ NPK were applied through urea, SSP, MOP. Entire P and K and 1/3 N was applied as basal, remaining N was applied in two equal splits at active tillering and panicle initiation stage. Both the treatments were received uniform plant protection and cultural management practices throughout the period of crop growth. Growth and yield attributes on 10 randomly selected hills were noted in each treatment plot. At harvest, grain yields from the net plots (5 m x 5 m) were recorded. Labour charges, cost of inputs were worked out to compute the cost of cultivation. Gross returns were calculated based on local market prices of paddy and straw. Benefit cost ratio was computed by dividing gross returns with cost of cultivation.

RESEARCH RESULT AND DISCUSSION

The findings of the present study as well as relevant discussion have been presented under following heads:

Yield parameters: Higher plant height of 68.2 cm was recorded with direct sowing by drum seeder compared to farmers practice (65.3cm) (Table 2). There were more number of tillers (395) and panicles (370) per m² in direct sowing by drum seeder compared to 365 tillers and

346 panicles per m² in farmer's practice which can be attributed to sowing of sprouted seeds at wider row spacing (20cm). Early establishment of seedlings in direct sown sprouted seed might be the reason for higher tillers per m² than conventional transplanting. These results are in accordance with Visalakshi and Sireesha (2013). Higher number of grains panicle⁻¹(182) and more panicle length (16.9 cm) were recorded in direct sowing by drum seeder compared to farmers practice (14.5 cm panicle length and 156 grains per panicle). Tiller to panicle conversion ratio was higher in direct sowing by drum seeder which might be due to favourable growth and better translocation of assimilates to the sink as it was revealed by more number of grains panicle⁻¹. Similar findings were reported by Halder and Patra (2007). Where as in farmers practice planting more seedlings per hill probably led to poor tillering which was also reflected in yield attributes such as number of panicles per metre square, panicle length and number of grain panicle⁻¹. Similar findings were observed by Shekhar et al. (2009). The direct seeded rice matured in 10 days earlier than transplanted rice. Wang and Sun (1990) noticed that duration can be shortened by 7-15 days in direct seeded rice compared to transplanted rice.

Grain yield: Grain yield was influenced by sowing methods. Direct sowing of paddy with drum seeder has recorded higher grain yield (6512 kg ha⁻¹) (Table 1) which was 6.4 per cent higher over farmers practice (6094 kg ha⁻¹). Higher number of grains panicle⁻¹ and panicle length (cm) might be the reason behind the yield increase in direct sowing method. Shekhar and Singh (1991) stated that direct sowing of sprouted seeds under puddled condition results in significant improvement in yield attributes like number of effective tillers and grain yield. Seeding of sprouted seeds with 8-row seeder after puddling increased grain yield (21.5%) over farmer's practice of transplanting (Halder and Patra, 2007). Lower paddy yields under farmers' practice might be due to planting 4 to 6 older seedlings (25 - 30 days old) with irregular spacing may lead to poor growth, poor tillering, lesser number of panicles per metre

square and yield. Earlier reports indicated that grain yield reduction is due to planting of older seedlings (Menete et al., 2008) at a higher density (San-oh et al., 2004).

Economics: A saving of Rs. 8990/- on cost of cultivation per hectare was realized indirect sowing of paddy with drum seeder besides increasing paddy yield. Gross returns (Rs. 1, 20,795/ ha) and net returns (Rs. 69, 265/ ha) (Table 1) was realised with drum seeder method compared to farmers practice (Rs. 1, 13,036 / ha gross returns and Rs. 52, 516/ ha net returns). This might be due to reduction in cost of cultivation and higher grain yield with drum seeder method. Higher Net returns of Rs. 69, 265/ha was obtained with drum seeder method due to lesser cost of cultivation and higher grain yield compared to farmers practice. The cost of cultivation was comparatively lesser in drum seeder method which resulted in an additional net profit of Rs.16749/ha as compared to conventional method of rice cultivation. Similar findings were reported by Halder and Patra (2007). Simultaneously benefit cost ratio was higher with drum seeder method (2.3) compared to farmers practice (1.8) because of lower cost of cultivation and increased yield with drum seeder method. Higher net returns and B:C ratio were recorded with drum seeder method due to no nursery raising and transplanting, less seed cost, reduced labour for weeding denoting lower cost of cultivation in drum seeder method resulted in increased profitability compared to farmers' practice. In drum seeder method cost of cultivation was reduced by 14.8 per cent whereas, gross returns and net returns were improved by 6.4 and 24.2 per cent, respectively over farmers practice.

Table 1: Yield Improvement in rice by direct sowing with rice drum seeder

Treatment	Average Yield kg/ha	Gross returns (Rs. ha⁻¹)	Cost of cultivation (Rs. ha⁻¹)	Net Returns (Rs)	C:B ratio	% Increase in yield
Direct sowing with drum seeder	6,512	1, 20,795	51530	69, 265	2.3	6.41

Farmers practice (Conventional transplanting method)	6,094	1, 13,036	60520	52, 516	1.86	
--	-------	-----------	-------	---------	-------------	--

Table 2: Growth parameters of paddy as influenced by direct sowing with drum seeder (Mean of 3 years)

Particulars	Direct sowing with drum seeder	Traditional method of transplanting (Farmers practice)
Seed rate (kg/ha)	30	68.7
Days to transplant	0	25-30
Cost of raising nursery (Rs./ha)	0	3950
Labour required for transplanting (for one hectare)	2	15-20
Spacing (cm)	20*5-8	Zig zag method
No. hills m ⁻²	26	30-35
Plant height (cm)	68.2	65.3
Number of tillers m ⁻²	395	365
Number panicles m ⁻²	370	346
Panicle length (cm)	16.9	14.5
Number of grains / panicle	182	156
Crop duration (days)	125	135

Extension Activities

Extension activities like Front line demonstrations, training programmes, exposure visits, field days, group discussions, Popular articles, kisan melas, radio and TV and newspaper coverage were conducted by KVK to popularize the drum seeder. A total no of 12 on and off campus training programmes were conducted for 360 beneficiaries.

Farmer Feedback

Farmers accepted the technology as it is viable and easy to practice. Good relief from labour shortage during the peak periods. As the crop duration is reduced by 7-10 days it facilitates in raising the summer pulse on conserved soil moisture without any moisture stress. The weed problem in the initial stage of the crop growth can be managed by application of selective herbicide (Jitendranath et.al 2012).

Conclusion: These demonstrations clearly established an advantage of direct sowing of paddy using drum seeder in puddled condition over conventional method of transplanting.

Seed rate can be reduced to 30 kg ha⁻¹ as against 68.7 kg ha⁻¹ in farmers practice. Direct sowing with drum seeder helps in reducing the cost on nursery raising and transplanting besides increasing yield by 6.4 per cent, reduces the crop duration and cost of cultivation. The cost of cultivation was reduced by 14.8 per cent and net returns increased by 24.2 per cent.

LITERATURE CITED

Chavan, S.P. and Palkar, S.M. (2010). Performance evaluation of paddy drum seeder. *Internat. J. Agric. Engg.*, 3 (2): 279- 287.

Halder, J. and Patra, A.K. (2007). Performance of eight-row drum seeder in direct seeded rice *Oryza Sativa* under puddled condition. *Indian J. Agric. Sci.*, 77 (12): 819823.

Jitendranth. S, C Chandrasekhararao and T G K Murthy (2012). Vari Sagu paddatulu (Telugu).

Menete, M.Z.L. Van, E.S.H.M. Brito, R.M.L., De Gloria, S.D. and Famba, S. (2008). Evaluation of system of rice intensification (SRI) component practices and their synergies on salt-affected soils. *Field Crops Res.*, 109: 34–44.

Ponnuswamy, K., Santhi, P., Kempu Chetty, N. and Subramaniam, M. (1999). Effect of various methods of establishment on growth and yield of rice. *Oryza*, 36: 294-305.

San-oh, Y., Mano, Y., Ookawa, T. and Hirasawa, T. (2004). Comparison of dry matter production and associated characteristics between direct-sown and transplanted rice plants in a submerged paddy field and relationships to planting patterns. *Field Crop Res.*, 87: 43–58.

Santhi, P., Ponnuswamy, K. and Chetty, N.K. (1998). Effect of seeding methods and efficient nitrogen management practices on the growth of lowland rice. *J. Ecobiol.*, 10: 123-132.

Shekhar, J. and Singh, C. M. (1991). Influence of methods and dates of stand establishment on growth and yield of rice. *Oryza*, 28: 45-48.

Shekhar, J., Mankotia, B.S. and Dev, S.P. (2009). Productivity and economics of rice (*Oryza sativa*) in systems of rice intensification in north western Himalayas. *Indian J. Agron.*, 54: 423-427.

Singh, G., Singh, R.G., Singh, O.P., Kumar, T., Mehta, R.K., Kumar, V. and Singh, P.P. (2005). Effect of weed management practices on direct seeded rice *Oryza sativa* under puddle low lands. *Indian J. Agron.*, 50:35-37.

Visalakshi, M. and Sireesha, A. (2013). Performance of drum seeder in direct sown paddy under puddled condition. *J. Res. ANGRAU*,41: 16-20.

Vyavasaya, Panchangam (2018). Professor Jayashankar Telangana State Agricultural University, Telangana. in the form of 142 PDF files.

Wang, H.Y. and Sun, T.S. (1990). The characteristics of machine direct-sown rice following wheat and the corresponding techniques. *Acta Agri. Uni. Jiangxiensis*, 12: 34-39.

UNDER PEER REVIEW