

Effect of Phosphorus and Zinc on Growth and Yield of Baby corn (Zeamays L.)

1. Abstract:-

A field study was carried out at the Crop Research Farm (CRF), Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj (UP) during the year 2021 zaid season. The experiment comprised of 9 treatments with different combinations of phosphorus and zinc levels replicated thrice in a Randomized Block Design. The primary goal of the experiment was to evaluate the Effect of phosphorus and zinc levels on growth and yield of Baby corn (*Zea Mays L.*). The phosphorus levels include (50, 60 and 70 kg/ha) and levels of zinc include (05, 15, 25 kg/ha). From the present investigation the profitable production of baby corn can be secure with application of phosphorus (70 kg/ha) and zinc (25 kg/ha).

Keyword : Phosphorus, Zinc, Growth and yield , Baby corn

1. Introduction

In addition to its diversified uses, maize is highly valued as animal feed, human food, and raw material in the industry. Next to rice and wheat, maize is the third most important cereal crop and has the highest production potential. Growing 'baby corn' for vegetable purposes is one recent development to diversify and add value to the food processing industry. A young, fresh, and finger-like green ear is harvested when the silk length is 2-3 cm, but before fertilization. (Pandey et al., 2000).

“Phosphorus is the second most important mineral nutrient after nitrogen. In terms of quantitative requirements for plants. It is necessary for growth, sugar and starch utilization, Photosynthesis, nucleus formation, cell division, fat and albumin production. Energy derived as a result of photosynthesis and carbohydrate metabolism, phosphate compounds store energy use in growth and reproduction” (Arya et al. 2001). In plants, it is readily translocated from older to younger tissues as roots, stems, and leaves are formed. An adequate amount of P promotes rapid growth and earlier maturity, as well as improving the quality of vegetative

growth. Deficiency of phosphorus results in crooked rows and small ear numbers in maize as the kernel twists.

In addition to lipids, proteins, and cofactors of auxins, it also plays an important role in the metabolism of nucleic acids **Mengel et al., (2001)**. The application of zinc has been proven to improve crop yields and quality **Hassan et al.,(2019)**. **Mousavi et al., (2007)** show that its deficiency reduces yields and deteriorates crop quality. In addition, higher Zn concentrations suppress cell differentiation and elongation, affecting biomass production **Tsonev et al., (2012)**.

2. Materials and methods

The research was conducted at the Crop Research Farm, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj, during the Zaid season 2021-22. It is located approximately 9 kilometers from Prayagraj city, near the Yamuna River, on the left side of the Prayagraj-Rewa Road, at 25.4089833, 81.8530037. The subtropical region of Uttar Pradesh, where Prayagraj is located, has scorching summers and mild winters. The annual rainfall here seems to be 1050 mm on average. “The data of the soil chemistry examination showed that the soil had a sandy loam texture, a pH of 7.2, small amounts of organic carbon (0.48 percent) and potassium (215.4 kg/ha), and small amounts of phosphorus that was easily accessible (13.6 kg/ha). The soil had a conductivity of 0.26 dS/m and was electrically conductive. Three replications have been used for every one of the nine treatment combinations. Tables 1 and 2 show the treatment details and treatment combinations, respectively. Phosphorus and zinc levels were maintained in accordance with the treatment combinations, and an analysis of each treatment was completed to determine the best treatment combination for Babycorn cultivation” [Krishna et al.2022]. The statistics were calculated and analysed using **Sunnam Hemanth Kumar** statistical approach, which is a one-way annova table used to compare more than two groups based on one factor with a F probability of 0.005% which was developed by Ronald Fisher in 1918..

3. Results and discussion

Data presented in table 1 and table 2 indicates significant increases in all growth parameters in chickpea at harvest, including plant height and dry weight per plant. Application of phosphorus 70 kg/ha and zinc 25kg/ha significantly increased yields, and remained at par

with with 60 kg/ha Phosphorus + 25 kg/ha Zinc and 70 kg/ha Phosphorus + 15 kg/ha Zinc results showed that significantly maximum plant dry weight 105.37 g/plant, highest plant height 158.17 cm, maximum Cobs/Plant (1.46), Length of Cob/plant (17.30 cm), Girth of Cob/plant (7.28 cm), Cob weight with husk (63.01 g), cob weight without husk (20.65 g), cob yield with husk (13.30 t/ha), Cob yield without husk (5.16 t/ha) at harvest. “Phosphorus P application increased plant height, dry weight, and number of cobs per plant, which could be attributed to improved early vegetative growth in terms of increased leaf area, dry matter accumulation, and a vigorous root system, which increased the number of cobs” [Masood et al. (2011)] “The source and sink relationship is responsible for the increase in cob yield caused by phosphorus application. It appears that increased photosynthate translocation from source to sink may have increased cob yield. Phosphorus increases yield due to its well-developed root system, increased N fixation and availability to plants, and favourable rhizosphere environments”[Hirpara et al. (2017)] “Zinc plays an important role in plant metabolism by influencing the activity of growth enzymes. It is also involved in carbohydrate metabolism, maintaining the integrity of protein synthesis, cellular membranes and regulation of auxin synthesis and pollen formation, which causes a greater number and length of cobs”[Anjum et al (2017)]. “The production of photosynthetic pigments and their transport to sinks is dependent on the presence of mineral nutrients, which has increased zinc uptake. Most photosynthetic pathways rely on enzymes and co-enzymes that are synthesised by mineral nutrients, and zinc application resulted in higher chlorophyll contents, synthesis of metabolites and growth-regulating substances, oxidation and metabolic activities, and ultimately better crop growth and development, which resulted in an increase in yield attributes of baby corn”[Arab et al. (2018) and Naik et al. (2020)]

3.1.Benefit cost ratio:-

In Table 2 The data presented in table 2 shows that benefit cost ratio (2.23) as influenced by different phosphorus and zinc levels in baby corn was found to be highest in treatment-9 with application of 70 kg/ha Phosphorus and 25 kg/ha Zinc and the minimum benefit cost ratio (1.97) was found to be in treatment-1 (application of 50 kg/ha Phosphorus and 5 kg/ha Zinc] as compared to other treatments

Table 1. Effect of Phosphorus and Zinc levels on growth attributes in Baby corn

| S.No | Treatments | Plant height | Plant Dry weight |
|------|-------------------------------------|--------------|------------------|
| 1 | 50 kg/ha Phosphorus + 5 kg/ha Zinc | 153.70 | 102.72 |
| 2 | 50 kg/ha Phosphorus + 5 kg/ha Zinc | 154.24 | 103.04 |
| 3 | 50 kg/ha Phosphorus + 25 kg/ha Zinc | 156.25 | 104.10 |
| 4 | 60 kg/ha Phosphorus + 5 kg/ha Zinc | 155.19 | 103.42 |
| 5 | 60 kg/ha Phosphorus + 15 kg/ha Zinc | 156.80 | 104.26 |
| 6 | 60 kg/ha Phosphorus + 25 kg/ha Zinc | 157.73 | 105.14 |
| 7 | 70 kg/ha Phosphorus + 5 kg/ha Zinc | 155.84 | 103.71 |
| 8 | 70 kg/ha Phosphorus + 15 kg/ha Zinc | 157.54 | 104.65 |
| 9 | 70 kg/ha Phosphorus + 25 kg/ha Zinc | 158.17 | 105.37 |
| | F test | S | S |
| | SEm (\pm) | 0.26 | 0.25 |
| | CD (P = 0.05) | 0.77 | 0.74 |

Table 2. Effect of Phosphorus and Zinc levels in yield attributes in Baby corn.

| S.No | Treatments | No. of Cobs/plant | Cob weight (g) | | Cob yield (t/ha) | | B:C ratio |
|---------------|-------------------------------------|-------------------|----------------|--------------|------------------|--------------|-----------|
| | | | With Husk | Without Husk | With Husk | Without Husk | |
| 1. | 50 kg/ha Phosphorus + 5 kg/ha Zinc | 1.12 | 60.93 | 17.63 | 11.75 | 3.73 | 1.27 |
| 2. | 50 kg/ha Phosphorus + 15 kg/ha Zinc | 1.16 | 60.75 | 18.42 | 12.03 | 3.93 | 1.32 |
| 3. | 50 kg/ha Phosphorus + 25 kg/ha Zinc | 1.29 | 61.78 | 19.47 | 12.72 | 4.48 | 1.70 |
| 4. | 60 kg/ha Phosphorus + 5 kg/ha Zinc | 1.21 | 61.06 | 18.79 | 12.28 | 4.14 | 1.46 |
| 5. | 60 kg/ha Phosphorus + 15 kg/ha Zinc | 1.33 | 62.20 | 19.78 | 12.88 | 4.60 | 1.80 |
| 6. | 60 kg/ha Phosphorus + 25 kg/ha Zinc | 1.41 | 62.84 | 20.39 | 13.16 | 5.01 | 1.97 |
| 7. | 70 kg/ha Phosphorus + 5 kg/ha Zinc | 1.25 | 61.42 | 19.17 | 12.48 | 4.30 | 1.63 |
| 8. | 70 kg/ha Phosphorus + 15 kg/ha Zinc | 1.37 | 62.49 | 20.14 | 13.09 | 4.83 | 1.88 |
| 9. | 70 kg/ha Phosphorus + 25 kg/ha Zinc | 1.46 | 63.01 | 20.65 | 13.30 | 5.16 | 2.08 |
| F test | | S | S | S | S | S | |
| SEm (±) | | 0.03 | 0.21 | 0.18 | 0.07 | 0.12 | |
| CD (P = 0.05) | | 0.09 | 0.63 | 0.55 | 0.21 | 0.36 | |

4. Conclusion:-

Based on the findings of this study, it is concluded that the profitable production of Baby corn can be ensured by applying phosphorus 70 kg/ha and zinc 25 kg/ha at the agroclimatic zone in Prayagraj, Uttarpradesh state, INDIA. These practises could be passed on to farmers in the Prayagraj, Uttarpradesh agroclimatic zone to boost their yields. Since the findings have been based on a single season of research.

ACKNOWLEDGEMENT

I express thankfulness to my advisor Dr. Rajesh Singh and all the faculty members of Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj -211007, Uttar Pradesh. For providing us essential facilities to undertake the studies.

References:-

- Anjum, S. A., Saleem, M. F., Shahid, M., Shakoore, A., Safeer, M., Khan, I., Farooq, A., Ali, I. and Nazir, U. 2017. Dynamics of soil and foliar applied boron and zinc to improve maize productivity and profitability. *Pakistan Journal of Agricultural Research*, 30(3): 294- 302.
- Arab, G.M., Dina A. Ghazi and El-Ghamry, A.M. 2018. Effect of zinc application on maize grown on alluvial soils. *J. Soil Sci. and Agric. Eng., Mansoura Univ.* **9**(9): 419-426.
- Arya, K.C. and Singh, S.N. 2001. Effect of different levels of phosphorus and zinc on yield and nutrients uptake of maize (*Zea mays* L.) with and without irrigation. *Indian Journal of Agronomy* 45(4) 717-721.
- Hassan, M.U.; Chattha, M.U.; Ullah, A.; Khan, I.; Qadeer, A.; Aamer, M.; Khan, A.U.; Nadeem, F.; Khan, T.A. Agronomic biofortification to improve productivity and grain Zn concentration of bread wheat. *Int. J. Agric. Biol.* 2019, 21, 615–620.
- Hirpara, D.V., Sakarvadia, H.L., Savaliya, C.M., Ranpariya, V.S. and Modhavadiya, V.L. (2017) Effect of different levels of boron and molybdenum on growth and yield of summer groundnut (*Arachis hypogaea* L.) under medium black calcareous soils of south Saurashtra region of Gujarat. *International Journal of Chemical Studies*, **5**(5), 1290-1293.

- M. M. 2020. Morpho-physiological impact of growth indices to Biofortification on growth and yield of sweet corn (*Zea mays* L. *Saccharata*). *Bulletin of Environment, Pharmacology and Life Sciences*, **9**(3): 37-43.
- Masood, T., R. Gul, F. Munsif, F. Jalal, Z. Hussain, N. Noreen, H. Khan, N. Din and H Khan. 2011. Effect of different phosphorus levels on the yield and yield components of maize. *Sarhad Journal of Agriculture*. Vol. 27, No.2: 167-170.
- Mengel, K.; Kosegarten, H.; Kirkby, E.A.; Appel, T. *Principles of Plant Nutrition*; Springer: Berlin, Germany, 2001.
- Mousavi, S.R.; Galavi, M.; Ahmadvand, G. Effect of zinc and manganese foliar application on yield, quality and enrichment on potato (*Solanum tuberosum* L.). *Asian J. Plant Sci.* 2007, **6**, 1256–1260.
- Naik, C., Meena, M. K., Ramesha, Y. M., Amaregouda, A., Ravi, M. V. and Dhanoji, Pandey, A.K., Prakash, V., Mani, V.P. And Singh, R.D., 2000, Effect of rate of nitrogen and time of application on yield and economics of Baby corn. *Indian J. Agron.*, **45**(2): 338-343.
- Sunnam Hemanth Kumar, Joy Dawson, Pole Shiva Kiran, Varsha Vyas V. Effect of Iron and Zinc Levels on Growth and Yield of Chickpea (*Cicer arietinum* L.). *International Journal of Current Microbiology and Applied Sciences*. 2020;**9**(11).
- Tsonev, T.; Lidon, F.J.C. Zinc in plants—An overview. *Emir. J. Food Agric.* 2012, **24**, 322–333.
- Krishna PM, Singh R, Indu T. Effect of phosphorus and zinc on yield and economics of baby corn (*Zea mays* L.). *The Pharma Innovation Journal* 2022; **11**(9): 1466-1469