

Original Research Article

EFFECT OF MICRONUTRIENTS ON YIELD OF ACID LIME (*Citrus aurantifolia* Swingle) cv. KAGZI LIME

ABSTRACT

The present investigation entitled “Effect of micronutrients on yield of [Acid-acid Lime lime](#) (*Citrus aurantifolia* Swingle) cv. Kagzi Lime” was carried out during *Ambe bahar* 2019 at ~~the~~ Horticultural Instructional Farm, C. P. College of Agriculture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar, District: Banaskantha, Gujarat. [The present Experiment-experiment consist fifteen treatments with three replications which](#) was laid out in a Randomized Block Design ~~with three replications. Total fifteen treatments were evaluated with respect to yield parameters of acid lime. The findings of the present studies indicated that the Among various treatments,~~ treatment ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.2 % (T₈) was significantly superior ~~over rest of treatments~~ with respect to yield ~~over rest of the other treatments. Further. Results-results~~ revealed that [the](#) maximum fruit diameter (4.58 cm), average fruit weight (44.06 g), fruit volume (43.83 cc), number of fruits per tree (946.00), fruit yield (41.68 kg per tree) and total fruit yield (115.46 quintal per hectare) were recorded with treatment ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.2 % (T₈) [as](#) compared to other treatments ~~and control~~.

Keywords: Micronutrients, yield, zinc sulfate, ferrous sulfate, borax and acid lime

INTRODUCTION

Acid Lime (*Citrus aurantifolia* Swingle), belongs to the family Rutaceae. Citrus is the third largest fruit crop grown in India, next to mango and banana. It is generally grown under both tropical and subtropical climatic conditions. It is a rich source of vitamin C and has good antioxidant properties. Fruits being acidic in nature, they are largely used for garnishing and flavoring several vegetarian and non-vegetarian dishes. Besides its value-added products like pickle, juice, squash *etc.*, lime peel oil, peel powder *etc* are also in great demand in soap and cosmetic industry (Debaje *et al.* 2011).

Zinc is required to obtain good fruit set and size. Its role in flowering is due to synthesis of tryptophan which is a precursor of auxin and promotes flowering. It also helps in the process of translocation of metabolites to the bud itself or to the site of bud development. Its deficiency produces small and narrow leaves, shorter shoot internodes and terminal dieback (Ryugo, 1988).

Iron is required for the synthesis of enzymes responsible for chlorophyll synthesis. ~~Iron deficiency is expressed as yellow leaves due to low levels of chlorophyll (chlorosis), which first appears on the younger upper leaves in interveinal tissues.~~ Severe iron deficiency may cause leaves to turn completely yellow or almost white and then brown as leaves die (Pandey and Sinha, 2006).

Comment [E1]: It is general statement, please delete and write the effect of micronutrients especially in acid lime

Boron is important element for flowering, fruiting, growth and quality of fruits. Boron also increase the chlorophyll content of leaves and play an important role in enzymatic activities. Foliar application of boron (B) improves tree growth, productivity and fruit quality in citrus. Deficiency of boron in citrus has serious consequences for tree health and crop production and also leads to low sugar content, granulation and excessive fruit abortion as well as rind thickness, symptoms that are seen regularly in fruit grown. So it has to be need of foliar application of boron for its role in the yield and fruit quality of citrus (Prasad *et al.* 2013).

Effective use of micronutrients in acid lime is one such research gap. Micronutrients can tremendously boost up acid lime flowering and fruiting quality. The problem of micronutrients deficiency in acid lime causes great concern to fruit growers and also flower drop as well as fruit drop major problem.

Comment [E2]: Which micronutrients please write

MATERIAL AND METHOD

Experiment was carried out during *Ambe bahar* of the year 2019 under field condition at the Horticultural Instructional Farm, C. P. College of Agriculture, Department of Horticulture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar, Dist. Banaskantha, Gujarat. The investigation was conducted on 15 years old plants of acid lime cultivar “Kagzi Lime”. All the plants selected were uniform in growth and size which planted at the distance of 6 m × 6 m and were subjected to uniform application of cultural practices like weeding, irrigation, manures, fertilizers and plant protection measures *etc.*

Experiment was laid out in a randomized block design with three replications. Total fifteen treatments were evaluated in the present study *viz.*, T₁: Control; T₂: ZnSO₄ 0.5 %; T₃: ZnSO₄ 1.0 %; T₄: FeSO₄ 0.5 %; T₅: FeSO₄ 1.0 %; T₆: Borax 0.2 %; T₇: Borax 0.4 %; T₈: ZnSO₄

0.5 % + FeSO₄ 0.5 % + Borax 0.2 %; T₉: ZnSO₄ 0.5 % + FeSO₄ 1.0 % + Borax 0.2 %; T₁₀: ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.4 %; T₁₁: ZnSO₄ 0.5 % + FeSO₄ 1.0 % + Borax 0.4 %; T₁₂: ZnSO₄ 1.0 % + FeSO₄ 0.5 % + Borax 0.2 %; T₁₃: ZnSO₄ 1.0 % + FeSO₄ 1.0 % + Borax 0.2 %; T₁₄: ZnSO₄ 1.0 % + FeSO₄ 0.5 % + Borax 0.4 %; T₁₅: ZnSO₄ 1.0 % + FeSO₄ 1.0 % + Borax 0.4 %.

RESULT AND DISCUSSION

Fruit diameter (cm)

The data related to effect of micronutrients on fruit diameter are presented in Table 1. The maximum fruit diameter (4.58 cm) was observed with treatment T₈ (ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.2 %) which was statistically at par with treatment T₉, T₁₀ and T₁₁. Whereas, the minimum fruit diameter (3.50 cm) was recorded with control plot the treatment (T₁-(control)).

Increased in The fruit diameter might be Increased due to the favorable effect was attributed to the fact that zinc is essential in the nitrogen metabolism (Asana *et al.*, 1971) and it also increase the synthesis of auxin which promote the cell size (Agarwala and Sharma, 1978). The cumulative effect of combination of Zn + Fe + B on faster cell division, cell expansion and increase volume of intercellular spaces in mesocarpic cells. It could also be due to higher mobilization of food and minerals from other parts of plants towards the developing fruit that are extremely active metabolic sink. Similar results were also found by Venu *et al.* (2016) in acid lime, Meena *et al.* (2016) in Nagpur mandarin, Trivedi *et al.* (2012) in guava, Razzaq *et al.* (2013) in mango, Yadav *et al.* (2014) in pomegranate, Meena *et al.* (2014) and Ambaliya and Masu (2018) in aonla.

Average fruit weight (g)

The data showed that the effect of micronutrients on average fruit weight are-is presented in Table 1. was found significant. The maximum average fruit weight (44.06 g) was observed with treatment T₈ (ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.2 %) which was statistically at par with treatment T₉, T₁₀, T₁₁, T₁₂, T₁₃ and T₁₄. Whereas, tThe minimum average fruit weight (34.40 g) was recorded with thein treatment T₁ (control).

~~Increased in fruit weight~~It might be due to the zinc micronutrients application which plays a vital role to promote starch formation, iron required to suitable cell enlargement and cell division and boron actively involved in transportation of carbohydrates in plants. Thus, the cumulative effect of combined treatment of Zn + Fe + B might have resulted higher fruit weight. The possible reason for increase in fruit weight by the micronutrients might be due to faster loading and mobilization of photo assimilates to fruits and involvement in cell division and cell expansion which ultimately reflected into more weight of fruit in treated plants (Brahamchari *et al.*, 1995). These findings are supported by the results obtained by Venu *et al.* (2016) in acid lime, Tariq *et al.* (2007) in sweet orange, Meena *et al.* (2016) in Nagpur mandarin, Nehete *et al.* (2011) and Razzaq *et al.* (2013) in mango, Modi *et al.* (2012) in papaya, Ningavva *et al.* (2014) in banana, Bhojar and Ramdevputra (2017) in guava and Ambaliya and Masu (2018) in aonla.

Fruit volume (cc)

The data regarding effect of micronutrients on fruit volume are presented in Table 1. Data showed that the effect of micronutrients on fruit volume was found significant. The maximum fruit volume (43.83 cc) was observed with treatment T₈ (ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.2 %) which was statistically at par with treatment T₉, T₁₀, T₁₁ and T₁₂. ~~Whereas~~However, the minimum fruit volume (32.96 cc) was recorded ~~with the in~~ treatment T₁ (control).

These might be due to the Zn plays vital role to promote starch formation, iron required to suitable cell enlargement and cell division and B actively involved in transportation of carbon in plants. Thus, the cumulative effect of combination of Zn + Fe + B on faster cell division and cell expansion reflected on higher fruit volume. Similar results were also noted by Venu *et al.* (2016) in acid lime, Tariq *et al.* (2007) in sweet orange, Meena *et al.* (2016) in Nagpur mandarin, Goswami *et al.* (2012), Jat and Kacha (2014) and Bhojar and Ramdevputra (2017) in guava, Yadav *et al.* (2014) in pomegranate and Ambaliya and Masu (2018) in aonla.

Number of fruits per tree

A perusal of data presented that effect of micronutrients on number of fruits per tree was found significant. The maximum number of fruits per tree (946.00) was observed with treatment T₈ (ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.2 %) which was statistically at par with treatment T₉, T₁₀, T₁₁ and T₁₂. ~~Whereas~~While, the minimum number of fruits per tree (698.83) was recorded ~~with the in~~ treatment T₁ (control).

Comment [E3]: Write number of fruits in one decimal

Comment [E4]: Write number of fruits in one decimal

All the micronutrients when sprayed alone or in combination involved directly in various physiological processes and enzymatic activity. This might have resulted into better photosynthesis, greater accumulation of starch in fruits and involvement of Zn in auxin synthesis and B in translocation of starch to fruits. The balance of auxin in plant regulates the fruit drop or retention in plants, which altered the control of fruit drop and increased the total number of fruits per tree. Similar findings have been observed by Venu *et al.* (2018) in acid lime, Gurjar and Rana (2014) and Reetika *et al.* (2018) in kinnow mandarin, Bhowmick *et al.* (2011) and Gurjar *et al.* (2015) in mango, Suman *et al.* (2016) in guava, Dhurve *et al.* (2018) in pomegranate and Ambaliya and Masu (2018) in aonla.

Fruit yield (kg per tree)

The data pertaining to effect of micronutrients on fruit yield are presented in Table 1. The maximum fruit yield (41.68 kg per tree) was observed with treatment T₈ (ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.2 %) which was statistically at par with treatment T₉, T₁₀ and T₁₁. Whereas, the minimum fruit yield (24.04 kg per tree) was recorded with the treatment T₁ (control).

An increased in fruit yield per tree might be due to cumulative effect of number of fruits, reduction in fruit drop and higher fruit weight by effect of foliar spray of micronutrients. Promotion of starch formation followed by rapid transportation of carbohydrates in plants is activated by Zn and B is well established. Iron (Fe) is highly associated with chlorophyll synthesis which later on boosted up the photosynthesis. Foliar spray of micronutrients might have affected the physiological processes resulting into higher yield. This finding is in the accordance with the result of Deshmuk *et al.* (2015) and Venu *et al.* (2018) in acid lime, Sajid *et al.* (2010) in sweet orange, Ilyas *et al.* (2015) in kinnow mandarin, Bhalerao and Patel (2015) in papaya, Gurjar *et al.* (2015) in mango, Suman *et al.* (2016) in guava, Abhijith *et al.* (2018), Ambaliya and Masu (2018) and Jangid *et al.* (2018) in aonla and Dhurve *et al.* (2018) in pomegranate.

Total fruit yield (quintal per hectare q ha⁻¹)

Data regarding to total fruit yield are presented in Table 1. Data showed that effect of micronutrients on total fruit yield was found significant. The maximum total fruit yield (115.46 quintal per hectare q ha⁻¹) was observed with treatment T₈ (ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.2 %) which was statistically at par with treatment T₉, T₁₀ and T₁₁. ~~Whereas~~ However, the

Formatted: Font: (Default) Times New Roman, 12 pt, Superscript

Formatted: Superscript

minimum total fruit yield (66.59 quintal per hectare) was recorded ~~with the~~in treatment T₁ (control).

The increase in total fruit yield by application of micronutrient treatments may be due to the direct or indirect involvement of nutrients which provide better mobilization of nutrients and metabolites for the growth and development of fruits by increase in metabolic activities and better cellular pathways. These activities improve their size, weight and volume, number of fruit and thereby synergistically increased the total fruit yield. These results are in confirmation with those of Deshmuk *et al.* (2015) and Venu *et al.* (2018) in acid lime, Sajid *et al.* (2010) in sweet orange, Ilyas *et al.* (2015) in kinnow mandarin, Gurjar *et al.* (2015) in mango, Suman *et al.* (2016) in guava and Abhijith *et al.* (2018) in aonla.


UNDER PEER REVIEW

Table 1: Effect of micronutrients on yield parameters (cm)

Code	Treatment	Fruit diameter (cm)	Average fruit weight (g)	Fruit volume (cc)	Number of fruits per tree	Fruit yield (kg per tree)	Total fruit yield (quintal per hectare $q\ ha^{-1}$)
T ₁	Control	3.50	34.40	32.96	698.83	24.04	66.59
T ₂	ZnSO ₄ 0.5 %	3.87	38.93	38.46	809.17	31.50	87.25
T ₃	ZnSO ₄ 1 %	3.72	36.89	36.42	774.67	28.58	79.16
T ₄	FeSO ₄ 0.5 %	3.80	38.04	37.96	797.17	30.33	84.01
T ₅	FeSO ₄ 1 %	3.68	36.21	35.87	758.83	27.48	76.11
T ₆	Borax 0.2 %	3.74	37.41	37.20	786.00	29.41	81.46
T ₇	Borax 0.4 %	3.61	35.90	35.28	752.83	27.03	74.86
T ₈	ZnSO ₄ 0.5 % + FeSO ₄ 0.5 % + Borax 0.2 %	4.58	44.06	43.83	946.00	41.68	115.46
T ₉	ZnSO ₄ 0.5 % + FeSO ₄ 1 % + Borax 0.2 %	4.38	43.15	42.58	901.00	38.88	107.68
T ₁₀	ZnSO ₄ 0.5 % + FeSO ₄ 0.5 % + Borax 0.4 %	4.32	42.79	41.96	881.67	37.72	104.49
T ₁₁	ZnSO ₄ 0.5 % + FeSO ₄ 1 % + Borax 0.4 %	4.28	42.03	41.06	868.67	36.51	101.13
T ₁₂	ZnSO ₄ 1 % + FeSO ₄ 0.5 % + Borax 0.2 %	4.18	41.45	40.87	851.67	35.30	97.79
T ₁₃	ZnSO ₄ 1 % + FeSO ₄ 1 % + Borax 0.2 %	4.13	41.15	40.29	847.17	34.92	96.72
T ₁₄	ZnSO ₄ 1 % + FeSO ₄ 0.5 % + Borax 0.4 %	4.08	40.43	39.25	834.00	33.72	93.40
T ₁₅	ZnSO ₄ 1 % + FeSO ₄ 1 % + Borax 0.4 %	3.96	39.04	38.82	821.83	32.08	88.87
	S.Em. ±	0.13	1.58	1.21	33.40	1.89	5.22
	C.D. (P = 0.05)	0.38	4.57	3.50	96.74	5.46	15.13
	C.V. %	5.69	6.93	5.38	7.04	10.01	10.01

Formatted: Superscript

UNDER PEER REVIEW



CONCLUSION

On the basis of experimental results, [it may be concluded that](#) the combined application of ZnSO₄ 0.5 % + FeSO₄ 0.5 % + Borax 0.2 % was found effective and promising ~~in~~ [for](#) increasing fruit diameter, fruit volume, fruit weight, number of fruits per tree, fruit yield per tree and total fruit yield per hectare of Acid Lime (*Citrus aurantifolia* Swingle) cv. Kagzi Lime.

REFERENCES

- Abhijith, Y. C.; Adiga, J. D.; Kishor, H. and Sindhu, C. (2018). Effect of micronutrients on yield and quality of aonla (*Embllica officinalis* Gaertn.) cv. NA-7. *International Journal of Current Microbiology and Applied Sciences*, **7**(3): 140-145.
- Agarwala, S. C. and Sharma, C. P. (1978). Plant nutrients- their functions and uptake. Soil fertility theory and practice, ICAR New Delhi, pp.26-32.
- Ambaliya, N. B. and Masu, M. M. (2018). Effect of multimicronutrient on fruit yield of aonla (*Embllica officinalis* Gaertn.) cv. Gujarat aonla-1. *International Journal of Chemical Studies*, **6**(5): 1359-1363.
- Asana, R. D.; Sarin, R. D. and Saxena, M. N. (1971). Handbook of manures and fertilization, ICAR New Delhi, pp. 160-94.
- Bhalerao, P. P. and Patel, B. N. (2015). Effect of foliar application of Ca, Zn, Fe and B on growth, yield and quality of papaya (*Carica papaya*) var. Taiwan Red Lady. *Indian Journal of Horticulture*, **72**(3): 325-328.
- Bhowmick, N. and Banik, B. C. (2011). Influence of pre-harvest foliar application of growth regulators and micronutrients on mango cv. Himsagar. *Indian Journal of Horticulture*, **68**(1): 13-107.
- Bhojar, M. G. and Ramdevputra, M. V. (2017). Effect of foliar spray of zinc, iron and boron on quality of guava (*Psidium guajava* L.) cv. Sardar L-49. *Trends in Biosciences*, **10**(39): 8109-8111.
- Brahmachari, V. S.; Mandal, A. K.; Kumar, R.; Rani Ruby (1995). Effect of growth substances on fruit-set and physico-chemical characteristics of Sardar guava (*Psidium guajava* L.). *Recent Horticulture*. **2**(2): 127-131.
- Debaje, P. P.; Shinde.; Ekta, D. and Ingale, H. V. (2011). Effect of plant growth regulators and nutrients on quality of Acid lime (*Citrus aurantifolia* Swingle). *The Asian Journal of Horticulture*, **6**(1): 253-255.
- Deshmukh, H. K.; Nimbolkar, P. K.; Paithankar, D. H. and Dewangan, R. K. (2015). Effect of plant growth regulators and micronutrients on growth and yield of acid lime (*Citrus aurantifolia* swingle) in hasta bahar. *International Journal of Agriculture, Environment and Biotechnology*, **8**(3): 615-620.
- Dhurve, M. K.; Sharma, T. R.; Bhooriya, M. S. and Lodha, G. (2018). Effect of foliar application of zinc and boron on growth, reproductive and yield of pomegranate cv. Ganesh in hast bahar. *International Journal of Chemical Studies*, **6**(5): 499-503.
- Goswami, A. K.; Shukla, H. S.; Kumar, P. and Mishra, D. S. (2012). Effect of pre harvest application of micronutrients on quality of Guava (*Psidium guajava* L.) cv. Sardar. *Horticulture Flora Research Spectrum*, **1**(1): 60-63.

- Gurjar, P. S. and G. S. Rana (2014). Influence of foliar application of nutrients and growth regulator on fruit drop, yield and fruit size and quality in Kinnow mandarin. *Indian Journal of Horticulture*, **71**(1): 109-111.
- Gurjar, T. D.; Patel, N. L.; Panchal, B. and Chaudhari, D. (2015). Effect of foliar spray of micronutrients on flowering and fruiting of Alphonso mango (*Mangifera indica* L.). *The Bioscan*, **10**(3): 1053-1056.
- Ilyas, A.; Ashraf, M. Y.; Hussain, M.; Ashraf, M.; Ahmed, R. and Ali, K. A. (2015). Effect of micronutrients (Zn, Cu and B) on photosynthetic and fruit yield attributes of *Citrus reticulata* Blanco var. Kinnow. *Pakistan Journal of Botany*, **47**(4): 1241-1247.
- Jangid, G.; Mandal, G. and Kumari, U. (2018). Effect of nutrients on yield and chemical characteristics of aonla (*Embllica officinalis* Gaertn) cv. Chakiya. *Journal of Medicinal Plants Studies*, **7**(1): 106-108.
- Jat, G. and Kacha, H. L. (2014). Response of guava to foliar application of urea and zinc on fruit set, yield and quality. *Journal of Agri Search*, **1**(2): 86-91.
- Meena, D.; Tiwari, R. and Singh, O. P. (2014). Effect of nutrient Spray on growth, fruit yield and quality of aonla. *Annals of plant and soil research*, **16**(3): 242-245.
- Meena, M. K.; Jain, M. C.; Singh, J.; Sharma, M.; Singh, B. and Maurya, I. B. (2016). Effect of pre- harvest spray of calcium nitrate, boric acid and zinc sulphate on yield and quality of Nagpur Mandarin (*C. reticulata* blanco). *International Journal of Current Research*, **8**(8): 36837-36842.
- Modi, P. K.; Varma, L. R.; Bhalerao, P. P.; Verma, P. and Khade, A. (2012). Micronutrient spray effects on growth, yield and quality of papaya (*Carica papaya* L.) cv. Madhu Bindu. *Madras Agricultural Journal*, **99**(7): 500-502.
- Nehete, D. S.; Padhiar, B. V.; Shah, N. I.; Bhalerao, P. P.; Kolambe, B. N. and Bhalerao, R. R. (2011). Influence of micronutrient spray on flowering, yield, quality and nutrient content in leaf of mango cv. Kesar. *The Asian Journal of Horticulture*, **6**(1): 63-67.
- Ningavva, B. V.; Kulapati, H.; Paramappa, M. K. and Nadukeri, S. (2014). Effect of soil application and foliar spray of zinc and boron on quality of ratoon banana cv. Grand Naine under hill zone of Karnataka. *Trends in Biosciences*, **7**(20): 3294-3296.
- Pandey, S. N. and Sinha, B. K. (2006) Plant Physi., (4th edition) Vikas Publication House, Nodia; pp 130-135.
- Prasad, H.; Prasad, D.; Bhan, C.; Bairwa, S. K.; Babu, S. and Pal, S. (2013). Effect of foliar application of urea, zinc sulphate, and 2,4-D on kinnow mandarin. *Journal of Progressive Agriculture*, **4**(1): 148-153.
- Razzaq, K.; Khan, A. S.; Malik, A. U.; Shahid, M. and Ullah, S. (2013). Foliar application of zinc influences the leaf mineral status, vegetative and reproductive growth, yield and fruit quality of Kinnow Mandarin. *Journal of Plant Nutrition*, **36**(10): 1479-1495.
- Reetika; Rana, G. S.; Rana, M. K.; Prince and Kant, G. (2018) Effect of foliar application of macro and micronutrients on fruit drop and yield of kinnow mandarin, *International Journal of Pure & Applied Bioscience*, **6**(2): 1163-1169.
- Ryugo, K. (1988). Fruit culture. John Wiley and Sons p. 259-261.
- Sajid, M.; Rab, A.; Ali, N.; Arif, M.; Ferguson, L. and Ahmed, M. (2010). Effect of foliar application of Zn and B on fruit production and physiological disorders in sweet orange cv. Blood orange. *Sarhad Journal of Agriculture*, **26**(3): 355-360.
- Suman, M.; Dubalgunde, S. V.; Poobalan, O. and Sangma, P. D. (2016). Effect of foliar application of micronutrients on yield and economics of guava (*Psidium guajava* L.) cv.

L-49. *International Journal of Agriculture, Environment and Biotechnology*, **9**(2): 221-224.

Tariq, M.; Sharif, M.; Shah, Z. and Khan, R. (2007). Effect of foliar application of micronutrients on the yield and quality of sweet orange (*Citrus sinensis* L.). *Pakistan Journal of Biological Sciences*, **10**(11): 18-23.

Trivedi, N.; Singh, D.; Bahadur, V.; Prasad, V. M. and Collis, J. P. (2012). Effect of foliar application of zinc and boron on yield and fruit quality of guava (*Psidium guajava* L.). *Horticulture Flora Research Spectrum*, **1**(3): 281-283.

Venu, A. and Delvadia, D. V. (2018). Effect of zinc, boron, and iron application on flowering and yield of acid lime (*Citrus aurantifolia* Swingle) cv. Kagzi lime. *International Journal of Microbiology Research*, **10**(4): 1164-1165.

Venu, A.; Delvadia, D. V.; Chitroda, R. L. and Bhalani, R. (2016). Effect of zinc, boron, and iron application on physio-chemical parameters of acid lime (*Citrus aurantifolia* Swingle) cv. Kagzi lime. *The Bioscan*, **11**(3): 1597-1599.

Yadav, V. K.; Jain, M. C.; Sharma, M. K.; Gupta, N. K. and Singh, J. (2014). Effect of micronutrients foliar feeding on growth and yield of pomegranate (*Punica granatum* L.) cv. Sindhuri. *Journal of Tropical Agriculture*, **32**(3-4): 469-473.

UNDER PEER REVIEW