

# BULKING BEHAVIOUR OF PROMISING POTATO VARIETIES IN NORTHERN REGION OF BANGLADESH

## ABSTRACT

The experiment was conducted at Breeder Seed Production Centre (BSPC), Debiganj, Panchagarh during the Rabi season of 2014-2015 to find out the suitable time of harvest for getting desirable yield of promising potato varieties. Six harvesting was done at 10 days interval started from 40 days after planting (DAP) and continued up to 90 DAP. Six promising potato varieties such as Bari Alu 40 (4.45w), Bari Alu 41 (5.183), Bari Alu42 (Agila), Bari Alu 43 (Atlas), Bari Alu 44 (Elgar) and Bari Alu 45 (Steffi) were included in the study. Tuber yield was increased up to 90 DAP for all the varieties. The highest tuber yield (55.86ton/ha) was recorded in Bari Alu 45 (Steffi) when harvested at 90 DAP. On the other hand, the lowest tuber yield was found in BariAlu43 (Atlas) when harvested at 90 DAP. The tuber yield increase rate of Bari Alu 45 (Steffi), Bari Alu 41 (5.183), Bari Alu 42 (Agila) and Bari Alu 40 (4.45w) were reasonable up to 90 DAP, while the increase rate of Bari Alu 43 (Atlas) was reasonable up to 80 DAP and Bari Alu 44 (Elgar) was reasonable up to 70 DAP. Bari Alu 44 (Elgar) may be harvested as early variety among the varieties. The percent dry matter was increased up to 90 DAP for all the varieties. The growth rate of tuber was higher at 60 – 70 DAP for all the varieties except Bari Alu 41 (5.183). In case of Bari Alu 41 (5.183) tuber growth rate was higher at 70-80 DAP.

*Keywords: Dry matter, Days to harvest, Potato, Tuber growth rate and Yield*

## 1. INTRODUCTION

In Bangladesh, the potato (*Solanum tuberosum* L.) is a significant cash crop (Rahman et al., 2019). In cool nations, it is also used as a food and cash crop. It can satisfy the demand for vegetables and supply the low-income population with the necessary nutrients (Islam et al., 2009; Miah and Hossain, 2012). In terms of production and area, it is the most widely grown vegetable in Bangladesh. However, despite being the seventh-largest producer of potatoes in the world, Bangladesh has a low output per unit area of this crop when compared to other nations that also cultivate potatoes (Islam et al., 2017; Anon, 2007). In Bangladesh, the average yield of potatoes is 19.55 t/ha, which is significantly less than the crop's potential productivity (BBS, 2013). A key consideration for choosing an early or late variety of potato, as well as one that would provide a higher yield, is the bulking behavior of the germplasm or variety. Throughout the entire growing period, the bulking rate varies from variety to variety. Numerous studies have attempted to comprehend the early bulking behavior of various varieties, and it has been reported that at the early harvest stage, a greater number of small-sized tubers are found (Rady, 1963; Songin and Gasirowska, 1980). Johanston et al., on the other hand, (1982) stated that varieties with fast bulking produced high-quality chips and those with slow bulking produced dark-colored chips. Additionally, the darkest chips were produced by varieties harvested late (Miller et al., 1975), but the tuber had more dry matter and starch (Rytel, 2009; Lisinska, 2006). However, the planting dates were reported to have an impact on the maximum marketable and total tuber yield as well as the number (Hassanpanah et al., 2009).

In addition to this, the quantity of starch and dry matter in the potato tubers is a key element in determining yield and the quality of processed potato products (Kumar et al., 2011). Furthermore, dry matter is a sign of higher processing quality because it reduces oil absorption and reduces frying time (Pavlista and Ojala, 1997) and better textured products (Genet, 1992) associated with high fat absorption at frying (Araújo et al. 2016). Moreover, to their genetic component, which affects the texture and nutritional value of the tubers, other factors that have an impact on tuber dry matter and starch content include soil type, fertilizer application, dates of planting and days to harvesting (Buono et al., 2009; Murniece et al., 2010; Hamouz et al., 2005; Herman et al., 1996; Marwaha et al., 2005).

**Comment [Gh1]:** Explain the general objective of study at first

**Comment [Gh2]:** Please state the month as known English one

**Comment [Gh3]:** What was the experiment design?

**Comment [Gh4]:** What was the main conclusion of study?

**Comment [Gh5]:** How is potato production across the world?

**Comment [Gh6]:** These references are so old

The primary goals of the current study, taking into account the aforementioned concerns, were to i) learn about the bulking behavior of promising potato varieties, ii) identify early varieties, and iii) determine the best time of harvesting for maximizing the production of large tubers needed for processing.

## 2. MATERIALS AND METHODS

The experiment was conducted at the BSPC Debiganj, Panchagarh during the **rabi** season of 2014-2015. Six promising varieties VIZ: Bari Alu 40 (4.45W), Bari Alu 41 (5.183), Bari Alu 42 (Agila), Bari Alu 43 (Atlas), Bari Alu 44 (Elgar), Bari Alu 45 (Steffi) and six harvesting time viz 40, 50, 60, 70, 80, and 90 days after planting (DAP) were used in the study. The two-factor experiment was laid out in a randomized complete block design (RCBD) with three replications. The unit plot was 3m x 1.8m with a spacing of 60cm x 25cm. Harvesting was done according to maturity of the crop. Fertilizer, Irrigation and necessary intercultural operations were done as per tuber crop research center (TCRC), Bangladesh Agricultural Research Institute (BARI) recommendation (Azad et al., 2012). Data on different growth yield and yield contributing characters were recorded. Data that were collected from the field experiments has been analyzed by MSTATC Program and means separation was done by Duncan's Multiple Range Test (DMRT).

**Comment [Gh7]:** State the season as English known month

**Comment [Gh8]:** Explain the factors

**Comment [Gh9]:** Please write the analysis data separately

## 3. RESULTS AND DISCUSSION

### 3.1 Effect of variety and date of harvest on the yield

The tuber yield of potato was significantly varied due to variety and harvesting time (Table-1). The highest tuber yield (55.86 ton/ha) was found in Bari Alu 45 (Steffi), which was statistically similar to Bari Alu 41 (5.183) and Bari Alu 42 (Agila), when potato was harvested at 90 DAP. The lowest tuber yield (40.24 ton/ha) was recorded in Bari Alu 43 (Atlas). Tuber yield of potato was increased sharply up to 70 DAP for all the varieties except Bari Alu 41 (5.183) and Bari Alu 45 (Steffi). After that increased rate was slightly lowered but continued. In case of Bari Alu 41 (5.183), tuber yield of potato was increased sharply up to 80 DAP and in case of Bari Alu 45 (Steffi), tuber yield of potato was increased sharply up to 90 DAP. Considerable accumulation of dry matter in the tuber after linear phase of bulking might be the reason of the higher yield at 90 DAP harvest compared to other days to harvest (Gregory, 1956), caused by significant reductions in the percentage of large tubers and the tuber specific gravity at early and advanced stage of the crop (Silva, 2004). The tuber yield increase rate of Bari Alu 45 (Steffi), Bari Alu 41 (5.183), Bari Alu 42 (Agila) and Bari Alu 40 (4.45w) were reasonable up to 90 DAP, while the increase rate of Bari Alu43 (Atlas) was reasonable up to 80 DAP and Bari Alu 44 (Elgar) was reasonable up to 70 DAP.

**Comment [Gh10]:** Please discuss more strongly about your results

**Comment [Gh11]:** This reference is so old.

**Table 1. Effect of Variety and date of harvest on the yield**

Variety	Yield (ton/ha)					
	40DAP	50DAP	60DAP	70DAP	80DAP	90DAP
V <sub>1</sub> =Bari Alu 40 (4.45w)	2.60	9.61	20.59	33.88	42.08	49.07
V <sub>2</sub> =Bari Alu 41 (5.183)	1.67	7.89	18.30	29.36	45.44	51.44
V <sub>3</sub> = Bari Alu 42 (Agila)	3.36	11.29	22.96	38.89	46.04	51.90
V <sub>4</sub> = Bari Alu 43 (Atlas)	3.81	10.23	19.29	29.23	35.24	40.24
V <sub>5</sub> = Bari Alu 44 (Elgar)	6.98	18.08	27.16	38.25	41.45	45.25
V <sub>6</sub> = Bari Alu 45 (Steffi)	3.18	11.73	24.50	36.77	45.82	55.86
CV (%)	9.76					

### 2.2 Effect of variety and date of harvest on the dry matter

Though early harvesting of potato tubers provides economic support to farmers but affects quality and tubers harvested at higher maturity stages contain more dry matter than immature ones (Misra et al., 1993). The dry matter of potato was significantly varied due to variety and harvesting time (Table 2).

The highest dry matter (20.48%) was found in Bari Alu 43 (Atlas), while the lowest dry matter (16.61%) was recorded in Bari Alu 45 (Steffi). Dry matter (%) was increased sharply up to 70 DAP for all the varieties, after that the increase rate was gradually decreased up to 90 DAP except Bari Alu 43 (Atlas) and Bari Alu45 (Steffi). In case of Bari Alu43 (Atlas) the dry matter gradually increased up to 90 DAP and in case of Bari Alu45 (Steffi), the dry matter gradually increased up to 80 DAP. According to Solaiman et al. (2015), dry matter was 21.72% higher at 110 DAP than it was at 80 DAP (16.94 percent), that was clearly reflected that the dry matter accumulation was increasing with higher harvest days which was similar to our findings in the present study.

**Table 2. Effects of Variety and date of harvest on the dry matter**

Variety	Dry matter (%)					
	40DAP	50DAP	60DAP	70DAP	80DAP	90DAP
V <sub>1</sub> =Bari Alu 40 (4.45w)	15.16	16.10	17.40	18.73	18.76	18.82
V <sub>2</sub> =Bari Alu 41 (5.183)	14.87	15.95	16.72	17.38	17.59	17.70
V <sub>3</sub> = Bari Alu 42 (Agila)	12.79	13.65	13.73	14.74	14.83	15.20
V <sub>4</sub> = Bari Alu 43 (Atlas)	14.01	15.81	17.32	18.93	19.40	20.48
V <sub>5</sub> = Bari Alu 44 (Elger)	12.47	13.51	15.74	16.25	16.89	16.95
V <sub>6</sub> = Bari Alu 45 (Steffi)	12.36	12.58	14.51	15.44	16.40	16.61
CV (%)	3.37					

### 2.3 Tuber growth rate (TGR) of six potato varieties at different DAP

Tuber growth rate (TGR) significantly varied among the varieties as well as in different growth stage (Table-3). The highest tuber growth rate (289.01 kg/ha/day) was found in Bari Alu41 (5.183) at 70-80 DAP. The lowest tuber growth rate (66.89 kg/ha/day) was recorded in Bari Alu44 (Elgar) at 80-90 DAP. Tuber growth rate was increased sharply up to 60-70 DAP for all the varieties. Contrary to our discovery, Sen et al (2014) achieved the highest tuber growth rate at 45-60 days per day, while Mahmoud (2012) found the highest TGR in Asterix at 50-60 days per day. After that it was declining except Bari Alu41 (5.183). In case of Bari Alu41 (5.183) tuber growth rate was increased up to 70-80 DAP, after that it was decreased. According to Klein Kopf et al., (2003), the maximum growth rate for each individual variety in the field can be determined physiological seed age, time of harvest and environmental conditions during tuber formation dedication.

**Table 3. Tuber growth rate (TGR) of six potato varieties at different DAP**

Variety	Tuber growth rate (Kg/ha/day)				
	40-50DAP	50-60DAP	60-70DAP	70-80DAP	80-90DAP
V <sub>1</sub> =Bari Alu 40 (4.45w)	115.30	203.54	276.30	154.84	134.07
V <sub>2</sub> =Bari Alu 41 (5.183)	101.01	180.13	204.30	289.01	111.19
V <sub>3</sub> = Bari Alu 42 (Agila)	111.13	161.13	257.99	109.53	106.10
V <sub>4</sub> = Bari Alu 43 (Atlas)	108.35	172.36	219.22	130.33	140.45
V <sub>5</sub> = Bari Alu 44 (Elger)	157.22	183.23	194.06	78.52	66.89
V <sub>6</sub> = Bari Alu 45 (Steffi)	108.25	207.93	212.23	183.72	176.38
CV (%)	38.75				

## 4. CONCLUSION

From the above discussion it may be concluded that Bari Alu 45 (Steffi) variety showed the highest yield (55.86 ton/ha). Bari Alu 41 (5.183) and Bari Alu 42 (Agila) gave the higher yield. The tuber yield increase rate of Bari Alu45 (Steffi), Bari Alu41 (5.183), Bari Alu 42 (Agila) and Bari Alu 40 (4.45w) were reasonable up to 90 DAP, while the increase rate of Bari Alu 43 (Atlas) was reasonable up to 80 DAP and Bari Alu 44 (Elger) was reasonable up to 70 DAP. Bari Alu 44 (Elger) may be harvested as early variety among the varieties. The dry matter (%) was increased up to 90 DAP for all the varieties. The tuber growth rate was higher at 60-70DAP for all the varieties except Bari Alu41 (5.183). In case of Bari Alu 41 (5.183) tuber growth rate was higher at 70-80 DAP, after that it was decreased. Ninety (90) DAP in the best harvesting time for getting large size of tuber and yield of potato. However, it

**Comment [Gh12]:** There is no need to repeat results here. What was the main conclusion.

needs more specific and systemic studies under different climatic condition to find out final recommendation.

## REFERENCES

**Comment [Gh13]:** References should be updated.

1. Anonymosu. Annual Report of Tuber Crops Research Centre published yearly TCRC. BARI. Joydebpur, Gazipur, Bangladesh. 2007.
2. Araújo, TH, Pádua JG, Spoto MH, Ortiz VD, Margossian PL, Dias CT, Melo PC. Productivity and quality of potato cultivars for processing as shoestrings and chips. *Horticultura Brasileira*. 2016; 34(4):554-560.
3. Azad AK, Wahab MA, Shaha MG, Nesa J, Rahman ML, Rahman MHH, Al-Amin M. *Krishi Prajukti Hatboi (Handbook on Agro-Technology)*. Bangladesh Agricultural Research Institute, Gazipur-1701, Bangladesh. 2012.
4. BBS. The year Book of Agricultural statistics of Bangladesh. Stat. Div. Minis. Planning, Govt. Peoples Repub. Bangladesh, Dhaka. 2013.
5. Buono V, Paradiso A, Serio F, Gonnella M, De Gara L, Santamaria P. Tuber quality and nutritional components of "early" potato subjected to chemical haulm desiccation. *J of food composition and analysis*. 2009;22(6):556-562.
6. Gasiórowska E, Songins W. The effects on seed tuber yield of the date of harvesting potatoes of the varieties Krab, Notee, Nysa and Proсна. *Biuletyn Instytutu Ziemmiaka*, 1980;25:25-34
7. Genet RA. Potatoes-the quest for processing quality. In *Proceedings of the Agronomy Society of New Zealand*. 1992;22:3-7.
8. Gregory LE. Some factors for tuberization in the potato plant. *American J of Botany*. 1956;43(4): 281-288.
9. Hamouz K, Lachman J, Dvorák P, Pivec V. The effect of ecological growing on the potatoes yield and quality. *Plant Soil and Environment*. 2005;51(9): 397.
10. Hassanpanah D, Ho sienzadeh AA, Allahyari N, 2009. Evaluation of planting date effects on yield and yield components of Savalan and Agria cultivars in Ardabil region. *Journal of Food, Agriculture & Environment*. 2009;7(3&4):525-528.
11. Herrman TJ, Love SL, Shafii B, Dwelle RB. 1996. Chipping performance of three processing potato cultivars during long-term storage at two temperature regimes. *Amer Potato J*. 1996;73(9): 411-425.
12. Hossain MA, Miah MAM. Post Harvest Losses and Technical Efficiency of potato storage systems in Bangladesh. 2012.
13. Islam MZ, Zaman MM, Hossain MM, Hossain A. Integrated nutrient management with liming for potato production in North-West region of Bangladesh. *Annual Report 2008-2009, Tuber Crops Research Centre, Bangladesesh Agricultural Res. Inst, Gazipur, Bangladesh*. 2009.
14. Islam S, Siddique MA, Rahman L, Saha AK, Nahiyen ASM, Raihan A. Study on field performance of six table potato varieties at debiganj ASRBC station. *Bangladesh Hort*. 2017;3(01):47-52.
15. Johnston GR, Rowberry RG. Determination of tuber sizing and accumulation of total solids contents of four potato varieties harvested at several dates. *Amer. Potato J*. 1962;39(1):29-35
16. Johnston GR, Rowberry RG. Science, a new early maturing, chipping potato variety. *Amer Potato J*. 1982;59(1): 39-43
17. Kleinkopf GE, Brand TL, Olsen N. Physiology of tuber bulking. *Idaho Potato Conference*. January 23, 2003.
18. Kumar M, Baishya LK, Ghosh DC, Gupta VK. 2011. Yield and quality of potato (*Solanum tuberosum*) tubers as influenced by nutrient sources under rainfed condition of Meghalaya. *Indian J of Agrono*. 2011;56(3): 260.
19. Lisińska G. Technological and nutritive value of the Polish potato cultivars. *Zeszyty Problemowe Postepow Nauk Rolniczych*. 2006;511:81-91.
20. Mahmud AA. Improvement of drought tolerant potato varieties. An unpublished Ph. D. Dissertation. Department of Horticulture, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Salna, Gazipur. 2012.

21. Marwaha RS, Pandey SK, Singh SV, Khurana SP. Processing and nutritional qualities of Indian and exotic potato cultivars as influenced by harvest date, tuber curing, pre-storage holding period, storage and reconditioning under short days. *Adv in Hort Sci.* 2005;30:140.
22. Millar RA, Harrington JD, Dukn GD. Effect of variety and harvest date on tuber sugars and chips colour. *Amer Potato J.* 1975;52 (12): 379-382
23. Misra JB, Anand SK, Chand P. Changes in processing characteristics and protein content of potato tubers with crop maturity. *J of Indian Potato Assoc.* 1993;20:150-154.
24. Murniece I, Karklina D, Galoburda R, Sabovics M. 2010. Reducing sugar content and colour intensity of fried Latvian potato varieties. *Latvijas Lauksaimniecības Universitātes Raksti.* 2010;24: 20-30.
25. Pavlista, AD, Ojala JC. Potatoes: Chip and French fry processing. *Processing Vegetables, Science and Technology.* Lancaster: Technomic Publishing, Inc. 1995;237-284.
26. Rahman MW, Islam S, Islam MM, Hossain MS. Effect of Sulphur on the Yield and Quality of Potato Varieties in Tista Meander Floodplain Soil of Bangladesh. *Inter J of Plant & Soil Sci.* 2019;30(4):1-8.
27. Rytel E. Influence of potato maturity on changes of non-starch polysaccharides and lignin content in tubers. *Zeszyty Problemowe Postepow Nauk Rolniczych.* 2004;500: 295-303.
28. Sen D, Rakshit A, Ghosh DC. 2014. Effect of transplanting dates and plant population on growth parameters of potato (*Solanum tuberosum* L.) raised from true potato seed (TPS). *Cercetări agronomice în moldova.* 2014;XLVII(1):97-106.
29. Silva LAS. Duração do ciclo vegetativo e sua relação com o potencial produtivo de genótipos de batata (Doctoral dissertation, Universidade Federal de Lavras.). 2004.
30. Solaiman AHM, Nishizawa T, Roy TS, Rahman M, Chakraborty R, Choudhury J, Hasanuzzama M. 2015. Yield, Dry Matter, Specific Gravity and Color of Three Bangladeshi Local Potato Cultivars as Influenced by Stage of Maturity. *J of Plant Sci.* 2015;10(3):108-115.