

Original Research Article

Effect of land use change on total phosphorus and its fractions in North-Western Himalayas

ABSTRACT

Aims: Conversion of land from forest to cropping has a serious effect on soil phosphorus and its fractions

Results: Land use is now widely understood to be a primary factor in environmental change across all time and space scales. The purpose of this research was to ascertain how different land uses affected the concentration of phosphorus in soil. Soil phosphorus (P) reserves are depleted when land is converted from natural vegetation to permanent agricultural cropping. The transformation of North-Western Himalayas from a forest-dominated to a grassland-dominated ecosystem is just one example of the diversity of land significantly less soil aggregation occurred when agricultural land was cleared of its native vegetation. Total organic carbon in soils was reduced when grassland was converted to cropland. Reduced total organic carbon (TOC) concentrations by 62-79% and organic phosphorus (Po) concentrations by 47-53%. Even though, the total silt+clay fraction's contribution was negligible, it contained a significant amount of C and Po reserves and the C/Po ratio has been holding fairly steady, they have proven to be more robust. This impact of cropping on soil P reserves has been demonstrated in research, but changing land use practices can alleviate these problems significantly.

Keywords: Land use, Phosphorus fractions, Pasture

1. INTRODUCTION

Phosphorus (P) is a macronutrient that plants need to grow and flourish. However, it is believed that about 5.7 billion hectares (ha) of land around the world has insufficient amounts of plant-available P [1]. Phosphate rock is used in crop production at the current rate of about 150 million tons per year [2]. In addition, global reserves of P are the smallest of the 14 essential plant nutrients [3], and the vast majority of them are located in just one country (Morocco). Understanding P behavior in soils, including its effect on long-term agricultural production, is thus crucial. Despite total P concentrations between 200 and 800 mg kg⁻¹ soil [4], only 0.1% of this P is available for plant uptake in the soil solution [5]. Since plants uptake P from the soil solution, which has low P concentrations, that fraction must be replenished to meet plant needs. Labile P replenishes. The first labile pool (Pi) is P in mineral forms that dissolve, releasing P, or P adsorbing to soil surfaces. Plant and enzyme activity mineralizes the second labile pool (Po) [6].

Long-term cultivation and cropping reduce soil P concentrations in low-input agricultural systems [7-9]. It was reported that both inorganic P and Po concentrations reduced significantly. However, much remains unknown about cropping, along with how land management techniques affect P loss and how soil aggregates sustain cycle P pools, particularly Po [9]. Only few studies have examined its effects on P forms. Fontaine et al. concluded that despite a 40% increased Po content in productive pasture soils, there wasn't any difference in P content.

2. MATERIAL AND METHODS

All the soil samples were air-dried, polished and pulverized using mortar and pestle and then sieved through a 2mm mesh sieve. The hydrometer method [11] was used to determine particle size distribution. Bl

Comment [H1]: The author should include A BRIEF DESCRIPTION AND A MAP OF THE STUDY AREA. In addition, a description of the method for selection of sites and procedure for soil sampling is required.

How many sites were sampled?

If the main aim of the study is to compare changes in P content in soils as a result of changes in land use, the methodology should include sites with different land uses at same time other soil factors remain constant, such as climate, geology and relief

Hartge's[12]method was used to estimate soil bulk density in which each core of soil sample was oven-dried at 105 °C for at least two to three days. The soil pH was determined potentiometrically 1:2.5 (w/v) soil-water supernatant suspension[13]. SOC was estimated using [14]. Citrate-dithionite-extractable Fe a2d Al were determined using method of Kuo[15]. Concentrations of P within the soil samples were examined using Colwell-P as well as sequential fractionation. For Colwell-P, concentrations were determined following extraction with NaHCO₃[16], with P measured using the molybdate blue colorimetric method [17]. Sequential extraction of P was conducted following a modified Hedley fractionation method[18].

Comment [H2]: This is a section for laboratory procedures.

3. RESULTS

Overall, the pH ranged from 5.5 to 6.8 in the soils of different land uses (Table 1). Soil pH levels were typically lower in areas with forest compared to those in areas where the land had been converted to agriculture (Table 1). The EC of the soils tested ranged from 0.01 to 0.21 dS m⁻¹, indicating no salinity hazard. A high percentage of clay was present in all soils, with values ranging from 25 % to 43 % (Fig.1). Average concentration of SOC ranged from 3.2 to 10.3 g Kg⁻¹ across all land uses with highest in forest soils and lowest in paddy soils. Total nitrogen varied between 0.4 to 7.7 g Kg⁻¹ with highest in forest soils and lowest in paddy soils. Fe_a varied between the land uses from 1.2-13.5 g Kg⁻¹ (Fig.2) with lowest in paddy soils and highest in forest soils, similar trend was found in case of Al_d among all the land uses (Table 1).

Table 1. Physico-chemical properties under different land uses

Land use	pH	EC (dSm ⁻¹)	Bulk Density (Mgm ⁻³)	Sand (%)	Silt (%)	Clay (%)	SOC (g Kg ⁻¹)	Total Nitrogen (g Kg ⁻¹)	Fe _a (g Kg ⁻¹)	Al _d (g Kg ⁻¹)
Forest	5.5	0.07	1.15	20	15	25	10.3	7.7	13.5	8
Pasture	5.8	0.01	1.21	15	20	41	6.7	5.6	9.4	3.2
Paddy	6.8	0.15	1.42	24	25	30	3.2	0.4	1.2	2.1
Orchard	6.7	0.21	1.51	30	30	43	4.1	1.1	1.6	2.8




Fig. 1 Mechanical analysis of different land uses




Fig. 2 Chemical properties of different land uses

Total P ranged from 526 to 901 mg kg⁻¹ in studied soils. Pasture soils had the highest total P concentration of 901 mg kg⁻¹ (Table 2). 65-70 % of P was Pi and 25-50 % Po in all the land uses. Converting forest soils to agricultural production (cropping) affected these P fractions. First, transition of soil to cropping reduced total P concentrations. However, conversion of forest soils to other land uses decreased Po's contribution to total P. In all soils, the C/Po ratio (for Po) also decreased significantly for all land uses for all three soils. Finally, transition from forest to cropping reduced Colwell-P concentrations. These P fractions within the soil were influenced by the transition from grazing to cropping, though the extent of this influence varied across land uses. At first, it was discovered that when land was used for farming (cropping), total P concentrations fell. No matter the land use, it was obvious that conversion significantly reduced the contribution of Po to total P. Similar declines in C/Po ratio (for Po) were observed across all land uses. For any given shift in land use, bulk soil Pi/Po ratios rose, with cropping leading the way and plantation showing the least change. Furthermore, Colwell-P concentrations declined dramatically across all land types.

Table 2 P concentrations influenced by land use, including for total P, inorganic P (Pi), organic P (Po), and the Po as a proportion of total P

Land use	Total P (mg Kg ⁻¹)	Total Pi (mg Kg ⁻¹)	Total Po (mg Kg ⁻¹)	Po/Total P (%)	Colwell-P (mg Kg ⁻¹)
Forest	813.2	536.2	439.1	54	65
Pasture	901.5	665.5	527.5	59	53
Paddy	526.4	402.5	125.6	23	36
Orchard	672.1	485.5	275.3	41	42

Comment [H3]: The author should explain why the sum of the Pi and Po results in higher values that reported as Total P in the Table 2.




Fig. 3 Phosphorus fractions under different land uses

DISCUSSION

Changes in both total P and Pi were found to be highly site-specific. However, cropping consistently reduced Po compared to soil with forest soils Kopittke et al. (19), who analyzed five Australian cropping soils, found that Po decreased significantly for four of them, on average by 28%. Crews and Brookes (20) found that after 116 years of cropping without fertilizer addition, Pi and Po did not change in the surface layer but showed depletion deeper within the soil profile. At the **Lamington site**, total organic carbon (TOC) decreased by 45 percent after land was converted from native vegetation to pasture, but neither total P nor P fractionation were affected. Townsend et al (21) studied the effects of pasture age on TOC and Po in an Oxisol, an Entisol, and a Mollisol from Brazil and Costa Rica and found similar results. For **Kingaroy**, however, a shift from cropland to pasture led to a 94% increase in both total P and Po. Although Pi decreased by 31%, the total P and Po were barely impacted by the conversion of this pasture to a plantation (Figure 2). Previous research has shown that after afforestation, total P and Po often decrease while plant-available Pi can increase Maranguit(22).

Before being used for grazing, the soil in the pastures was cropped. Firstly, it was seen that the Po fraction was reduced by 63% after cropping, but then increased to almost the same levels as in the soil with native vegetation after being converted to pasture or plantation. Switching from grazing to cropping resulted in a significant drop in total organic carbon (85% in the bulk soils due to net mineralization. Total organic carbon (TOC) concentrations were higher in soil that had been converted from cropland to pasture, but not to pre-farm levels. Finally, related to the preceding points, we noticed that converting cropland to pasture also partially restored values for C/Po and Pi/o to levels closer to those for the soil with forest landuse.

4. CONCLUSION

We analyzed how different types of land use affected the amount, fractionation, and distribution of phosphorus throughout the North-Western Himalayas. In addition to a 45% drop in soil C, we discovered that shifting land uses significantly altered P distribution. Changing the land use from forested to agricultural

Comment [H4]: This increase in total P and Po was without phosphorus fertilizer application?

Comment [H5]: The author should discuss why possibly is the sustain for pasture prove those increases.

resulted in significant reductions in total organic carbon, total nitrogen, and phosphorus. More research into the impact of management practice on increasing P availability in these soils is needed.

REFERENCES

1. Rayne, N. and L. Aula, *Livestock manure and the impacts on soil health: A review*. Soil Systems, 2020. **4**(4): p. 64.
2. Hellal, F., et al., *Importance of phosphate rock application for sustaining agricultural production in Egypt*. Bulletin of the National Research Centre, 2019. **43**(1): p. 1-11.
3. Cordell, D., J.-O. Drangert, and S. White, *The story of phosphorus: global food security and food for thought*. Global environmental change, 2009. **19**(2): p. 292-305.
4. Johnston, A. and P. Poulton, *Phosphorus in agriculture: A review of results from 175 years of research at Rothamsted, UK*. Journal of environmental quality, 2019. **48**(5): p. 1133-1144.
5. Naikoo, N.B., et al., *Aggregation-dependent phosphorus adsorption under different land uses of district Kupwara of Kashmir Valley*. Journal of Plant Nutrition and Soil Science, 2022.
6. Borie, F., et al., *Revisiting the nature of phosphorus pools in Chilean volcanic soils as a basis for arbuscular mycorrhizal management in plant P acquisition*. Journal of Soil Science and Plant Nutrition, 2019. **19**(2): p. 390-401.
7. Mäder, P., et al., *Arbuscular mycorrhizae in a long-term field trial comparing low-input (organic, biological) and high-input (conventional) farming systems in a crop rotation*. Biology and fertility of Soils, 2000. **31**(2): p. 150-156.
8. Stinner, B.R. and J.M. Blair, *Ecological and agronomic characteristics of innovative cropping systems*, in *Sustainable agricultural systems*. 2020, CRC Press. p. 123-140.
9. O'Neil, J.M., et al., *The rise of harmful cyanobacteria blooms: the potential roles of eutrophication and climate change*. Harmful algae, 2012. **14**: p. 313-334.
10. Fonte, S.J., et al., *Pasture degradation impacts soil phosphorus storage via changes to aggregate-associated soil organic matter in highly weathered tropical soils*. Soil Biology and Biochemistry, 2014. **68**: p. 150-157.
11. Bouyoucos, G.J., *Hydrometer method improved for making particle size analyses of soils 1*. Agronomy journal, 1962. **54**(5): p. 464-465.
12. Blake, G. and K. Hartge, *Peterson, RG, and LD Calvin. 1986. Bulk density Chapter 13 pp. 363-376*. Klute, A. Methods of soil analysis. Part, 1986. **1**.
13. Jackson, M.L., *Soil chemical analysis-advanced course*. Soil Chemical Analysis-Advanced Course., 1969(Edition 2).
14. Walkley, A. and I.A. Black, *An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method*. Soil science, 1934. **37**(1): p. 29-38.
15. Kuo, S., *Phosphorus*. Methods of soil analysis, 1996: p. 869-919.
16. Colwell, J., *The estimation of the phosphorus fertilizer requirements of wheat in southern New South Wales by soil analysis*. Australian Journal of Experimental Agriculture, 1963. **3**(10): p. 190-197.
17. Murphy, J. and J. Riley, *Colorimetric method for determination of P in soil solution*. Anal. Chim. Acta, 1962. **27**: p. 31-36.
18. Hedley, M., J. Stewart, and B. Chauhan, *Changes in inorganic and organic soil phosphorus fractions induced by cultivation practices and by laboratory incubations*. Soil Science Society of America Journal, 1982. **46**(5): p. 970-976.]

19. Kopittke, P. M., Dalal, R. C., Hoeschen, C., Li, C., Menzies, N. W., & Mueller, C. W. (2020). Soil organic matter is stabilized by organo-mineral associations through two key processes: The role of the carbon to nitrogen ratio. *Geoderma*, 357, 113974.
20. Crews, T. E., & Brookes, P. C. (2014). Changes in soil phosphorus forms through time in perennial versus annual agroecosystems. *Agriculture, ecosystems & environment*, 184, 168-181.
21. Townsend, A. R., Asner, G. P., Cleveland, C. C., Lifer, M. E., & Bustamante, M. M. (2002). Unexpected changes in soil phosphorus dynamics along pasture chronosequences in the humid tropics. *Journal of Geophysical Research: Atmospheres*, 107(D20), LBA-34.
22. Maranguit, D., Guillaume, T., & Kuzyakov, Y. (2017). Effects of flooding on phosphorus and iron mobilization in highly weathered soils under different land-use types: Short-term effects and mechanisms. *Catena*, 158, 161-170.

UNDER PEER REVIEW