

Distribution of microorganisms and fractionation of ~~sulfur~~sulphur in anthropogenic wetlands under long-term elevated CO₂(eCO₂) soil

Comment [DRY1]: Follow same English style either British or American
Most frequently used "sulphur" in British English

Formatted: Justified

ABSTRACT

The fate of extra carbon accessing soil under elevated CO₂ levels, as well as the repercussions for plant nutrition, is primarily determined by soil microbe activity. However, most increased CO₂ research has reported changes usually increases in soil organic carbon and reduction in the pH of the soil which is merely the first step in understanding how soil processes are altered. We analyzed these variables by assessing enzyme activities and identifying the individual components impacted by high CO₂ and those that reflect changes in soil organic matter pools. The majority of the microbial variables studied showed a significant increase under eCO₂ conditions. The rise in dehydrogenase activity suggests that the increased biomass of bacteria coincided with an increase in their activity. The rise in phosphatase activities implies that organic matter breakdown is being stimulated overall. The ~~sulfur~~sulphur fractions had a significantly increased number of substrates consumed by soil microorganisms under increasing CO₂. ~~Furthermore~~Moreover, direct examination of data from these perspective steep shifts in soil biological activity points to possible areas of investigation

Keywords: *Elevated CO₂, Soil organic matter, ~~Sulfur~~Sulphur fractions, Microbial distribution, Soil dehydrogenase.*

1. INTRODUCTION

The world population is predicted to reach 9.1 billion by 2050, requiring a 60-110 % increase in the food supply [1]. Which would ~~necessitate demand~~ agricultural productivity enhancement given the restricted scope of arable land growth. Food security is a serious concern in nations with an increasing population and shrinking cultivable land space. Rice (*Oryza sativa* L.) ~~is one of the most important staple crops, feeding almost half of the world population~~the world's most significant cereal crop. ~~It has fed more people over a longer period of time than any other crop.~~ Rice supplies more than half of the dietary requirements for 1.3 billion people, while rice provides 25-50 percent of the overall food requirements for 400 million individuals [2]. It is also a key source of calorie intake and nutrition, accounting for more than one-fifth of the calories ingested by humans globally. Rice is India's most significant staple food, consumed by around 65 percent of the population [3]. It is grown in almost all states, but West Bengal (13.79 percent), Uttar Pradesh (13.34 percent), Andhra Pradesh including Telangana (12.84 percent), Punjab (11.01 percent), Odisha (6.28 percent), Chhattisgarh (5.61 percent), Tamil Nadu (5.54 percent), Bihar (5.19 percent), Assam (4.41 percent), Haryana (3.88 percent), and Madhya Pradesh are the major rice producing states (3.86 percent).and it provides two-thirds of a person's calorie supply and one-half of total protein consumption [4]. Rice is special in that it can be produced in damp conditions when many other crops cannot. Such damp areas are common across Asia. However, soil fertility and fertilizer control are critical to a strong rice crop. Carbon and nutrient dynamics, which vary throughout agricultural ecosystems, are critical in soil fertility and crop yield [5]. As global temperatures rise due to greenhouse gas (GHG) emissions, sustainable agricultural output will be a major challenge in the future [6]. CO₂ is a significant GHG, accounting for approximately 60% of the entire greenhouse impact [7].

Formatted: Space Before: 0 pt, After: 8 pt

Comment [DRY2]: Authority is not in italics

Formatted: Font: Not Italic

CO₂ emissions have increased in recent decades, elevating the atmospheric CO₂ level from 280 ppm at the start of the industrial revolution to 410 ppm now [8]. Human actions like fossil fuel combustion, indiscriminate land use, land cover, and so on are to blame for rising GHG concentrations in the atmosphere. In addition to water vapor and solar irradiance, well-mixed GHGs such as CO₂, nitrous oxide (N₂O), methane (CH₄), and halocarbons contribute to global temperature rises and climate change [9]. Global surface temperature is rising significantly, and crops may be unable to be cultivated in the future if such events continue [10]. According to an international team of 17 academics from 17 countries, temperature rises due to global warming could subsequently be double what climate models assess. Sustained CO₂ increases in the atmosphere would have an effect on soil and plant nutrition. [11]

Comment [DRY3]: Please add recent reference it may be increased nearby 420 ppm

Sulphur, one of the most essential components for all plants and animals, is regarded as the fourth most important nutrient for agricultural crop development after nitrogen, phosphorus, and potassium. The relevance of sulphur in Indian agriculture is growing due to the recognition of its function in improving crop yield, not only for oil seeds, pulses, legumes, and forages but also for numerous kinds of cereal [12]. The sulphur deficit in crops is becoming more common as a result of the continued use of sulphur-free fertilizers, high-yielding crop types, intense multiple cropping systems, and increased productivity. Sulphur deficiency affects around 41% of India's farmed land. In India, crops remove around 1.26 million tons of sulphur, while fertilizers restore approximately 0.76 million tonnes [13]. Furthermore, the recovery of additional sulphur from external sources is approximately 8-10%. The continued depletion of local sulphur deposits throughout the post-green revolution period has resulted in a sulphur deficit in many sections of the country [14]. Sulphur exists in both inorganic and organic forms in soils, and the proportion of inorganic to organic sulphur varies greatly depending on the composition of the soil, its depth, and the management variables to which the soil is subjected. In most arable land top soils, total soil sulphur, which includes inorganic and organic binding forms, varies between 250 and 2500 kg/ha. Water soluble and adsorbed SO_4^{2-} composes inorganic "S" which is widely assumed to be the direct source of plants. In general, it accounts for less than 5% of total soil S. SO_4^{2-} is present in soil solutions only present in trace amounts that vary continually, and its concentration at any one moment is determined by the balance of plant uptake, and fertilizer input, mineralization, and immobilization. Understanding the S transformations in connection to crop availability in soil is required to utilize S fertilizers more efficiently.

Formatted: Superscript

Recent studies reveal that if temperatures rise by more than 0.5°C, the effects will be disastrous for millions of the world's poorest people due to reduced food output. As eCO_2 causes, an increase in global temperature in the future, soil health, nutrient availability, and crop output are all expected to suffer. Adaptation studies are being conducted all around the world, although they are often insufficient. India is far behind in terms of studies on the effects of climate change on agriculture. As a result, the current study was carried out to investigate the impacts of eCO_2 on the dynamics of microbes and sulphur fractions in the soil.

2. MATERIALS AND METHODS

2.1 Site description

The study was conducted in the A1 field in the Wetland Farms, TamilNadu Agricultural University, Coimbatore, India (11°005'13"N, 76° 93'045"E) which is under long-term elevated (eCO_2) 550±30 for a decade. Soil sampling was conducted on a randomized quadrant basis during the year 2018. A total of three samples with three replications have been collected from (i) Long-term elevated CO_2 open-top chamber (OTC) (eCO_2) (ii) OTC without any CO_2 , ambient (aCO_2), and (iii) Soil samples from a normal low-land rice ecosystem in the ambient bulk soil (ABS).

2.2 Enumeration of functional guilds

Enumeration of different functional bacterial groups. The media employed for the isolation of sulphur-oxidizing bacteria is mineral salt thiosulphate medium [15]. For the enumeration of methanogens, the [16] method was followed. For the enumeration of sulphur-reducing bacteria by [17] method. Methanotrophs enumerated by NMS medium. Diazotrophs and heterotrophs were enumerated by the method given by [18].

2.3. Soil enzyme analysis

All enzyme activities can be carried out by processing the soil size to < 2mm and following the colorimetric method for the analysis of the dehydrogenase [19], acid and alkaline phosphatase by the method given by [20] beta-glucosidase [21] in the soil was estimated by the standard method given by [24], aryl sulphatase by [22], and fluorescent diacetate hydrolysis was carried out by the standard method given by (Schnrer and Rosswall, 1982).

Comment [DRY4]: Follow journal format for reference

2.4. Sulphur fractions in soil

Fractionation of Sulphur

Comment [DRY5]: Should be revise according to journal format.

Less than 2mm of soil samples were used in the sequential extraction of soil **sulfursulphur** fractions [24]. The smaller fractions (water soluble) were extracted with demineralized water for the first time at (w/v) 1:10. After 30 minutes of shaking, each sample was analyzed. The sample was spun at 10,000 rpm for 10 minutes, and the homogenate was assessed using the turbidity method. Exchangeable **sulfursulphur** fractions were extracted at a 1:10 ratio with 0.032M NaH₂PO₄ (w/v) proportion after 30 minutes of shaking, each sample was centrifuged at 10,000 rpm for 10 minutes and the supernatant was analyzed by turbidity method. For precipitated **sulfursulphur**, the remainder of the soil sample was extracted by 1M HCl. The ratio of extraction was (w/v) 1:20. After 60 minutes of shaking, the samples were ready centrifuged at 10,000 rpm for 10 minutes, and analyzed by turbidity method.

Comment [DRY6]: ??

2.5. Available **sulfursulphur**

Five grams of soil was taken in a polythene shaking bottle and put it in a shaker container. 25 ml 0.15 percent CaCl₂ solution, shake for 30 minutes, filter with Whatman no.42 filter paper, pipette out 5 ml extract in 25 ml volumetric flask, then add 10 ml sodium acetate and acetic acid buffer after that, add 1 gram of barium chloride crystals and mix for 1 minute. After that, add 1 ml of 0.25 percent gum acacia solution and dilute to 25 ml with distilled water and shake until the barium chloride crystals dissolve 2-8 minutes after 420nm absorption. Calculate the concentration of accessible **sulfursulphur** in the soil sample by plotting the graph with a standard curve using K₂SO₄.

2.6. Statistical analysis

All the datasets recorded were analyzed by SPSS version 16.0. means of values compared by the least significant difference (LSD) at a 5 % level of probability

3. RESULTS AND DISCUSSION

3.1 Effect of eCO₂ concentrations on soil chemical properties. Soil pH, electrical conductivity, and Organic carbon in soil.

The pH of the soil **was decreased fell** as CO₂ concentration rise (Table 1). The lowest pH value (7.9) was **discovered-noticed** with 550 ppm eCO₂, while the highest pH value (8.2) was observed in ambient bulk soil. According to [25], eCO₂ is anticipated to raise CO₂ concentration in soil via the generation of H₂CO₃, and hence pH in the soil is likely to drop. The majority of plant nutrients become accessible when soil pH varies from 6.5 to 8.0. When submerged, the pH of the soil normally reaches 7.0. According to [26], an increase in CO₂ concentration affects **sed** soil pH. However, investigations have revealed that when a large quantity of CO₂ was added to the soil system, the pH changed insignificantly. It is also insignificant when compared to the depth of the soil. This may be due to the solubility of carbon dioxide (CO₂) in water, which results in the formation of undissociated carbonic acid. Acidic root exudates from plants and other microorganisms, on the other hand, lower pH and alter the solubility relationship of many plant nutrients. Acetic acid is the most common anaerobic respiration intermediate. During the decomposition process, all organic matter, including fats, amino acids, and carbohydrates, is converted to acetic acid. [27–29]. An increase in the eCO₂ increases the soil's organic carbon. The soil organic carbon was found higher in the eCO₂ range of 6.9 mg/kg⁻¹ while the lowest was observed in the ambient bulk soil range of 5.7 mg/kg⁻¹. These findings support [30]. The suggestion is that physical fractionation techniques can aid in identifying changes in the soil carbon pool under higher CO₂ conditions, even in trials with high intrinsic variability and beginning C content, as in our experiment. The increased sensitivity of SOC to changes in C dynamics with rising CO₂ than total C is most likely due to considerably quicker turnover in SOM pools (Table-1&3). The greater proportion of new SOC relative to total new soil versus the proportion of total SOC related to total new soil C indicates a quicker turnover of SOC. SOC content primarily through boosting photosynthate synthesis and allocation to the rhizosphere, hence increasing C intake into the soil [31]. Soil organic matter is a key factor that affects all physical, chemical, and biological activity in soils. It is soil's essence that enhances soil health, agricultural productivity, and soil aggregates. More C sequestration in soils equals lower CO₂ levels in the atmosphere, reducing the potential for global warming [32].

Comment [DRY7]: mg/kg

Comment [DRY8]: mg/kg

Comment [DRY9]: If possible add some recent reference

Table 1. Soil physical properties from three different soils elevated, ambient, and ambient bulk soil. All the values are means of ±SD

Formatted: Justified

Soil parameters	Elevated (eCO ₂)	Ambient	Ambient bulk soil
pH	7.9±0.1 ^a	8.1±0.3 ^b	8.2±0.4 ^b
EC	0.76±0.03 ^b	0.75±0.02 ^b	0.8±0.01 ^a
Bulkdensity(mg/m ³)	1.49 ±0.1 ^a	1.49±0.1 ^a	1.49±0.2 ^a
SOC (g/kg ⁻¹)	6.9±0.1 ^a	5.8±0.4 ^b	5.7±0.2 ^b

Comment [DRY10]: Correct unit g/kg

3.2. Microbial distribution

The development of bacteria has shown an increasing tendency as CO₂ levels rise (Fig 1). However, insignificant increase in microbial populations of bacteria, fungi, actinomycetes, heterotrophs, and diazotrophs. There was a significant difference observed in methanotrophs, Sulphur oxidizers, methanogens, and phosphorous solubilizing bacteria at ($p<0.01$) and ($p<0.05$) levels of significance among the three soils. Insignificant decrease of SRB under eCO₂ is observed this may be due to the proliferation of methanogens by utilizing CO₂ as a substrate and limiting the substrates required for the sulphidogenesis. The highest microbial populations were seen in soil treated with eCO₂ followed by ambient CO₂ soil, respectively. Variations in the microbial population may have altered the breakdown of organic molecules. Furthermore, bacteria account for around one-quarter of all living biomass on the planet, and their biomass is employed as an early indication of changing physical and chemical features of soils. Increased microbial growth was associated with higher levels of plant-derived C in the rhizosphere under eCO₂. Which is consistent with previous research using other plant species [33–35]. Increase in the eCO₂ increases the soil temperature and thus influences the total the microbial communities of bacteria, diazotrophs, fungi, actinobacteria, methanotrophs, and diazotrophs (Lou et al., 2006). In present soil hike in total anaerobic population under eCO₂ and decrease in methanotrophic population compared bulk soil as shown in fig 1. Similar results were obtained by [36]. Increases in the enzyme activities in soils boost up the carbon pools and diazotrophic population which was observed in our study (Fig 1 & Table 2). Similar results were observed by [37]. Significantly higher microbial ratios under enhanced CO₂ treatments than under ambient CO₂ imply that CO₂ amplification of C inputs may still play a significant role in defining community structure in N-rich agroecosystems, as seen in many forests and grasslands [38]. Organic matter concentration, water availability, and soil physical and chemical qualities all have an impact on microbial biomass.

~~Fig.1. Distribution of different groups of microbes. Bacteria, fungi, actinomycetes, heterotrophs, diazotrophs, methanotrophs, sulfur-oxidizing bacteria, sulfur reducing bacteria, methanogens, and phosphate solubilizing bacteria from three different soils elevated(eCO₂), ambient(A), and ambient bulk soil (ABS).~~

Formatted: Justified, Indent: Left: 0", Hanging: 0.39"




Fig.1. Distribution of different groups of microbes. Bacteria, fungi, actinomycetes, heterotrophs, diazotrophs, methanotrophs, sulphur-oxidizing bacteria, sulphur reducing bacteria, methanogens, and phosphate solubilizing bacteria from three different soils elevated(eCO₂), ambient(A), and ambient bulk soil (ABS).

3.3. Enzyme activities in soil

Soil enzymes profit from biological and metabolic activities in soil, which form the foundation of the terrestrial ecosystem. Elevated atmospheric CO₂ levels clearly have an effect on the soil enzyme activity that is influenced by the physical and chemical properties of soil, soil biological community, and plant either directly or indirectly. All the values are calculated by using a standard graph with an ($R^2 > 0.90$). There were significant changes observed in all enzyme activity in the soils at ($p < 0.01$) and ($p < 0.05$) levels of significance respectively. Where the highest values for all the samples were observed in the soil treated under long-term elevated CO₂ as shown in Table 2. Similar results were also obtained by [39]. The stimulation in enzyme activities by increasing CO₂ demonstrated that increased microbial biomass was accompanied by increased activity [40]. It is recognized that increased CO₂ levels can boost carbon input into the soil by improving the allocation of fixed carbon below ground [41]. This procedure can provide some information on future enzyme activity or the sum of active enzymes. However, because enzyme activities (particularly immobilized enzymes) are affected by edaphic circumstances, actual activity may differ from potential activity [42].

Table 2. Soil enzyme activities from three different soils elevated, ambient, and outside ambient. All the values are means of \pm SD. Significant change in all the enzyme activities at 1% and 5% level of significance.

Soil parameters	SDH ($\mu\text{g TPF g}^{-1}$ soil 24 hr ⁻¹)	BG ($\mu\text{g pNP g}^{-1}$ soil)	FDA ($\mu\text{g fluorescein released g}^{-1}$ soil)	ALP ($\mu\text{g pNP g}^{-1}$ soil)	ACP ($\mu\text{g pNP g}^{-1}$ soil)	UREASE ($\mu\text{g NH}_3 \text{g}^{-1}$ soil)	ARS ($\text{nmol g}^{-1} \text{min}^{-1}$)
eCO ₂	10.02 \pm 0.23 ^a	132.35 \pm 1.29 ^a	4.04 \pm 0.04 ^a	57.69 \pm 0.21 ^a	36.04 \pm 0.12 ^a	48.16 \pm 0.39 ^a	12.20 \pm 0.11 ^a
A	5.38 \pm 0.27 ^c	130.38 \pm 1.54 ^a	3.92 \pm 0.08 ^b	49.01 \pm 0.15 ^b	31.41 \pm 0.19 ^a	41.36 \pm 0.34 ^b	9.55 \pm 0.12 ^b

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Justified

Comment [DRY11]: Foot can be used all full form of short term

ABS	8.22±0.26 ^b	118.92±1.34 ^b	2.96±0.09 ^c	45.93±0.18 ^b	36.42±0.26 ^a	34.85±0.26 ^c	8.94±0.16 ^c
-----	------------------------	--------------------------	------------------------	-------------------------	-------------------------	-------------------------	------------------------

3.4. Sulfur Sulphur fractions

The content of S and sulfur sulphur fractions in the soil increased significantly due to eCO₂ in OTC, the available sulfur sulphur ranging from 13.49 to 17.13 mg kg⁻¹ (Fig 2). The largest available S content was discovered in the 550 ppm CO₂ treated OTC 17.13 mg kg⁻¹ followed by ambient soil 14.48 mg kg⁻¹) and the lowest value was reported in ambient bulk soil 13.49 mg kg⁻¹. Fractions of sulfur sulphur in the soils follow the order (AS≥WSS<ES<PS) Shown in Fig 2. The capacity of soils to adsorb sulfate varies greatly due to the clay composition of soils containing different proportions of hydrous oxides of Al and Fe. However, [43] found that eCO₂ had little effect on soil S content. According to [44], diminished pH enhances sulphate adhesion in soil systems.

Fig 2. Distribution of sulphur fractions available sulphur(AS), water-solublesulphur(WSS), exchangeablesulphur(ES), and precipitated sulphur(PS) from three different soils elevated, ambient, and ambient bulk soil.

Formatted: Justified




Fig 2. Distribution of sulphur fractions available sulphur(AS), water-solublesulphur(WSS), exchangeablesulphur(ES), and precipitated sulphur(PS) from three different soils elevated, ambient, and ambient bulk soil.

3.5. Total Carbon, Nitrogen, Sulfur Sulphur, and Available NPK in soil.

The quantity of CO₂ within the OTC had a detrimental influence on available soil N, P, and K content, which increased as CO₂ concentration increased significantly at 1% and 5 % levels of significance (Table-3). The maximum soil NPK concentration, was found with eCO₂, followed by ambient and ambient bulk soil respectively as shown in Table-3, Plant requirement for nitrogen (N) is predicted to grow as atmospheric CO₂ levels rise (eCO₂). Nitrogen supplementation improves the impact of CO₂ on plant production. Furthermore, eCO₂ greatly improved N absorption [45, 46]. As a result, the amount of the response in plant productivity is primarily determined by how well plant N absorption can keep up with eCO₂-induced stimulation of carbohydrate formation and growth. Plants may positively control a number of physiological processes, such as enzyme production and root development, to boost plant nutrient

acquisition capacity for optimum eCO₂ adaptation [47]. Significant increases in available P requirement by plants are predicted under increased CO₂ levels due to photosynthetic stimulation and subsequent growth responses. Elevated CO₂ affects P acquisition by altering root shape and increasing rooting depth. Furthermore, with high CO₂, the amount and content of root exudates are anticipated to alter due to changes in carbon fluxes through the glycolytic pathway and the tricarboxylic acid cycle. As a result, root exudates may induce P mobilization via alkylation of P from low solubility P complexes, shifts in the biochemical environment, and alters indigenous soil microbial activity.[48].Significant increases in the available potassium in the soil due to lowering of pH in soil due to the formation of carbonic acid and solubility of the ions increases in soil. Increases in the available potassium under elevated CO₂ have been reported by[49].In terms of total carbon nitrogen and sulfursulphur, no significant difference was found among the three different treatments observed.

Table 3 Distribution of carbon, nitrogen, and sulfursulphur in three different soils elevated, ambient, and ambient bulk soil. All the values are means of ±SD. Significant difference found among available N, P, K at (p<0.01) and (p<0.05) level of significance.

Soil parameters	eCO ₂	A	ABS
Available N <u>kg/ha</u> ⁻¹	285.6±2.2 ^a	268.8±3.6 ^a	246.4±3.9 ^b
Available P <u>kg/ha</u> ⁻¹	37.01±0.56 ^a	32.48±0.94 ^b	28.95±0.85 ^c
Available K <u>kg/ha</u> ⁻¹	490±12 ^a	470±14 ^b	454±18 ^c
Total N (%)	0.0784±0.005 ^a	0.0795±0.09 ^a	0.0756±0.06 ^a
Total S(%)	0.42±0.08 ^a	0.41±0.09 ^a	0.41±0.06 ^a
Total C(%)	0.88±0.02 ^a	0.87±0.02 ^a	0.87±0.01 ^a

Comment [DRY12]: full form should be in footnot

Formatted Table

Comment [DRY13]: Unit should be either kg/ha or kg ha-1

Conclusion

Elevated CO₂ levels are to blame for rising global temperatures, which may have a significant influence on soil health, crop output, and plant nutrient usage efficiency. In the current experiment, the findings showed that the responses of soil microbes and their community structure to elevated CO₂ changed significantly in the rice low land ecosystem which is under long-term elevated. S fractionation in rice field is a powerful tool for understanding S dynamics and biogeochemistry in wetland ecosystems. eCO₂ greatly influences the fractionation of sulfursulphur.The study found that all sulfursulphur fractions, including available S, water soluble S, and exchangeable S, precipitated S and total S were significantly influenced. Precipitated sulfursulphurwasdiscovered to be the most dominant among the observed fractions. These changes may be largely related to CO₂-induced changes in soil C, N and S availability.

Comment [DRY14]: Blame is scientific word, please replace with speific term

Ethical approval

Not applicable.

Consent for Publication

All authors have reviewed the manuscript and agree to its publication.

Reference

1. Wani SH, Sah SK (2014) Biotechnology and abiotic stress tolerance in rice. J Rice Res 2:e105
2. Lavakush, Yadav J, Verma JP, et al (2014) Evaluation of PGPR and different concentration of

Comment [DRY15]: Follow journal common format in all reference

phosphorus level on plant growth, yield and nutrient content of rice (*Oryza sativa*). *Ecol Eng* 62:123–128. <https://doi.org/https://doi.org/10.1016/j.ecoleng.2013.10.013>

3. Singh KM, Singh D (2020) Challenges of ensuring food and nutritional security in Bihar. *Challenges Ensuring Food Nutr Secur Bihar* (January 10, 2020)
4. Alam MA, Rahman MM, Khan MAR, et al (2021) Modulation of Yield and Yield Attributes of Transplanted Aman Rice through Organic and Inorganic Fertilization in Terrace Soil. *Eur J Agric Food Sci* 3:97–105
5. Alam MA, Rahman MM, Biswas JC, et al (2019) Nitrogen transformation and carbon sequestration in wetland paddy field of Bangladesh. *Paddy Water Environ* 17:677–688. <https://doi.org/10.1007/s10333-019-00693-7>
6. Rahman MM (2013) Nutrient-use and carbon-sequestration efficiencies in soils from different organic wastes in rice and tomato cultivation. *Commun Soil Sci Plant Anal* 44:1457–1471
7. Rastogi M, Singh S, Pathak H (2002) Emission of carbon dioxide from soil. *Curr Sci* 82:510–517
8. Kahn H (2009) *On Escalation: Metaphors and Scenarios* (Vol. 1)
9. Mackay A (2008) Climate change 2007: impacts, adaptation and vulnerability. Contribution of Working Group II to the fourth assessment report of the Intergovernmental Panel on Climate Change. *J Environ Qual* 37:2407
10. Risbey JS (2008) The new climate discourse: Alarmist or alarming? *Glob Environ Chang* 18:26–37
11. Razzaque MA, Haque MM, Khaliq QA, Solaiman ARM (2009) The effect of different nitrogen levels and enrichment CO₂ on the nutrient contents of rice cultivars. *Bangladesh J Sci Ind Res* 44:241–246
12. Singh AK, Amgain LP, Sharma SK (2000) Root characteristics, soil physical properties and yield of rice (*Oryza sativa*) as influenced by integrated nutrient management in rice-wheat (*Triticum aestivum*) system. *Indian J Agron* 45:217–222
13. Tiwari KN, Gupta BR (2006) Sulphur for sustainable high yield agriculture in Uttar Pradesh. *Indian J Fertil* 1:37
14. Sharma U, Subehia SK, Rana SS, et al (2014) Soil sulphur fractions and their relationship with soil properties and rice (*Oryza sativa* L.) yield under long-term integrated nutrient management in an acid Alfisol. *Res Crop* 15:
15. Vidyalakshmi R, Sridar R (2007) Isolation and characterization of sulphur oxidizing bacteria. *J Cult Collect* 5:73–77
16. Hungate RE (1969) Chapter IV A roll tube method for cultivation of strict anaerobes. In: *Methods in microbiology*. Elsevier, pp 117–132
17. Postgate JR (1963) Versatile medium for the enumeration of sulfate-reducing bacteria. *Appl Microbiol* 11:265–267
18. Döbereiner J (1989) Isolation and identification of root associated diazotrophs. In: *Nitrogen fixation with non-legumes*. Springer, pp 103–108
19. Casida Jr LE, Klein DA, Santoro T (1964) Soil dehydrogenase activity. *Soil Sci* 98:371–376
20. Tabatabai MA, Bremner JM (1969) Use of p-nitrophenyl phosphate for assay of soil phosphatase activity. *Soil Biol Biochem* 1:301–307
21. Eivazi F, Tabatabai MA (1990) Factors affecting glucosidase and galactosidase activities in soils. *Soil Biol Biochem* 22:891–897
22. Tabatabai MA, Bremner JM (1970) Arylsulfatase activity of soils. *Soil Sci Soc Am J* 34:225–229
23. Schnierer J, Rosswall T (1982) Fluorescein diacetate hydrolysis as a measure of total microbial activity in soil and litter. *Appl Environ Microbiol* 43:1256–1261

24. Morche L (2008) S-fluxes and spatial alterations of inorganic and organic sulphur fractions in soil as well as their accumulation and depletion in the rhizosphere of agricultural crops by partial use of the radioisotope ³⁵S. Bonn Rheinischen Friedrich-Wilhelms-Universität Bonn (in Ger)
25. Carrillo Y, Dijkstra FA, Pendall E, et al (2014) Plant rhizosphere influence on microbial C metabolism: the role of elevated CO₂, N availability and root stoichiometry. *Biogeochemistry* 117:229–240
26. Celia MA, Peters CA, Bachu S (2002) Geologic storage of CO₂: leakage pathways and environmental risks. In: AGU Spring Meeting Abstracts. pp GC32A-03
27. Cardon ZG (1995) Influence of rhizodeposition under elevated CO₂ on plant nutrition and soil organic matter. *Plant Soil* 187:277–288
28. Lindsay WL (1979) Chemical equilibria in soils. John Wiley and Sons Ltd.
29. Patil RH, Colls JJ, Steven MD (2010) Effects of CO₂ gas as leaks from geological storage sites on agro-ecosystems. *Energy* 35:4587–4591
30. Canadell JG, Pitelka LF, Ingram JSI (1995) The effects of elevated [CO₂] on plant-soil carbon below-ground: a summary and synthesis. *Plant Soil* 187:391–400
31. Satapathy SS, Swain DK, Pasupalak S, Bhadoria PBS (2015) Effect of elevated [CO₂] and nutrient management on wet and dry season rice production in subtropical India. *Crop J* 3:468–480
32. Six J, Carpentier A, van Kessel C, et al (2001) Impact of elevated CO₂ on soil organic matter dynamics as related to changes in aggregate turnover and residue quality. *Plant Soil* 234:27–36
33. Drigo B, Kowalchuk GA, Knapp BA, et al (2013) Impacts of 3 years of elevated atmospheric CO₂ on rhizosphere carbon flow and microbial community dynamics. *Glob Chang Biol* 19:621–636
34. Da Costa DP, Dias ACF, Cotta SR, et al (2018) Changes of bacterial communities in the rhizosphere of sugarcane under elevated concentration of atmospheric CO₂. *GCB Bioenergy* 10:137–145
35. Wang YH, Yu ZH, Li YS, et al (2019) Elevated CO₂ alters the structure of the bacterial community assimilating plant-derived carbon in the rhizosphere of soya bean. *Eur J Soil Sci* 70:1212–1220
36. Liu D, Tago K, Hayatsu M, et al (2016) Effect of elevated CO₂ concentration, elevated temperature and no nitrogen fertilization on methanogenic archaeal and methane-oxidizing bacterial community structures in paddy soil. *Microbes Environ* 31:349–356
37. Kumar U, Nayak AK, Sahoo S, et al (2020) Combined effects of elevated CO₂, N fertilizer and water deficit stress on diazotrophic community in sub-humid tropical paddy soil. *Appl Soil Ecol* 155:103682
38. Feng X, Simpson AJ, Schlesinger WH, Simpson MJ (2010) Altered microbial community structure and organic matter composition under elevated CO₂ and N fertilization in the duke forest. *Glob Chang Biol* 16:2104–2116
39. Dhillon SS, Roy J, Abrams M (1995) Assessing the impact of elevated CO₂ on soil microbial activity in a Mediterranean model ecosystem. *Plant Soil* 187:333–342
40. Ebersberger D, Niklaus PA, Kandeler E (2003) Long term CO₂ enrichment stimulates N-mineralisation and enzyme activities in calcareous grassland. *Soil Biol Biochem* 35:965–972
41. Jones TH, Thompson LJ, Lawton JH, et al (1998) Impacts of rising atmospheric carbon dioxide on model terrestrial ecosystems. *Science* (80-) 280:441–443
42. Freeman C, Liska G, Ostle NJ, et al (1995) The use of fluorogenic substrates for measuring enzyme activity in peatlands. *Plant Soil* 175:147–152
43. Fangmeier A, De Temmerman L, Black C, et al (2002) Effects of elevated CO₂ and/or ozone on nutrient concentrations and nutrient uptake of potatoes. *Eur J Agron* 17:353–368

- |
44. Shainberg I, Sumner ME, Miller WP, et al (1989) Use of gypsum on soils: A review. *Adv soil Sci* 1–111
 45. Ainsworth EA, Long SP (2005) What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytol* 165:351–372
 46. Langley JA, Megonigal JP (2010) Ecosystem response to elevated CO₂ levels limited by nitrogen-induced plant species shift. *Nature* 466:96–99
 47. Li Y, Yu Z, Liu X, et al (2017) Elevated CO₂ increases nitrogen fixation at the reproductive phase contributing to various yield responses of soybean cultivars. *Front Plant Sci* 8:1546
 48. Jin J, Tang C, Sale P (2015) The impact of elevated carbon dioxide on the phosphorus nutrition of plants: a review. *Ann Bot* 116:987–999
 49. Reddy KR, Zhao D (2005) Interactive effects of elevated CO₂ and potassium deficiency on photosynthesis, growth, and biomass partitioning of cotton. *F Crop Res* 94:201–213

UNDER PEER REVIEW