

Review Article

Studies on Phosphorous dynamics in soils: A Review

Abstract

More than two-thirds of the world's soils are afflicted with phosphorus shortage (P). Agricultural output suffers because this phosphorus is unavailable to all but a tiny percentage of plants. To lessen the severity of the environmental impact, phosphorus-deficient soils have traditionally been fertilized. Improving agricultural management practices for sustainable crop output, including lowering phosphorus runoff, is the emphasis of this study. Investigating chemical fractionation mechanisms to distinguish between occluded P, acid-extractable calcium-bonded P, and non-occluded P is necessary for increasing Pi in mung bean (*Vigna radiata*) types and establishing net grazing systems. Phosphate (P) is known to be transferred between pools due to weathering, with the highest P retention rates found in clay-rich soils. Soil with a finer texture is better able to absorb and fix phosphorus, which means more nutrients and water can be made available to the plant when mung bean varieties are inoculated into it. More photosynthesis means more accumulated dry stuff. The P-treated variety of mung bean had the greatest yield index (13.28). Pods per plant (46.02), pods per crop (8.20), test weight (40.63 gm), pod weight (8.0 g), and seed weight (1.0 g) were all significantly increased with DAP (1.83 g). An increase in nodule count, leaf area index, plant height, grain yield, total chlorophyll content, and straw output of up to 40% was observed at higher P₂O₅ concentrations (2988 kilogrammes per hectare). Each plant developed the maximum number of nodules after receiving injections of PSB and *Aspergillus awamori*. The highest levels of chlorophyll were found in its leaves, and its plants were the largest and most productive overall. Liming improved mungbean yields in acid piedmont soil by raising the pH and altering other chemical characteristics. They claim that increasing India's output of pulses is necessary for the country to attain food security in the future. If farmers and extension workers want to see higher quality pulse yields and greater long-term profitability, they need to incorporate P nutrition into their balanced nutrient management programmes.

Keywords: Phosphatic Fertilizer, grain yield, photosynthetic yield; plant productivity

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Comment [DC1]: alphabetical order

Introduction

Phosphorus (P) is a macronutrient required by plants for its growth. It accounts for between 0.2 and 0.8 percent of the dry weight of a plant. Phosphorus is necessary for a wide variety of physiological and biochemical processes in plants. Phospholipids, nucleic acids, coenzymes, enzymes, and nucleotides include phosphorus (Mehrvaz *et al.*, 2008). In the early stages of plant growth, enough P availability is also necessary for the laying down of the plant's reproductive parts' primordia (Satyaprakash *et al.*, 2017; Kalayu, 2019). The typical amount of phosphorus found in soil is 0.05 percent (w/w). However, only 0.1 percent of this phosphorus is available for plant uptake (Zhu *et al.*, 2011). Phosphorus fertilizers have traditionally been used as a method for addressing phosphorus deficiency in the soil. To maintain consistent amounts of phosphorus in soil and plant systems, fertilizers in 52,3 billion tonnes are sprayed yearly (FAO, 2017). This tremendous quantity of phosphorus, which has been transformed into inorganic phosphorus and precipitated by metal cations in the soil such as iron, aluminium, magnesium, and calcium, is only used by plants at a rate of around 0.2 percent, or 10 micrograms (Soumare *et al.*, 2020). The phosphate deposit in agricultural soils is sufficient to ensure that optimum crop production may be maintained worldwide for about 100 years (Kalay, 2019). In addition, the use of fertilizers contributes to a variety of environmental problems, including the contamination of groundwater and the eutrophication of streams (Yu, *et al.*, 2011 and Alori *et al.*, 2017). Agricultural management strategies need to be developed to improve the efficiency of phosphorus fertilization, increase crop productivity, and reduce environmental pollution caused by phosphorus loss from the soil. These goals will be achieved by reducing the amount of phosphorus loss from the soil.

Forms of Phosphorous in soil system

It is common for soils to have greater concentrations of total P than other key nutrients, such as N or K. More than 80% of P, on the other hand, is inert and so inaccessible to plants (Xu *et al.*, 2020). Figure 1 shows that inorganic and organic forms of P (Pi/Po) are found in soil, and the quantities of each alter over time. Most soil P comes from biological tissues where phosphorus is found in organic substances such as nucleotides, phospholipids, phosphoproteins, and coenzymes (Cross *et al.*, 1995). It is also possible that soil nutrient cycle mechanisms (e.g., nitrogen cycling) are responsible for the dispersion of primary Pi into Po forms. Because of the widespread usage of Po-containing products, including fire

Formatted: Highlight

retardants, plasticizers, insecticides, and antifoam agents, new Po sources have emerged in the environment, increasing soil Po concentrations and diversity (Adams *et al.*, 1992).

Due to its weak connections with the soil particles, Po is more quickly leached than the more stable forms of primary (variscite, apatite, and strengite) and secondary minerals (calcium, aluminium phosphates, and iron) found in soil (Gebrim *et al.*, 2010). Chemical fractionation processes were created in the early studies to determine the three forms of Pi: occluded P (containing 2nd NaOH extractable P and residual Pi), acid-extractable calcium-bonded P and non-occluded P. Fe and Al oxide coatings and concretions absorb occluded P during diffusive penetration and the development of the soil. Occluded P is more difficult to remove from Fe, and Al oxide surfaces than non-occluded P ions that make up non-occluded P. Soil Pi content decreases from 684 g/g in Entisols to 200 to 430 in Ultisol and Oxisols as the soils develop. Plants or soil bacteria may take up Pi and use it in the secondary Po cycle, or it can seep into streams and deposit phosphorus (P) in ocean sediments (Maltais-Landry *et al.*, 2014). A variety of types and amounts of Pi may be found in the soil. Plant structures (such stems, roots, and leaves), dead soil animals, and microbes gradually return the P they absorbed to the soil when mineralization has occurred. It is important to note that even though P has a wide range of biochemical activities in soil that are crucial to the health of the ecosystem, its cycles are slower than those of C, N, and S. A wide range of geochemical processes (weathering, precipitation/dissolution, and solid-phase transitions) have a long-term impact on soil phosphorus availability and distribution. Soil microorganisms immobilize and mineralize plant-available phosphorus (P) in soil organic matter over a short time (10–2 to 100 years). However, very little is known about the function these biological activities play in limiting soil phosphorus availability due to geochemical processes (Hesterberg *et al.*, 1999; Prietzel *et al.*, 2016 and Hao *et al.*, 2020,)




Figure 1: The Phosphorous cycle

Comment [DC2]: source

Release of inorganic P in soil from fertilizer application

Despite the fact that the soil contains a large amount of total P, only a little amount (less than 1%) gets dissolved in it (Bunemann *et al.*, 2015). There are many different forms of phosphorus, including organic P which includes nucleic acids, phospholipids, and orthophosphate), Al/Fe-bound inorganic P (which is found in acidic soils), and Ca-bound inorganic P (which is found in alkaline soils) (in alkaline soil). Occluding-P and Fe/Al-P percentages rise with rising P concentrations in the soil (Bai *et al.*, 2017). Progressively, these compounds release soluble P (0.1 percent of the total soil P), which is slowly released over time. Soluble P, labile soil P, and nonlabile soil P are all various types of P in the soil. For plants to flourish they must have both primary and secondary orthophosphate ($H_2PO_4^-$) and HPO_4^{2-} forms of P (Zhu *et al.*, 2018). Acidic soils contain primary orthophosphates, whereas alkaline soils have secondary orthophosphates (Rashid *et al.*, 2005). Stunted growth, delayed maturity of crops, and poor seed development are all symptoms of a phosphorus deficit in plants. A shortage of total phosphorus in the soil, along with the fixation of extra phosphorus via the use of chemical fertilizers and organic sources such as manure, results in a deficiency of phosphorus in the soil. The overall amount of phosphorus in soil may vary depending on several factors, including pH, organic matter content, particle size distribution, and mineralogy of the parent material (Havlin *et al.*, 2005). The most important controls on P

dynamics in soil are the sorption and desorption capacities of soil P, particle size distribution, metal oxides, and soil pH. (Li *et al.*, 2016). It has been shown that weathering is a factor in the movement of P across various pools. The largest phosphate retention capacity is found in clay-rich soils, which subsequently adsorbs/fixes more P than coarse-textured soils (Brady *et al.*, 2008).

Phosphorus fertilizer is very necessary to take full use of the greatest production potentials offered by various crop plants. Since the ratio of phosphorus that is applied determines the amount of phosphorus that is fixed, the amount of phosphorus that is fixed by fertilizer that is broadcast is significantly higher than the amount of phosphorus that is fixed by fertilizer that is applied in bands due to the narrow soil to fertilizer ratio in the latter situation (Rashid *et al.*, 1993). One of the primary reasons for the low output in crops is the poor performance of phosphatic fertilizer. Fertiligation resulted in a more significant reaction than broadcast and banding. Mungbean fertigation yielded the maximum mungbean production, as well as P absorption, P recovery, and agronomic efficiency, compared to broadcasting methods. Phosphate fertilizer costs have risen worldwide due to the rising demand for greater yields, which has placed a strain on land resources (Weber *et al.*, 2014; Chowdhury *et al.*, 2017). Soil P has been replenished by using phosphate fertilizers. Rock phosphate mining is the primary source for making phosphate fertilizers. Chemical methods must dissolve the phosphate that can only be obtained from mined rock phosphate (Khan *et al.*, 2009). Because crops have a high capacity to fix phosphate and a low ability to utilize it, the rock phosphate is applied to the soil each cropping season (Withers *et al.*, 2001). Phosphate fertilizers that have been applied indiscriminately have created a fixed P pool that may be explored and used by various bacteria, enzymes, organic acids, manures, and humic substances. Organic acids and enzymatic activity released by phosphate solubilizing microorganisms aid plant absorption (Dash *et al.*, 2017). However, soil type, parent material, organic matter concentration, soil pH, soil salinity, and metal oxides all influence phosphate dynamics in the soil.

3. Dynamics of Phosphorous in the soil affected by organic sources

Various phosphorus pools in the soil, such as Al/Fe₂ bound P or Ca-bound P, replenish the P that plants take up from the soil solution (Saleque *et al.*, 2004). With mass flow taking over from diffusion, the function of P in the process is diminished (Hinsinger, 2001). The

availability of Fe, Al, and Ca, which aid in the fixation of P, is also influenced by soil pH. Sodium phosphate is reduced in alkaline soils by forming several insoluble phosphate derivatives. Phosphorus is depleted from acidic soils by precipitates of hydroxyl phosphates generated by the interaction of iron and aluminium with phosphate ions (Harris *et al.*, 2006). Cation and anion concentrations at the exchange sites influence soil adsorption capacity for phosphate, although divalent cations (Ca^{2+}) are significantly more attracted to P adsorption in clays than monovalent cations (Na^+) are in soils.

Formatted: Highlight

Formatted: Highlight

Alkaline calcareous soils have a high concentration of calcium phosphate minerals (CaP) that plants may take up, although most of the P is fixed as CaP. In addition to supplying P, clay minerals in soil play a significant role (Mehdi *et al.*, 2007). Both the fixation and transformation of phosphorus are intricate processes, and it may be challenging to pin down the appropriate kinetics for them. Phosphate may be in the soil in several different forms, including solution P, active P, as well as fixed P.




Figure 2: Various types of total Phosphorous in the soil matrix.

Comment [DC3]: source

Phospholipids, inositol phosphates, and nucleic acids are the three different kinds of organic P compounds. Phospholipids are the most common kind. Inositol phosphates include mono-

to hexaphosphate esters as their primary building blocks (Dash *et al.*, 2017). One of the most common organic phosphates (Ca–Mg salts of phytic acid) is also phyton. There are three kinds of enzymes that can break down organic P, which include phosphates and phytases as well as phosphatases (Ingle and Padole, 2017).

As a result of using goat dung in combination with low-rates of P fertilizers, the amount of the less-labile NaOH-Pi fraction in the soil may be reduced, ensuring that the soil's labile P levels are maintained at an adequate level. P needs for Smallholder farmers may be met with this cost-effective technique for enhancing fertilizer P usage efficiency (Gichangi *et al.*, 2009).

After two and 16 weeks of composting in labile pools, it was shown that the Pi percentage soil was more likely to be dominated by compost P. However, the amount of labile P (51-58 percent), as well as total Pi (61-98 percent) in soil, was the same or nearly identical to that obtained from inorganic KH_2PO_4 alone. Compost from potato trash had a higher proportion of organic P (Po) than those from the other composts investigated when Po was extracted using NaHCO_3 . While this study indicated that Pi was the most necessary form of compost P, it is still hard to assess the availability of compost P after it is applied to soil (Gagnon *et al.*, 2012).

Additionally, it was shown that the soil formed from Basalt might lose some of its ability to bind P when organic molecules were introduced through the treatments. The intensity of P adsorption decreased when chicken dung as compost was added to soils originating from granite and river alluvial deposits but rose following the application of organic fertilizer. The three adsorption isotherm equations for basalt-derived soils all fit well. Langmuir >Freundlich >Temkin adsorption models were used for soils generated from granite and alluvial deposits as parent materials. It was shown that organic fertilizer and chicken manure had no effect on soils generated from basalt's MPBC or SPR, however organic fertilizer and incubation time had a significant impact on these parameters in soils formed from granite and alluvial deposits (Yu *et al.*, 2013).

As in both soils, a sandy loam acidic soil (Aquepts) and a silt loam alkaline soil, extraction efficiency was in the sequence of Bray and Kurtz-II>Mehlich-3> Kelowna>Olsen amended with diverse organic sources, such as CM, PM, CW, and triple superphosphate

Comment [DC4]: meaning

(TSP). As evaluated by NaHCO_3 , organic additions improve the effectiveness of the additional P, but the efficiency of Triple superphosphate with days of incubation decreases (Ara *et al.*, 2018).

Nitrogen (N), as well as Leaf litter, had an effect on the bacterial action and it led to an increase in phosphorus orthophosphate (Po), which resulted in a rise in the NaOH-Pi fractions, as a result, the quantity of Po detected in the NaOH-Pi portion increased overall (Kunito *et al.*, 2018). To increase P availability, organic matter may reduce P-selectivity and the maximum phosphate buffering capacity. At a SOM concentration of 75.3 grams per kilogram (g/kg), there is a growing tendency toward greater availability.

Formatted: Highlight

The relationship between different sources and sinks of phosphorus by using an intense cropping system that included rice, maize, and green gram (residual) was determined. Inorganic P fractions were for 71.9–86.0 percent of total P in surface soil, whereas organic P fractions accounted for 14.0–28.1 percent. The percentages in the subsoil ranged from 75.9–81 percent and 18.1–24.1 percent. The following is the order of the pools: Up to the first 15 centimetres of the soil, the order is $\text{Fe-P} > \text{RS-P} > \text{Al-P} > \text{Ca-P} > \text{Occluded P} > \text{Saloid P}$; however, the lowest 15–30 centimetres of soil have the following order: "Occluded P" = "Al-P" > "saloid P" in the RS-P. Both fractions provide a large quantity of phosphorus to the surface layer of the soil. In terms of inorganic phosphorus forms, there is a positive association between surface soil P absorption and Fe-P as well as a significant relation between the subsurface soil P uptake and Al-P (Chauhan *et al.*, 2012).

Conclusion

Phosphorus plays an important role in pulses nutrition mainly for mungbean production. Most of the Indian soils are deficient in phosphorus leading to affect the nutrition of pulse crop as pulses occupy an important position in food and nutritional security in India. India produces over 280 million tonnes of foodgrains every year with an increase of four folds since independence. Increased efforts to produce more food have resulted in tremendous shift in cropping systems towards cereal-cereal based systems. Accordingly, vision of Indian Institute of Pulse Research, 2030, the projected pulse requirement by the year 2030 would be 32 million tons with an anticipated required growth rate of 4.2%. The growing Indian population enhanced the pulses demand. The burgeoning human population in India needs

Formatted: Highlight

Formatted: Highlight

higher pulses production for fulfilling the dietary protein requirement. It requires to mitigating the demand of protein. In this regard, balanced fertilization with NPK along with biofertilizers has been proved beneficial in pulses both under rainfed and irrigated conditions. Hence, phosphorous needs to be taken care of while balanced fertilization in pulses. A concerted effort by farmers, researchers, development agencies, and government are needed to ensure that India becomes self-sufficient in pulses in the future. Therefore, farmers and extension functionaries must recognize that it is highly needed to include P nutrition in their nutrient management practices in balanced manner in present day intensive farming to harvest higher pulse yields of superior quality earning maximum sustainable profitability.

Reference:

- Adams, M. A., & Pate, J. S. (1992). Availability of organic and inorganic forms of phosphorus to lupins (*Lupinus* spp.). *Plant and Soil*, 145(1), 107-113.
- Alori, E. T., Glick, B. R., & Babalola, O. O. (2017). Microbial phosphorus solubilization and its potential for use in sustainable agriculture. *Frontiers in microbiology*, 8, 971.
- Amsalu, S., & Beyene, S. (2020). Effects of lime and phosphorous application on chemical properties of soil, dry matter yield, and phosphorus concentration of barley (*Hordeum vulgare*) grown on Nitosols of Emdibir, Southern Ethiopia. *Journal of Soil Science and Environmental Management*, 11(4), 131-141.
- Ara, I., Islam, M., Kashem, M., & Osman, K. T. (2018). A comparative study of phosphorus availability in an acidic soil and an alkaline soil amended with organic and inorganic phosphorus sources. *Journal of soil science and plant nutrition*, 18(2), 466-478.
- Bai, J., Ye, X., Jia, J., Zhang, G., Zhao, Q., Cui, B., & Liu, X. (2017). Phosphorus sorption-desorption and effects of temperature, pH and salinity on phosphorus sorption in marsh soils from coastal wetlands with different flooding conditions. *Chemosphere*, 188, 677-688.
- Batjes, N. H., & Sombroek, W. G. (1997). Possibilities for carbon sequestration in tropical and subtropical soils. *Global change biology*, 3(2), 161-173.
- Brady NC, Weil RR. 2008. *The Nature and Properties of Soils*, 14th ed., Upper Saddle River, NJ: Prentice Hall.

- Branch, S., & Maria, S. (2017). Evaluation of the functional properties of mung bean protein isolate for development of textured vegetable protein. *International Food Research Journal*, 24(4), 1595-1605.
- Bünemann, E. K. (2015). Assessment of gross and net mineralization rates of soil organic phosphorus—A review. *Soil Biology and Biochemistry*, 89, 82-98.
- Chauhan, B. S., Mahajan, G., Sardana, V., Timsina, J., & Jat, M. L. (2012). Productivity and sustainability of the rice–wheat cropping system in the Indo-Gangetic Plains of the Indian subcontinent: problems, opportunities, and strategies. *Advances in agronomy*, 117, 315-369.
- Chen, Y. S., Chen, S. C., Kao, C. M., & Chen, Y. L. (2003). Effects of soil pH, temperature and water content on the growth of *Burkholderia pseudomallei*. *Folia microbiologica*, 48(2), 253-256.
- Chowdhury RB, Moore GA, Weatherley AJ, Arora M. 2017. Key sustainability challenges for the global phosphorus resource, their implications for global food security, and options for mitigation. *J Clean Prod* 140:945–963.
- Cross, A. F., & Schlesinger, W. H. (1995). A literature review and evaluation of the Hedley fractionation: Applications to the biogeochemical cycle of soil phosphorus in natural ecosystems. *Geoderma*, 64(3-4), 197-214.
- Dash, N., Pahari, A., & Dangar, T. K. (2017). Functionalities of phosphate-solubilizing bacteria of rice rhizosphere: techniques and perspectives. In *Recent advances in Applied Microbiology* (pp. 151-163). Springer, Singapore.
- Gagnon, B., Demers, I., Ziadi, N., Chantigny, M. H., Parent, L. E., Forge, T. A., ... & Buckley, K. E. (2012). Forms of phosphorus in composts and in compost-amended soils following incubation. *Canadian journal of soil science*, 92(5), 711-721.
- FAO, IFAD, UNICEF, WFP, and WHO. 2017. Building resilience for peace and food security. The state of food security and nutrition in the world. Rome: Food and Agricultural Organization of the United Nations. Accessed June 25, 2020. <http://www.fao.org/3/a-I7695e.pdf>
- Gebrim, F. D. O., Novais, R. F., Silva, I. R. D., Schulthais, F., Vergütz, L., Procópio, L. C., ... & Jesus, G. L. D. (2010). Mobility of inorganic and organic phosphorus forms under

- different levels of phosphate and poultry litter fertilization in soils. *Revista Brasileira de Ciência do Solo*, 34(4), 1195-1205.
- Gichangi, E. M., Mnkeni, P. N., & Brookes, P. C. (2009). Effects of goat manure and inorganic phosphate addition on soil inorganic and microbial biomass phosphorus fractions under laboratory incubation conditions. *Soil Science and Plant Nutrition*, 55(6), 764-771.
- Haynes, R. J., & Ludecke, T. E. (1981). Yield, root morphology and chemical composition of two pasture legumes as affected by lime and phosphorus applications to an acid soil. *Plant and soil*, 62(2), 241-254.
- Halim, A., Siddique, M. N. E. A., Sarker, B. C., Islam, M. J., Hossain, M. F., & Kamaruzzaman, M. (2014). Assessment of nutrient dynamics affected by different levels of lime in a mungbean field of the Old Himalayan Piedmont soil in Bangladesh. *J. Agric. Vet. Sci*, 7, 101-112.
- Hao, J., Knoll, A. H., Huang, F., Schieber, J., Hazen, R. M., & Daniel, I. (2020). Cycling phosphorus on the Archean Earth: Part II. Phosphorus limitation on primary production in Archean ecosystems. *Geochimica et Cosmochimica Acta*, 280, 360-377.
- Harris, J. N., New, P. B., & Martin, P. M. (2006). Laboratory tests can predict beneficial effects of phosphate-solubilising bacteria on plants. *Soil Biology and Biochemistry*, 38(7), 1521-1526.
- Havlin JL, Beaton JD, Tisdale SL, Nelson WL. 2005. Soil Fertility and Fertilizers: An Introduction to Nutrient Management, 7th ed. Upper Saddle River, NJ: Prentice-Hall
- Hesterberg, D., Zhou, W., Hutchison, K. J., Beauchemin, S., & Sayers, D. E. (1999). XAFS study of adsorbed and mineral forms of phosphate. *Journal of Synchrotron Radiation*, 6(3), 636-638.
- Hinsinger, P. (2001). Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review. *Plant and soil*, 237(2), 173-195.
- Hossain, M. A., & Hamid, A. (2007). Influence of N and P fertilizer application on root growth, leaf photosynthesis and yield performance of groundnut. *Bangladesh Journal of Agricultural Research*, 32(3), 369-374.
- Ingle, K. P., & Padole, D. A. (2017). Phosphate solubilizing microbes: An overview. *Int J Curr Microbiol Appl Sci*, 6(1), 844-852.

- Kalayu, G. (2019). Phosphate solubilizing microorganisms: promising approach as biofertilizers. *International Journal of Agronomy*, 2019.
- Khan, A. A., Jilani, G., Akhtar, M. S., Naqvi, S. M. S., & Rasheed, M. (2009). Phosphorus solubilizing bacteria: occurrence, mechanisms and their role in crop production. *J agric biol sci*, 1(1), 48-58.
- Kumar, S., Wani, J. A., Panotra, N., Lone, B. A., Qayoom, S., & Fayaz, A. (2017). Effect of phosphorus and sulphur on nutrient and amino acids content of soybean [Glycine max (L.) Merrill] under 'Eutrochrepts'. *Legume Res*, 40, 716-720.
- Kumawat, N., Kumar, R., & Sharma, O. P. (2009). Nutrient uptake and yield of mungbean [Vigna radiata (L.) Wilczek] as influenced by organic manures, PSB and phosphorus fertilization. *Environ Ecol*, 27(4B), 2002-2005.
- Kumawat, P. K., Tiwari, R. C., Golada, S. L., Godara, A. S., Garhwal, R. S., & Choudhary, R. (2013). Effect of phosphorus sources, levels and biofertilizers on yield attributes, yield and economics of black gram (Phaseolus mungo L.). *Legume Research: An International Journal*, 36(1).
- Kunito, T., Hiruta, N., Miyagishi, Y., Sumi, H., & Moro, H. (2018). Changes in phosphorus fractions caused by increased microbial activity in forest soil in a short-term incubation study. *Chemical Speciation & Bioavailability*, 30(1), 9-13.
- Li, W., Zheng, Y., Zhang, H., Liu, Z., Su, W., Chen, S., ... & Lei, B. (2016). Phytotoxicity, uptake, and translocation of fluorescent carbon dots in mung bean plants. *ACS applied materials & interfaces*, 8(31), 19939-19945.
- Mahmud, K., Makaju, S., Ibrahim, R., & Missaoui, A. (2020). Current progress in nitrogen fixing plants and microbiome research. *Plants*, 9(1), 97.
- Maltais-Landry, G., Scow, K., & Brennan, E. (2014). Soil phosphorus mobilization in the rhizosphere of cover crops has little effect on phosphorus cycling in California agricultural soils. *Soil Biology and Biochemistry*, 78, 255-262.
- Margenot, A. J., Singh, B. R., Rao, I. M., & Sommer, R. (2016). Phosphorus fertilization and management in soils of Sub-Saharan Africa. *Soil phosphorus*, 151-208.
- Meena, R. S., Dhakal, Y., Bohra, J. S., Singh, S. P., Singh, M. K., Sanodiya, P., & Meena, H. (2015). Influence of bioinorganic combinations on yield, quality and economics of Mungbean. *Am J Exp Agric*, 8(3), 159-166.

- Mehdi, S. M., Obaid-ur-Rehman, A., Ranjha, M., & Sarfaraz, M. (2007). Adsorption capacities and availability of phosphorus in soil solution for rice wheat cropping system. *World Appl Sci J*, 2(4), 244-265.
- Mondal, A. K., & Mondal, S. (2011). Circumscription of the families within Leguminales as determined by cladistic analysis based on seed protein. *African Journal of Biotechnology*, 10(15), 2850-2856.
- Mehrvarz, S., Chaichi, M. R., & Alikhani, H. A. (2008). Effects of phosphate solubilizing microorganisms and phosphorus chemical fertilizer on yield and yield components of barley (*Hordeum vulgare* L.). *J. Agric. Environ. Sci*, 3(6), 822-828.
- Penn, C. J., & Camberato, J. J. (2019). A critical review on soil chemical processes that control how soil pH affects phosphorus availability to plants. *Agriculture*, 9(6), 120.
- Prietzl, J., Harrington, G., Häusler, W., Heister, K., Werner, F., & Klysubun, W. (2016). Reference spectra of important adsorbed organic and inorganic phosphate binding forms for soil P speciation using synchrotron-based K-edge XANES spectroscopy. *Journal of synchrotron radiation*, 23(2), 532-544.
- Rashid, A. (2005). Effect of intercropping on the yield and net monetary return of sorghum production. *Sarhad Journal of Agriculture (Pakistan)*.
- Rashid, A., & Din, J. (1993). Phosphorus deficiency diagnosis and fertilizer placement for mungbean grown in alkaline soils. *Journal of Agronomy and Crop Sciences*, 12, 135-172.
- Rasool, B., Ramzani, P. M. A., Zubair, M., Khan, M. A., Lewińska, K., Turan, V., ... & Iqbal, M. (2021). Impacts of oxalic acid-activated phosphate rock and root-induced changes on Pb bioavailability in the rhizosphere and its distribution in mung bean plant. *Environmental Pollution*, 280, 116903.
- Saleque, M. A., Naher, U. A., Islam, A., Pathan, A. B. M. B. U., Hossain, A. T. M. S., & Meisner, C. A. (2004). Inorganic and organic phosphorus fertilizer effects on the phosphorus fractionation in wetland rice soils. *Soil Science Society of America Journal*, 68(5), 1635-1644.
- Satyaprakash, M., Nikitha, T., Reddi, E. U. B., Sadhana, B., & Vani, S. S. (2017). Phosphorous and phosphate solubilising bacteria and their role in plant nutrition. *International Journal of Current Microbiology and Applied Sciences*, 6(4), 2133-2144.

- Singh, A., Baoule, A. L., Ahmed, H. G., Dikko, A. U., Aliyu, U., Sokoto, M. B., ... & Haliru, B. (2011). Influence of phosphorus on the performance of cowpea (*Vigna unguiculata* (L) Walp.) varieties in the Sudan savanna of Nigeria. *Agricultural Sciences*, 2(03), 313.
- Singh, R., Singh, P., Singh, V., & Yadav, R. A. (2018). Effect of phosphorus and PSB on yield attributes, quality and economics of summer greengram (*Vigna radiata* L.). *Journal of pharmacognosy and phytochemistry*, 7(2), 404-408.
- Soumare, A., Boubekri, K., Lyamlouli, K., Hafidi, M., Ouhdouch, Y., & Kouisni, L. (2020). From isolation of phosphate solubilizing microbes to their formulation and use as biofertilizers: status and needs. *Frontiers in bioengineering and biotechnology*, 7, 425.
- Speiser, C., Baumann, T., & Niessner, R. (2000). Morphological and chemical characterization of calcium-hydrate phases formed in alteration processes of deposited municipal solid waste incinerator bottom ash. *Environmental Science & Technology*, 34(23), 5030-5037.
- Hinsinger, P., Betencourt, E., Bernard, L., Brauman, A., Plassard, C., Shen, J., ... & Zhang, F. (2011). P for two, sharing a scarce resource: soil phosphorus acquisition in the rhizosphere of intercropped species. *Plant physiology*, 156(3), 1078-1086.
- Swaminathan, R., Singh, K., & Nepalia, V. (2012). Insect-pests of green gram *Vigna radiata* (L.) Wilczek and their management. *Agricultural science*, 10, 197-222.
- Uebersax, M. A., Urrea, C., & Siddiq, M. (2022). Physical and Physiological Characteristics and Market Classes of Common Beans. *Dry Beans and Pulses: Production, Processing, and Nutrition*, 57-80.
- Varma, D., Meena, R. S., & Kumar, S. (2017). Response of mungbean to fertility and lime levels under soil acidity in an alley cropping system of Vindhyan Region, India. *Int J Chem Stud*, 5(4), 1558-1560.
- Venkatarao, C. V., Naga, S. R., Yadav, B. L., Koli, D. K., & Jagga Rao, I. (2017). Effect of phosphorus and biofertilizers on growth and yield of mungbean [*Vigna radiata* (L.) Wilczek]. *Int. J. Curr. Microbiol. App. Sci*, 6(7), 3992-3997.
- Weber, O., Delince, J., Duan, Y., Maene, L., McDaniels, T., Mew, M., ... & Steiner, G. (2014). Trade and finance as cross-cutting issues in the global phosphate and fertilizer market. In *Sustainable Phosphorus Management* (pp. 275-299). Springer, Dordrecht.

Comment [DC5]: Alphabetical order

- Withers PJA, Withers PJA, Edwards AC, Foy RH. 2001. Phosphorus cycling in UK agriculture and implications for phosphorus loss from soil. *Soil Use Manage* 17:139–149.
- Xu, X., Mao, X., Van Zwieten, L., Niazi, N. K., Lu, K., Bolan, N. S., & Wang, H. (2020). Wetting-drying cycles during a rice-wheat crop rotation rapidly (im) mobilize recalcitrant soil phosphorus. *Journal of Soils and Sediments*, 20(11), 3921-3930.
- Yi-Shen, Z., Shuai, S., & FitzGerald, R. (2018). Mung bean proteins and peptides: Nutritional, functional and bioactive properties. *Food & nutrition research*, 62.
- Yu, X., X. Liu, T. H. Zhu, G. H. Liu, and C. Mao., 2011. “Isolation and characterization of phosphate-solubilizing bacteria from walnut and their effect on growth and phosphorus mobilization.” *Biology and Fertility of Soils* 47:437-446.
- Yu, W., Ding, X., Xue, S., Li, S., Liao, X., & Wang, R. (2013). Effects of organic-matter application on phosphorus adsorption of three soil parent materials. *Journal of soil science and plant nutrition*, 13(4), 1003-1017.
- Zhu, F., Qu, L., Hong, X., & Sun, X. (2011). Isolation and characterization of a phosphate-solubilizing halophilic bacterium *Kushneria* sp. YCWA18 from Daqiao Saltern on the coast of Yellow Sea of China. *Evidence-Based Complementary and Alternative Medicine*, 2011.
- Zhu, J., Li, M., & Whelan, M. (2018). Phosphorus activators contribute to legacy phosphorus availability in agricultural soils: A review. *Science of the Total Environment*, 612, 522-537.