

## Original Research Article

### EVALUATION OF DIFFERENT VARIETIES OF CORIANDER (*Coriandrum sativum* L.)

#### ABSTRACT

A research experiment was carried out at the Horticulture Research Farm department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj, during the rabi season of year 2021-2022. The experiment were laid out in Randomized Block Design comprising of 8 varieties viz., V<sub>1</sub>: Kashmiri, V<sub>2</sub>: Japani 57, V<sub>3</sub>: PanthHaritha; V<sub>4</sub>: Green King, V<sub>5</sub>: Sughanda, V<sub>6</sub>: Simco 55, V<sub>7</sub>: Simpcos 33, V<sub>8</sub>: Simco with three replications. The observations were recorded as per the growth, yield, and quality parameters. The results revealed that the varieties V<sub>5</sub> (Sughanda) was found to be the most suitable over all the other varieties in relation to growth (value), yield (value) and quality (value) of Coriander.

**Keywords:** Coriander, growth, yield and Quality, Varieties.

#### INTRODUCTION

India, often referred to as the "Home of spices", grows about 63 spices and many of them, like coriander, cumin, fennel, and fenugreek are counted among the major seed spices of the country on the basis of area, and production and their importance to the national economy. In India, seed spices are grown in an area of approximately 5.17 lakh hectare with an annual production of about 3.35 t. It has been estimated that the world demand of seed spices by 2000 A.D. would be about 40 lakh tonnes, of which India has a larger role to play (Anonymous, 1993).

Coriander (*Coriandrum sativum*) is an important seed spice crop belonging to family Umbelliferae. It is highly cross pollinated crop being grown as irrigated as well as unirrigated. Practically all parts of coriander have their own particular appeal in foods, through tender stem, leaves, flowers and fruits are in greater demand. Coriander leaves

**Comment [CA1]:** Put your objective here

**Comment [CA2]:** Why this is in upper case, I did not understand?

**Comment [CA3]:** If the experiment was conducted two consecutive years, then were is ok but make experiment plural

**Comment [CA4]:** Remove V2.....V8, here need to write serial number just write name of the varieties

**Comment [CA5]:** Is it correct?

**Comment [CA6]:** In abstracts, you have to present result, so put the values.

**Comment [CA7]:** It may be t/ha. If overall India's production, it is wrong.

**Comment [CA8]:** We are living in 2022, we have passed 2000, a long ago. **Must change first paragraph**, google some recent articles, or visit ICAR web site you will find recent information.

**Comment [CA9]:** It is apiaceae, earlier it was Umbelliferae. You have written both, it should be one.

contain special type of flavour and people use it in the preparation of vegetable and “chatni”. Coriander seeds are used as an ingredient in ‘curry powder’ and ‘spice powder’. Its seeds also contain proteins and sugars. The coriander oil is used for preparation of ‘chocolate’ and sweets, coriander oil is also used in flavouring beverage, whisky, gin and other liquors.

Coriander is mainly a crop of tropics and subtropics, mainly cultivated in Morocco, Romania, France, Spain, Italy, Holland, Russia, Burma, Pakistan, Turkey, India, Mexico, Argentina and to some extent in England and U.S.A. (Nethajiet *al.* 1979)

~~The coriander (*Coriandrum sativum*) of family Apiaceae. It is an annual herb. Mainly cultivated for its fruit as well as its tender green leaves. The fruit have a fragrant odour and a pleasant aromatic taste. The seeds are also used as condiment.~~

Coriander is native to regions spanning from Southern Europe and Northern Africa to Southwestern Asia. It is a soft plant growing to 50 cm (20 in) tall. The leaves are variable in shape, broadly lobed at the base of the plant, and slender and feathery higher on the flowering stems.

In India, major coriander growing area is lying in semi arid climate where coriander is cultivated on conserved moisture in rabi season. India is the largest producer of coriander (*Coriandrum sativum* L.). Coriander is cultivated on an area of 530.5 thousand hectares in India with a production of 482.0 thousand MT and productivity is 900 kg per hectare.

Coriander is widely cultivated in countries like India, Morocco, Romania, France, Spain, Italy, Holland, Yugoslavia, Russia, Myanmar, Turkey, Mexico and to some extent in England and USA. In India, it is mainly cultivated in the states of Rajasthan, Gujarat, Andhra Pradesh, Madhya Pradesh and Tamil Nadu covering an area of 351710 hectares with an annual production of 259480 tonnes (Anonymous, 2007). In Rajasthan, coriander is largely cultivated in the districts of Jhalawar, Baran, Kota, Bundi, Chittorgarh, Banswara, Tonk, SawaiMadhopur, Nagpur and Jodhpur covering an area of 136741 hectares with an annual production of 142353 tonnes and productivity of 1041.04 kg/ hectare (Anonymous, 2005-06).

Coriander (*Coriandrum sativum* L.) is an important seed spice crop belongs to family Apiaceae (Umbelliferae) with a chromosome number of  $2n=22$ . Mediterranean region is the centre of origin of this crop. Coriander is an annual herbaceous cross pollinated crop. Its name has been derived from Greek word “Koris” means bed-bug, because of unpleasant, fetid bug like odour of the green un ripened fruits. Coriander ranks first among the seed

**Comment [CA10]:** The information provided here is not matching with first paragraph. Provide consistence information. Put this information in first paragraph and provide reference.

**Comment [CA11]:** You have repeated information many times. Just make three paragraphs:  
1. Coriander in relation to India  
2. Brief introduction about coriander  
3. Need of this experiment, why did you conduct this experiment, why is there a need for variety evaluation for example to provide options to farmers, earlier varieties are less productive etc

spices with respect to export. It is the major ingredient of curry powder up to 40% by weight and also forms an important ingredient for several alcoholic beverages, particularly in Seeds are also used as tonic, carminative, diuretic, stomachic and as an aphrodisiac. Oleoresin from coriander is used as a flavouring agent and as an ingredient in pharmaceutical formulation and in perfumery (Singh et al., 2006) [14]. Coriander fruits are an important spice of many countries of Europe, Northern Africa, West, Central and South Asia. In the Mediterranean region, coriander cultivation dates back to ancient Egypt; in Europe, coriander is known since the middle ages (Anonymous, 2012).

Yield in this crop is governed by a multitude of component characters and is subject to genotype and environment interaction. One of the reasons for low oil yield is due to lack of potential genotype suited to a particular region. Though, coriander is grown on commercial scale for its herbage and also seeds in most parts of the state and in North Eastern Transitional tract of Karnataka in particular, besides the fact that coriander being an economically major seed spice and is put to a variety of different uses and is also important from the earnings of foreign exchange, the crop has received little or no concerted efforts for its improvement especially in this region. Hence, the present investigation was undertaken.

## **MATERIALS AND METHODS**

This ~~experiment e~~ investigation on ~~“Evaluation of Different Variety of Coriander (Coriander Sativum L.)”~~ was carried out at the Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj U.P., which is located at 25°24'46.14" N latitude, 81°50'49.95" E longitude and 98 m above the mean sea level during Rabi season (December, 2021 – May, 2022). The minimum temperature during crop season was ~~to be~~ 21.38°C and the maximum was ~~to be~~ 37.82°C. The experiment was ~~laid out~~ ~~at~~ ~~covered in a~~ Randomized Block Design comprising of ~~eight~~ varieties of coriander with three replications. Climate, meteorological conditions, Plan layout, cultural practice and techniques employed for growth studies.

Comment [CA12]: Make this sentence more clear.

The following biometrical observations were recorded for the randomly selected plants. Plant height, number of primary branches per plant, number of secondary branches per plant, and number of leaves per plant were recorded at 30, 60 and 90 DAS. Days of 50% flowering, number of flowers per umbellets, number of umbellets per plant, number of seeds per umbel, seed yield per plant (g), seed yield per plot (g), seed yield per ha (q), TSS, Ascorbic acid, etc.

As per the existing market prices, the input and output costs were computed treatment wise and different economic parameters viz., Cost of cultivation, gross return, net return and benefit-cost ratio were calculated.

**Comment [CA13]:** Just mention, which statistical tool that you used to analyse your data.

## **RESULT AND DISCUSSION**

The data related to plant height at 30, 60 and 90 DAS is presented in table 1. It is evident that all the varieties tried in this experiment produced considerable amount of changes in plant height at successive stages of growth.

**Comment [CA14]:** Make one paragraph for each parameter. When you are writing the name of the variety, you need to write v1, v2 etc.

At 30 DAS, the maximum plant height was recorded in V<sub>2</sub> (Japani 57) (21.16 cm) followed by V<sub>6</sub> (Simco 55) (20.83 cm), V<sub>8</sub> (Simco) (20.16 cm), V<sub>7</sub> (Simpco s 33) (17.86), V<sub>1</sub> (Kashmiri) (17.56) and V<sub>5</sub> (Sughanda) (16.33) which were on par. Whereas the minimum plant height (14.26) was found to be in V<sub>3</sub> (Pant Haritha).

At 60 DAS, the maximum plant height was recorded in V<sub>2</sub> (Japani 57) (68.50 cm) which were on par. Whereas the minimum plant height (48.5 cm) was found to be in V<sub>3</sub> (Pant Haritha).

At 90 DAS, the maximum plant height (100.11 cm) was recorded in V<sub>2</sub> (Japani 57) followed by V<sub>6</sub> (Simco 55) (99.56 cm) which were on par. Whereas the minimum plant height (76.06 cm) was found to be in V<sub>3</sub> (Pant Haritha).

The differences in plant height among the varieties might be due to the genetic makeup of the plant and its expression to the growing soil and environmental conditions. The variation in plant growth of different coriander varieties were also observed by **Kalid asueta**., (2008) and **Verma et al.**, (2014) in coriander.

**Formatted:** Tab stops: 5.32", Left

The data related to number of leaves per plant at 30, 60 and 90 DAS is presented in table 1. It is evident that all the varieties tried in this experiment produced considerable amount of changes in number of leaves per plant at successive stages of growth.

At 30 DAS, the maximum number of leaves was recorded in V<sub>2</sub> (Japani 57) (23.28) followed by V<sub>6</sub> (Simco 55) (23.26), V<sub>8</sub> (Simco) (23.15), V<sub>7</sub> (Simpco s 33) (23.10), V<sub>1</sub> (Kashmiri) (21.30) and V<sub>5</sub> (Sughanda) (21.26) which were on par. Whereas the minimum number of leaves (20.00) was found to be in V<sub>3</sub> (Pant Haritha).

At 60 DAS, the maximum number of leaves was recorded in V<sub>2</sub> (Japani 57) (38.33) followed by V<sub>6</sub> (Simco 55) (37.31) which were on par. Whereas the minimum number of leaves (30.11) was found to be in V<sub>3</sub> (Pant Haritha).

**Comment [CA15]:** All N should be in lower case

At 90 DAS, the maximum Number of leaves (62.66) was recorded in V<sub>2</sub> (Japani 57) which were on par. Whereas the minimum Number of leaves (48.17) was found to be in V<sub>3</sub> (Pant Haritha).

The differences in Number of leaves among the varieties might be due to the genetic makeup of the plant and its expression to the growing soil and environmental conditions. The variation in plant growth of different coriander varieties were also observed by **Verm aetal., (2014)** and **Meena et al., (2014)** in coriander.

The data related to Number of primary branches per plant are presented in table 1. The maximum number of primary branches per plant V<sub>2</sub> (Japani 57) (10.58) followed by V<sub>6</sub> (Simco 55) (8.97), V<sub>8</sub> (Simco) (8.63), V<sub>7</sub> ( Simpcos 33) (8.53), V<sub>1</sub> (Kashmiri) (8.50), V<sub>5</sub> (Sughanda) (8.13), V<sub>4</sub> (Green King) (7.87) and V<sub>3</sub> (Pant Haritma) (7.80) which were on par.

Increase in number of primary branches could be attributed to particular genotype in condition on which it is grown. Similar variations among varieties with respect to primary branches were reported by several workers **Venkatareddy et al., (1986)**, **Shridharet al., (1990)**, **Selvarajane et al., (2002)** and **Velayudham, (2004)** in coriander genotype grown in different agro climatic condition.

**Comment [CA16]:** Is it in the journal format? if not, do not make it bold.

The data related to Number of secondary branches per plant are presented in table 1. The maximum number of secondary branches per plant V<sub>2</sub> (Japani 57) (18.95) followed by V<sub>6</sub> (Simco 55) (18.93), V<sub>8</sub> (Simco) (16.67) and V<sub>7</sub> ( Simpcos 33) (15.90) which were on par. Whereas the minimum number of secondary branches per plant was found to be in V<sub>3</sub> (Pant Haritma) (10.73).

These variations in secondary branches may be due to its genetic characters and due to interaction with environment. There are several reports indicating variation with respect to number of secondary branches by several workers in coriander, Hari Prasad rao and **Srinivasrao (2001)**, **Selvarajane et al., (2002)** and **Saxena et al., (2005)**.

The data related to the coriander in days 50 % flowering are presented in table 1. The minimum days taken for 50 % flowering per plant V<sub>2</sub> (Japani 57) (57.27) and whereas the maximum days to 50% flowering was found in V<sub>3</sub> (Pant Haritma) (44.93) respectively.

Growth is one of the important characters, which determines the earliness of the variety which is controlled by the genetic factors. Similar results were reported by

**Moniruzzaman et al., (2013)** and **Malik and tehlan (2013)**.

The data related to the coriander in number of flowering per umbellet are presented in table 1. The minimum number of days taken for flowering per umbellet V<sub>2</sub> (Jaypani 57) (8.3) followed by V<sub>6</sub> (Simco 55) (8.33), V<sub>8</sub>(Simco) (8.03), V<sub>7</sub> ( Simpcos 33) (7.9), V<sub>1</sub> (Kashmiri) (7.56), V<sub>5</sub>(Sughanda) (7.53), V<sub>4</sub> (Green King) (7.53) and V<sub>3</sub> (Pant Haritma) (6.83) which were on par.

Growth is one of the important characters, which determines the earliness of the variety which is controlled by the genetic factors. Similar results were reported by **Moniruzzaman et al., (2013)** and **Malik and tehlan (2013)**.

The data related to number of umbellets per plant are presented in table 1. The highest number of umbellets per plant was recorded in V<sub>2</sub> (Japani 57) (239.63). Whereas the minimum number of umbellets per plant was found to be in V<sub>3</sub> (Pant Haritma) (141.23).

This variation in the yield could be attributed to genotypic character and response of the particular genotype to the specified environment conditions. These results are more or less in consonance with the results of **Giridhar and Sarada (2005)**, **Moniruzzaman et al.(2013)** and **Malik and Tehlan (2013)**.

The data related to number of seeds per umbel are presented in table 1 it shows that there were significant differences among the varieties during the growth stages of the crop.

The highest number of seeds per umbel (57.73) was recorded in V<sub>2</sub> (Japani 57) (28) followed by V<sub>6</sub> (Simco 55) (27.4) and V<sub>8</sub>(Simco) (26.16) which were on par. Whereas the lowest number of seeds per umbel was recorded in V<sub>3</sub> (Pant Haritma) (21.9).

This variation in the yield could be attributed to genotypic character and response of the particular genotype to the specified environment conditions. These results are more or less in consonance with the results of **Giridhar and Sarada (2005)**, **Moniruzzaman et al.(2013)** and **Malik and Tehlan (2013)**

The data related to seed yield per plant are presented in table 1. The highest seed yield per plant was recorded in V<sub>2</sub> (Japani 57) (6.03) followed by V<sub>6</sub> (Simco 55) (4.86), V<sub>8</sub>(Simco) (4.78), V<sub>7</sub> ( Simpcos 33) (4.6), V<sub>1</sub> (Kashmiri) (4.56), V<sub>5</sub> (Sughanda) (3.8), V<sub>4</sub> (Green King) (3.51) and V<sub>3</sub> (Pant Haritma) (3) which were on par.

The difference in seed yield per plant might be due to the genotypic difference

and ecological variation. The results are more or less in consonance with the results of **Moniruzzaman et al. (2013)** and **Malik and Tehlan (2013)**.

The data related to seed yield per plot are presented in table 2 and it shows that there were significant differences among the varieties during the stages of the crop growth.

The highest seed yield per plot was recorded in V<sub>2</sub> (Japani 57) (478.67). Whereas the lowest seed yield per plant was recorded in V<sub>3</sub> (Pant Haritma) (101.13).

Considerable variations in the seed yield among the genotypes and varieties of coriander were also reported earlier by several workers in coriander, **Velayudham (2004)**, **Giridhar and Sarada (2005)**, **Chaulagain et al., (2011)**, **Palanikumar and Rajamani (2012)**, **Malik and Tehlan (2013)** and **Moniruzzaman et al., (2013)**.

The data related to seed yield per plot are presented in table 2. It shows that there were significant differences among the varieties during the stages of the crop growth.

The highest seed yield per plot was recorded in V<sub>2</sub> (Japani 57) (22.03) followed by V<sub>6</sub> (Simco 55) (21.03) and V<sub>8</sub> (Simco) (20.30) was statistically on par. Whereas the lowest seed yield per plant was recorded in V<sub>3</sub> (Pant Haritma) (16.03).

Considerable variations in the seed yield among the genotypes and varieties of coriander were also reported earlier by several workers in coriander, **Velayudham (2004)**, **Giridhar and Sarada (2005)**, **Chaulagain et al., (2011)**, **Palanikumar and Rajamani (2012)**, **Malik and Tehlan (2013)** and **Moniruzzaman et al., (2013)**.

The data related to TSS Brix are presented in table 3. The highest TSS Brix was recorded in V<sub>2</sub> (Japani 57) (6.03) followed by V<sub>6</sub> (Simco 55) (4.86), V<sub>8</sub> (Simco) (4.78), V<sub>7</sub> (Simco s 33) (4.6), V<sub>1</sub> (Kashmiri) (4.56), V<sub>5</sub> (Sughanda) (3.8), V<sub>4</sub> (Green King) (3.51) and V<sub>3</sub> (Pant Haritma) (3) respectively.

The difference in TSS Brix might be due to the attributed lower starch content which is being a primary metabolite, may effect the accumulation of secondary metabolites like TSS. This is in confirmation with the studies of earlier workers **Agarwal et al., 1990**; **Rajagopalan et al., 1996**; **Prabhu and Balakrishnamurthy 2006**; **Velayudham et al., 2004**. The results are in conformity results of the (**palanikumar and rajamani, 2012**).

The data related to Ascorbic acid are presented in table 3. The highest Ascorbic acid was recorded in V<sub>2</sub> (Japani 57) (208.60) and the lowest Ascorbic acid content was found in

variety V<sub>3</sub> (Pant Haritma) (151.72) respectively.


The difference in TSS Brix might be due to the attributed lower starch content which is being a primary metabolite, may affect the accumulation of secondary metabolites like TSS. This is in confirmation with the studies of earlier works **Agarwal et al., 1990; Rajagopalan et al., 1996; Prabhu and Balakrishnamurthy 2006; Velayudham et al., 2004**. The results are in conformity with the results of the **(palanikumar and rajamani, 2012)**.

## **CONCLUSION**

From the present investigation, ~~it is concluded that~~ the ~~Variety V<sub>2</sub> coriander variety J (japani 57)~~ was found ~~the to be the~~ most suitable ~~in rabi season for Prayagraj~~ over ~~all the other variety the others~~ in relation to plant height, number of primary branches per plant, number of secondary branches per plant, number of leaves per plant, days of 50% of flowering, number of flowers per umbellet, number of umbels per plant, number of seeds per plant (g), seed yield per plot (g), seed yield per ha (q), gross return, net return, and highest cost benefit ratio (1.68:1) of coriander.

**Table 1. Evaluation of different varieties of coriander in Plant Height, No. of leaves per plant, No. of primary branches, No. of secondary branches, Days of 50% flowering, No. of flower per umbellet and No. of umbellets per plant.**

UNDER PEER REVIEW



Notation	Varieties	Days of 50% flowering	Plant Height (cm)			No. of leaves per plant			No. of primary branches	No. of secondary branches	No. of flower per umbellet	No. of umbellets per plant
			30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS				
V1	Kashmiri	49.00	17.56	52.58	86.5	21.30	33.79	52.31	8.50	15.30	7.56	172.03
V2	Japani-57	44.93	21.16	68.5	100.11	23.28	38.33	62.67	10.58	18.95	8.43	239.63
V3	PanthHaritha	57.27	14.26	48.5	76.06	20.00	30.11	48.17	7.80	10.73	6.83	141.23
V4	Green King	46.09	14.51	48.7	77.63	20.27	31.01	49.13	7.87	14.47	7.5	156.16
V5	Sughanda	47.97	16.33	50.66	81.8	21.26	33.06	49.17	8.013	15.00	7.53	160.26
V6	Simco 55	53.93	20.83	57.7	99.56	23.26	37.31	59.163	8.97	18.93	8.33	218.70
V7	Simpeco s 33	49.50	17.86	52.6	88.48	23.10	35.58	55.93	8.53	15.90	7.9	178.20
V8	Simco	51.10	20.16	52.86	89.7	23.15	35.97	57.83	8.63	16.67	8.0	191.16
	F- Test	S	S	S	S	S	S	S	S	S	S	S
	S.Em±	0.34	0.38	0.31	0.34	0.23	0.24	0.22	0.15	0.26	0.14	0.30
	SE(d±)	0.48	0.54	0.45	0.48	0.33	0.34	0.31	0.21	0.38	0.20	0.43
	CV (%)	1.00	5.59	1.53	1.01	0.70	0.71	0.64	0.45	0.79	0.42	0.90
		1.77	1.12	0.93	1.00	2.82	1.85	1.05	4.66	4.45	4.83	0.44

**Comment [CA17]:** If it is for significant, you have to write it below your table. Standard practice is to provide asterisk (\*) for different level of significance.

**Comment [CA18]:** I did not understand this row.

**Table 2. Evaluation of different varieties of Coriander in No. of seed per umbel, Seed yield per plant (g), Seed yield per plot (Kg), Seed yield per ha (q)**

Notation	Varieties	No. of seed per umbel	Seed yield per plant (g)	Seed yield per plot (Kg)	Seed yield per ha (q)
V1	Kashmiri	23.5	4.56	244.67	18.40
V2	Japani-57	28	6.03	478.67	22.03
V3	PanthHaritha	21.9	3	101.13	16.03
V4	Green King	22.43	3.51	154.33	16.80
V5	Sughanda	22.76	3.8	207.67	17.53
V6	Simco 55	27.4	4.86	372.00	21.03
V7	Simpcos 33	24.5	4.60	252.00	19.23
V8	Simco	26.16	4.7	316.33	20.30
	F- Test	S	S	S	S
	S.Em±	0.20	0.08	1.70	0.17
	SE(d±)	0.28	0.11	2.40	0.24
	CV (%)	0.58	0.24	4.98	0.50
		2.09	4.98	1.66	2.35

**Table 3. Evaluation of different varieties of Coriander based on quality parameters TSS °Brix and Ascorbic acid (mg/100gm).**

<b>Notation</b>	<b>Varieties</b>	<b>TSS°Brix</b>	<b>Ascorbic acid (mg/100gm)</b>
V1	Kashmiri	5.36	141.00
V2	Japani-57	5.98	160.50
V3	PanthHaritha	4.83	127.33
V4	Green King	4.84	131.00
V5	Sughanda	5.26	137.00
V6	Simco 55	5.96	157.67
V7	Simpcos 33	5.66	146.67
V8	Simco	5.76	154.83
	F- Test	S	S
	S.Em±	0.09	0.67
	SE(d±)	0.14	0.94
	CV (%)	0.29	1.96
		4.67	1.20

## REFERENCE

- Agarwal, S. and Sharma, R.K., 1990** Variability of quality aspects of seed spices and future strategy. *Indian Cocoa, Arecanut and spices J.*, **13**(4): 127-129.
- Agrawal, S., Sharma, R.K. and bhatt, B.N., 1990**, Quality evaluation in coriander. *Indian Cocoa, Arecanut and spices J.*, **13**(4): 137-137.
- Anonymous (2012)**. Cultivation of coriander for spice and quality seed. *Spice India*, **25**(4): 7-15
- Anonymous, 1993**. All India coordinated research project on spices. Annual report, National research centre for spices Calicut.
- Ashish Kumar Maurya, M.L. Kushwaha, Vikas Kumar Jain and Neeraj Singh (2016)** *Evaluation of chilli (Capsicum annum L.) genotypes for yield and performance against diseases. Progressive Research Journal* ISSN: 0973-6417, Vol 11: 4615-4617.  
<https://www.researchgate.net/publication/311309867> Evaluation of chilli *Capsicum annum L* genotypes for yield and performance against diseases
- Abdul Kaium, M. Islam, S. Sultana, Mahjuba Akter, E. Hossain, A. Mahjuba ( 2015)** Yield and Yield Contributors of Coriander (*Coriandrum Sativum L.*) as Influenced by Spacing and Variety. *International Journal of scientific and research publications*, Vol5, Issue 3, ISSN 2250-3153.  
<https://www.researchgate.net/publication/273141866> Yield and Yield Contributors of Coriander *Coriandrum Sativum L* as Influenced by Spacing and Variety .
- Bhati, D.S. 1988**. Effect of leaf plucking on growth, yield and economics of coriander varieties under semi-arid conditions. *Indian J. Agronomy* **33** (3) 242-244.
- Chaulagain, R., Pant, S. S., Thapa, R. B. and Sharma, M. D. (2011)**. Performance of coriander cultivars for green leaf production under late sowing condition. *The J. Agri. and Env.*, **12**:67-73.
- Duwal, A. Nepal, S. Luitel, S. Acharya, R. Pathak, P.R. Poudel and J. Shrestha (2019)** Evaluation of coriander (*Coriandrum sativum L.*) varieties for growth and yield parameters. *Nepalese Journal of Agricultural Sciences*, vol. **18** ISSN 2091- 0428.

- Giridhar, k. and Sarada, C., 2005.** Identification of coriander (*Coriandrum sativum* L.) genotypes for varieties of Andhra Pradesh. *Nat. symp. Cur. Trends in onion, garlic, chilies and seed spices-production, marketing and utilization, SYMSAC-II*, 25-27 November, NRCOG, Rajgurunagar, pp-92.
- Edison, S. and joshi A. 1990.** New varieties to improve productivity of seed spices. *Indian cocoa, Areca, spices J. XIII* (4): 121-23.
- Hariprasadrao, N. and Srinivasrao, G. (2001).** Studies on the performance of exotic and indigenous coriander (*Coriandrum sativum* L.) genotypes for greens. *The Andhra Agric. J.*, **48**(3-4): 324-326.
- Jackson, M.L. (1958)** Soil chemical analysis Second edition Indian reprint, prentice hall of India, New Delhi. 498.
- Kader, M. Jem, M.D. Abdul kader and Muthuswami, S. 1984.** Coriander- A crop of good prospectus for Tamil nadu. *Indian Cocoa, Areca, Spices, J. VIII* (1): 5-6.
- Kalidasu G, Sarada C and Reddy T Y (2008).** Efficacy of biofertilizers on the performance of rainfed coriander (*Coriandrum sativum* L.) in vertisols. *J Spices and Aromatic Crops* **17**(2): 98- 102.
- Lavanya Gandepalli and V.M. Prasad (2020)** Evaluation of coriander (*Coriander sativum* L.) Varieties in prayagaraj Agro-climatic conditions. ISSN: 2319-7706 vol9. No. (12).
- Malee Ram Jhajhra, D.K. Rana and Arjun Lal Ola (2017)** Evaluation of fenugreek (*Trigonella Foeniculum- Graecum* L.) Varieties under sub-tropical Condition of Garhwal Himalayas. Research Article. ISSN 2278-6783.
- Malik, T.P., and Tehlan, S.K., (2013)** Performance of coriander (*Coriander sativum* L.) varieties for growth and seed yield. *International Journal seed spices*, **3**: 89-90.
- M. Ahsan Altaf, Rabia Shahid, M. Asad Altaf, M. Mohsin Altaf (2019)** Effect of NPK, Organic manure and their combination on growth, yield and nutrient uptake of chilli (*Capsium Annum* L.) *Horticultural International Journal* . Volume 3 Issue 5- 2019. DOI: [10.15406/hij.2019.03.00135](https://doi.org/10.15406/hij.2019.03.00135)
- Moniruzzamani, M., Rahman, M.M., Hossain, M.M., Sirajul, K.A.J.M., and khaliq, Q.A., (2013)** Evaluation of coriander (*Coriander sativum* L.) genotypes for seed yield and yield contributing characters. *Bangladesh J. Agri. Res.*, **38**(2): 189-202.

- MukeshAwasthi, Devi Singh and Vijay Bahadur (2021)** Varietal evaluation of chilli (*Capsicum annuum*) for growth, yield and quality in Prayagraj Agro climatic condition. *The pharma Innovation Journal*; 10(10): 1267-1269.
- Nethaji, Seetharaman, R. and Arjun. G. (1979).** Coriander the wonder spice. Indian Farmers Digest 12:33-34.
- Nirmalya Dhal, VM Prasad, Samir E topno, Vijay bahadur and Shailesh Marker (2021)** Varietal trails of chilli (*Capsicum spp.*) varieties on the basis of growth and yield in PrayagrajAgroclimaticconditons. *The pharma Innovation Journal*; 10(10): 424-426.
- Olsen, S.R., Cole, C.V., Watnahe, F.S. and Dean, L.A. (1954)** Estimate of available Phosphorus in soil by extraction with sodium bicarbonate US. Deptt. Agricciric. 939.
- Prasad, R., Kumar, V.,and Prasad, K., (2014).** Nanotechnology in sustainable agriculture:Present concerns and future aspects. *African J. of Biotechnology*
- Palanikumar, M. and Rajamani, K., (2012)** Evaluation of coriander (*CoriandrumsativumL.*) genotypes for fresh, dry biomass yield and oil content under different seasons., *j. Crop Res.*, 44(1&2):194-202.
- PratapNaikwade (2014)** Evaluation of Leaf Litter Compost and Vermicompost on Yield and Nutrient Uptake of Trigonella.
- <https://www.researchgate.net/publication/314899010> Evaluation of Leaf Litter Compost and Vermicompost on Yield and Nutrient Uptake of Trigonella. *Indian Journal of Applied Research* 4(2): 5-3.
- Phurailatpam, A.K., Geetha, K.A., Meena, R. S., Maiti, S., (2014)** Evaluation of coriander (*CoriandrumsativumL.*) cultivars for yield and yield contributing characters in Gujarat *J. Spices and Aromatic crops.*, 25(1): 7-12.
- Prabhu, T. and Balakrishnamurthy, G., (2006)** Evaluation of coriander (*CoriandrumsativumL.*) accessions under irrigated conditions for growth, yield and quality. *Proc. Nat. Sem. Emerging trends in production, uality, Processing and export of spices.*,28-29 March, Coimbatore pp 13.
- Rajagopalan, A., Azhakiyamanavalan., R.S. and Abdul-khader, M.D., 1996.** Evaluation of coriander cultivars for yield. *Indian Cocoa, Arecanut and spices J.*, 20(1):13-14.

- Rao, T.S. Rao, J.N. and Dashrathi, T.B. 1979.** CJ-2 (Lam Selection)- A promising coriander type for Andhra Pradesh. *Indian Cocoa, Areca and spices J.III* (2): 44-45
- Rao, T.S. and Babu, M.K. 1979.**Improvement of minor spices in Andhra-Pradesh. *Indian Cocoa, Areca. And Spices J.III* (1): 3-4.
- Ravi Pujari, B.B. Sunanda, A.R. Kurubar, Jaiprakash Narayan, T. Chetan and Satish Kale (2019).** Collection and Evaluation of Coriander Varieties for Growth and Seed Purpose in UKP Command Area. *Int.J.Curr.Microbiol.App.Sci.* 8(6): 3125-3130. doi: <https://doi.org/10.20546/ijcmas.2019.806.372>
- Rs Mishra and Vp Pandey (2015)** Evaluation of varieties of coriander (*Coriandrum sativum*L.) for resistance to stem gall disease and seed yield, current Advances in Agricultural Sciences, vol - 7, Pages (151-153).
- Sharma, R.K. and Bhati, D.S. 1985.** Performance of fennel varieties under irrigated conditions. *Indian cocoa, Areca. Spices J. IX* (17): 16.
- Sharma, R.K. and Bhati, D.S. 1987.**Evaluation of fenugreek varieties under irrigated conditions. *Indian cocoa, Areca. Spices J. X* (4): 89-91.
- Sharma, R.K. and Bhati and Agarwal, H.R. 1989.** Performance of Coriander varieties under irrigated conditions. *Indian cocoa, Areca. Spices J. XI* (3): 95-96..
- Saxena, R. P., Pandey, V. P., Datta, J. and Gupta, R. K. (2005).** Performance of coriander entries at Kumarganj, Faizabad. *Nat. Symp. Cur. Trends in Onion, Garlic, Chillies and Seed Spices-Production, Marketing and Utilization, SYMSACII*, 25-27 November, NRCOG, Rajgurunagar, pp.55-56.
- Selvarajan, M., Chezhiyan, N., Muthulakshmi, P. and Ramar, A. (2002).** Evaluation of coriander genotypes for growth and yield. *South Indian Hort.*, **50**(4-6):458-462.
- Shridhar, Sulikeri, G.S. and Hulamani, N.C. (1990).** Performance of coriander (*Coriandrum sativum*L.) genotypes.
- Subbiah, B.V., and Asija, G.L., (1956)** A rapid procedure for the estimate of available Nitrogen. *International journal of Soil Current Science.* 25: 259-260.

- Toth, S.J. and Prince, A.L. (1949)** Estimate of cation exchange capacity and exchangeable Ca, K, Na, content of soil by flame photometer technique soil. *Science direct* **67**: 439-445.
- Velayudham, A. (2004)**.Evaluation and effects of organics with bio-inoculants in coriandervar.Co3. M.Sc. (Hort.) Thesis,Univ.Agric. Sci.,Dharwad.
- Venkatareddy,P.,Sriramarao,T.,Narasimharao,S.B.S.andNarisireddy,A.(1986)**. Geneticvariabilityincoriander.*Indian,ArecanutandSpicesJ.*,**10**(3):90-92.
- Verma P, Doshi V and Solanki R K. (2014)**. Genetic variability assessed in Coriander(*Coriandrumsativum* L.) over years under environmental conditions of SouthEasternRajasthan (HadotiRegion). *IntJ SeedSpices* **4**(2):94-95.
- Wakley, A. and Black, I.A. (1947)** Critical examination of rapid method for determining organic carbon in soils, effect of variance in digestion conditions and inorganic soil constituents. *Soil science.* 632:251.
- Wilcox.L.V.(1950)** Electrical Conductivity Am. Water work Associate journal. **42**: 775-776.
- Yogesh Kumar Agarwal, Ramchandra, Hemant Kumar and Anil Kumar** Performance analysis and economics of coriander (*coriandrumsativum* l.) Cultivation under subabul (*leucaenaleucocephala*) based alley cropping system. Vol 20 No. 1, **2020** pp. 1970-1974