

# **Effect of Hydromethanolic Fruit Pulp Extract of *Azanza garckeana* (Malvaceae) on Sexual Behavior and Histology of Female Wistar rats**

## **ABSTRACT**

**Aim:** To evaluate the effect of hydromethanolic fruit pulp extract of *Azanza Garckeana* on sexual behavior and histology of female Wistar rats.

**Study design:** Twenty-five female Wistar rats and five (5) male Wistar rats were used for the study, the female animals were randomly divided into 5 groups (n= 5): Group I served as normal control and received 1 ml/kg distilled water, group II served as positive control and received 100 µg estradiol valerate and 50 µg progesterone 48 hr and 5 hr respectively before the sexual behavioral test, while group III, IV and V received 250 mg/kg, 500 mg/kg and 1000 mg/kg body weight of the extracts respectively for 14 days.

**Place and Duration of study:** Department of Human Physiology (animal house), Ahmadu Bello University, Zaria, between September 2021 and November 2021.

**Methodology:** Sexual behavioral tests were performed with sexually experienced male rats 1:1 after the treatment period in a unilevel pacing chamber and the process videotaped for proper scoring. Proceptive behaviors and Lordosis was extrapolated as index for the level of proceptivity and receptivity respectively in the female animals. The proximity time of the female to the male was also recorded. At the end of the experiment, the rats were anaesthetized with combined doses of ketamine and diazepam injection 0.4 mg/kg. Blood sample was collected for hormonal assays which were done using ELISA kit. The ovaries were dissected out and fixed in 10% formalin for histological examination. Phytochemical screening was also carried out on the extract to determine the presence and absence of secondary metabolites.

**Results:** The hormonal assay result revealed significant ( $p = .000$ ) increase in the serum Estradiol levels across the treatment groups but no significant change in progesterone level except at 500mg/kg dose where it was significantly increased when compared to both negative and positive control. Receptivity and proceptivity in the extract treated groups and positive control groups increased significantly when compared with the negative control group, indicating enhanced sexual behaviour. The proximity time also significantly increased ( $p = .000$ ) in the extract treated groups when compared to negative control. The histological sections of ovaries revealed increased follicle development in the treated groups when compared to negative control. The presence of flavonoid and saponin in the fruit pulp extract could be the reason for the high aphrodisiac activity.

**Conclusion:** This study shows *Azanza garckeana* hydromethanolic fruit pulp extract demonstrated aphrodisiac property and could be used for the treatment of women with hypoactive sexual desire.

**Keywords:** *Azanza garckeana*, sexual behavior, histology, estrogen, progesterone, lordosis quotient, paracopulatory behavior

## 1. INTRODUCTION

The World Health Organization (WHO) defines sexual health as a state of physical, emotional, mental and social well-being in relation to sexuality, not merely the absence of disease, dysfunction or infirmity [1]. Good sexual health requires a positive and respectful approach to sexuality and sexual relationships, as well as having pleasurable and safe sexual experiences [2]. Sexual relationship is amongst the most important social and biological relationships in human life as it has proven to enhance an individual's quality of life and well-being [3]. Sexual and other physiologic functions diminish with advancing age [4]. Apart from aging people, healthy young people who are unable to carry out sexual act efficiently or entirely, normally suffer from psychogenic, organic, or mixed etiologies leading to sexual dysfunction [5]. Sexual dysfunction refers to difficulties that occur during the sexual response cycle that prevent one from experiencing satisfaction from sexual activity [6]. Female sexual dysfunction (FSD) may manifest as apparently unrelated emotional disorders that could degrade one's quality of life and destroy relationships, amongst family, social sphere and even the workplace [7], these problems makes some individuals resort to using sexual enhancing substances.

Over a long period of time, people have devised ways to increase sexual desire and improve sexual health. This has led to the discovery and use of substances called aphrodisiacs [8]. The term aphrodisiac has been used for substances that enhance sexual activity and are helpful in treating sexual disorders [9]. Aphrodisiacs are defined as drugs, food, scents or devices that can arouse or increase sexual drive or boost libido, most aphrodisiacs carry out their function either by altering the level of specific neurotransmitters or specific sex hormones such as progesterone, estrogen and testosterone in the body [10]. The use of plants or their products to treat sexual disorders or improve on sexual performance has a long history in most countries, and their investigations in animals have proven that they are effective in improving sexual desire and sexual behavior in animals [11]. *Azanza garckeana* plant locally called Goron tula (Hausa, Nigeria), variously it is called snotappel (Afrikaans), chinga, mukole (Bemba), azanza, tree hibiscus, snot apple, quarters, wild hibiscus, African chewing gum (English), muneko (Lozi), mukole (Lunda), uxhakuxhaku (Ndebele), mkole (Nyanja), mutohwe (Shona), mtobo (Swahili), muneko (Tongan) and morajwa (Tswana) [12]. *Azanza garckeana* grows naturally in semi-arid areas [13], in Nigeria *Azanza*

*garckeana* is found abundantly in Tula, a town in Kaltungo Local Government Area of Gombe State, Nigeria. It is also found around Kankiya, Katsina State and Daggish Kali hills of Zah district, Michika local government area of Adamawa State [14].

Among the Tula people of North-Eastern Nigeria, ripened fruits are consumed as an aphrodisiac and for treatment of infertility in men. Other parts of the plants have been formidably studied with numerous reports of isolated compounds [15], in Zimbabwe, an infusion of the roots is dropped into the ear to treat earache or orally as an antiemetic or to treat cough, chest pain, menstruation and in large doses as an abortifacient [16,17]. A tea of the stems and leaves is taken to treat liver problems. A poultice of the pounded fruit is applied to abscesses to sufficiently thin out inflamed tissue encompassing the infected cavity or draw pus to a head so the abscess may rupture [18]. The fruit of this important ethno-medicinal plant has not been previously investigated with view of isolating its medicinal principles [19].

## **2. MATERIALS AND METHODS**

### **Reagents**

Ketamine, methanol, formalin, diazepam, Estradiol valerate tablets, progesterone tablets, ELISA kits (Estradiol and progesterone), methylated spirit.

### **Instruments**

Bowl, cages, clinical hand gloves, cotton wool, drinkers, uni-level pacing chamber, hand towel, cannula, permanent markers, sawdust, stop watch, syringe, weighing balance.

### **Experimental Animals**

Thirty-seven (37), age between 2-3 months (110-140 g) female Wistar rats and five (5) sexually experienced male Wistar rats (200-300 g) were used for the research. The animals were obtained from animal house of Physiology department, Ahmadu Bello University (ABU), Zaria. The animals were housed in plastic cages and kept at room temperature, they were fed with standard animal feed and water *ad libitum*. The study was carried out in accordance with the principles of laboratory animal care and standard experimental procedure.

## **Fruit pulp sourcing and extraction**

Fresh fruit of *Azanza garckeana* was sought from Kaltungo local government of Gombe state and confirmed taxonomically in the Department of Botany, A.B.U Zaria. Voucher number ABU07276. The fruit pulp was air dried after removal of the seeds and pulverized. The pulverized sample was macerated in 70% aqueous methanol and allowed to stand for 72 hours and then filtered. The resulting filtrate was concentrated using rotary evaporator and evaporated to dryness on a water bath at 50°C. The dried extract was stored properly in a container before further experiments.

$$\% \text{yield} = \frac{\text{weight of dry extract (g)}}{\text{weight of dry fruit (g)}} \times 100$$

$$\% \text{yield} = \frac{278.5}{1000} \times 100 = 27.8\%$$

## **Methods**

### ***Determination of the Estrous Phase***

The phases of the estrous were determined using vaginal smear according to the method of Long and Evans [20]. This was done based on cytology of vaginal smears obtained daily in the morning (9 – 10 am). Vaginal secretions were collected using a 1 ml syringe filled with 0.02 ml of normal saline (0.9% NaCl) by inserting the tip into the rat vagina but not deeply. The vaginal fluid was placed on glass slides. A different glass slide for each cage, one drop was collected with a clean tip from each rat and viewed unstained under a light microscope with 10 and 40 x objective lenses. Rats in their estrous phases were used for the experiment.



**Image I:** Vaginal secretion viewed with x40 magnification showing all cell types indicating Metestrus. NE= nucleated epithelial cells AC= anucleated cornified cell L= leucocytes.

### **Experimental Design**

Twenty-five (25) adult female Wistar rats were randomly divided into 5 groups of five (5) animals each (n = 5).

**Table 1: Experimental design**

<b>Groups (n=5)</b>	<b>Treatment</b>	<b>Dosage / Kg body weight</b>
<b>I</b>	Distilled water	1 ml
<b>II (positive control) [21]</b>	Estradiol valerate	100 µg 48 hours before testing orally
	Progesterone	50 µg five (5) hours before testing orally
<b>III</b>	Fruit pulp extract	250 mg
<b>IV</b>	Fruit pulp extract	500 mg
<b>V</b>	Fruit pulp extract	1000 mg

The five sexually experienced male animals were used for the sexual behavior study using 5: 1 ratio.

### **Assessment of Sexual Behavior**

A unilevel pacing chamber (60 L x 40 W x 40 H cm) was used. The chamber was bisected by a transparent Plexiglass wall containing 3 small holes permitting the female to enter or exit the half of the cage in which the male was confined (Image II). The holes were too small for the male rats to pass through, thus allowing the female to control or “pace” the sexual interaction. During the first two weeks of the laboratory work the animals (female and male) were habituated separately to the pacing chamber twice a week for two weeks. To become sexually experienced, male rats received 4 training test sessions (twice a week for 2weeks) with non-experimental receptive females. On the day of the study, experimental female rat (in their estrous phase) were placed in one half of the chamber and a stimulus male was introduced into the other half after 5 minutes. Appetitive and consummative behaviors were evaluated for a period of 30 minutes per female animal. All frequencies of hopping, darting, and ear wiggling during the test was added and divided by the length of the test to establish a standard measure of proceptive behavior [22]. Lordosis quotient was calculated by dividing the number of lordosis responses by the total number of mounts and multiplied by 100.

$$Lq = \frac{NL}{Nm} \times 100$$

Lq: lordosis quotient NL: number of lordosis response Nm: total number of mounts



## **Image II: Copulatory unilevel pacing chamber.**

### ***Assessment of Hormonal levels***

After oral administration of the extract for fourteen (14) days, rats were anaesthetized with combined doses of ketamine and diazepam injection 0.4mg/kg given intra-peritoneally. Each rat was dissected to expose the heart and blood samples were collected in plain bottles by cardiac puncture. The blood sample was centrifuged and the serum obtained was used for hormonal analyses [23]. The serum level of estradiol and progesterone was analyzed using enzyme linked immunosorbent assay (Elisa) kits by the method of Tietz [25] following the manufacturer's manual. The principle of the immunoassays follows the typical competitive binding scenario. Competition occurs between unlabeled hormone (present in standards, controls and samples) and an enzyme-labelled hormone (conjugate) for a limited number of antibody binding sites on the microwell plate.

### ***Histological studies***

At the end of the behavioral study, the ovaries were carefully dissected out and was fixed in 10% formalin and processed for embedding. The ovarian tissue was carefully sectioned using a rotary microtome, stained and the histological architecture examined.

### **2.1 Statistical Analysis**

The data obtained from the study is expressed as mean  $\pm$  Standard error of mean (SEM). The significant differences among the groups were assessed using one-way analysis of variance (ANOVA) followed by Turkey's *post hoc* test. *P* values < 0.05 was considered statistically significant. All data were analyzed using IBM SPSS Verksion 23.0

## **3. RESULTS**

**Table 2: Phytochemical screening results for hydromethanolic fruit pulp extract of *A. garckeana***

S/N	Phytoconstituents	Test	Inferences
1	Alkaloids	Dragendorff test	
2	Cardiac Glycosides	Keller-Kiliani test	+
3	Saponins	Frothing test	+
4	Phenolic compounds	Lead acetate test	+
5	Tannins	Ferric Chloride test	+
6	Steroids	Salkowski test	+
7	Carbohydrates	Molisch test	+
8	Flavonoids	Shinoda test	+
9	Terpenoids	Liebermann Burchard test	+
10	Antraquinones	Bontragers test	

Keys: + PRESENT

### 3.2: Serum estradiol and progesterone hormone levels in female Wistar rats treated with hydromethanolic fruit pulp extract of *Azanza garckeana*.

There was no significant difference noted in the level of estradiol hormone between the three doses of the extract. However, a significant increase ( $p = 0.000$ ) was observed in the groups administered with the standard drug (positive control), 250mg/kg extract, 500mg/kg extract and 1000mg/kg extract when compared to the control group that received distilled water. Although the rise in the level of estrogen at the higher dose of 1000mg/kg is not up to that of the lower doses of 250mg/kg and 500mg/kg of the extract (table 3). The level of estradiol in the groups treated with the extract was not significantly different from that of the positive control group (table 3).

There was no significant difference in progesterone hormone levels in the groups administered 250mg/kg and 1000mg/kg of the extract when compared to both positive and negative control groups. However, there was a significant increase  $p = 0.000$  in the group administered 500mg/kg extract when compared to the control group and the positive control as shown in table 3.


Table 3: Effect of hydromethanolic fruit pulp extract of *Azanza garckeana* on the serum level of estradiol and progesterone hormone in female Wistar rats

Groups	Estradiol level	Progesterone level
Negative control	41.80±6.61 <sup>a</sup>	8.92±0.68 <sup>a</sup>
Positive control	108.40±3.91 <sup>b</sup>	7.86±0.93 <sup>a</sup>
250mg/kg Extract	104.40±2.30 <sup>b</sup>	7.54±0.34 <sup>a</sup>
500mg/kg Extract	112.40±4.72 <sup>b</sup>	11.50±0.39 <sup>b</sup>
1000mg/kg Extract	70.00±1.58 <sup>b</sup>	6.86±0.38 <sup>a</sup>

Data represented mean ± SEM, values with different super superscript letters <sup>a,b</sup> =  $p < 0.05$  show the level of significance between the groups. Positive control= Combined estrogen and progesterone (Estradiol valerate 100 µg and progesterone 50 µg).

### 3.3: Effect of hydromethanolic fruit pulp extract of *Azanza garckeana* on the proceptive (soliciting) behavior of female Wistar rats

A significant ( $p = 0.000$ ) increase in the proceptive behavior was observed in the groups administered the standard drug, 250 mg/kg extract, 500 mg/kg extract and 1000 mg/kg dose of extract when compared to the negative control group that received distilled water (figure 1). The level of proceptivity amongst all the treated groups including the positive control was not statistically different. However, administration of 250mg/kg extract resulted in a higher level of proceptive behavior when compared to the positive control that received the standard drug as show in figure 1.




**Figure 1: Effect of hydromethanolic fruit pulp extract of *A. garckeana* on proceptive behavior of female Wistar rats.**

Data represented mean  $\pm$  SEM, values with error bars having different super superscript letters a,b =  $p < 0.05$  are significantly different. Combined estrogen and progesterone (Estradiol valerate 100  $\mu$ g and progesterone 50  $\mu$ g).

### **3.4 Effect of hydromethanolic fruit pulp extract of *Azanza garckeana* on the receptivity of female Wistar rats using the lordosis quotient as the measure of evaluation.**

A significant ( $p = 0.000$ ) increase in the receptivity was observed in the group administered 1000mg/kg dose of *A. garckeana* when compared to the normal control, positive control and 250mg/kg *A. garckeana* extract treated group respectively (figure 2). A significant ( $p = 0.000$ ) increase in the receptivity was also observed in the 250mg/kg and 500mg/kg extract treated group when compared with the positive and normal control (figure 2).



**Figure 2:** Effect of hydromethanolic fruit pulp extract of *A. garckeana* on receptive behavior of female Wistar rats. Data represented as mean  $\pm$  SEM, values with error bars having different super superscript letters a,b,c,d =  $p < 0.05$  are significantly different.

### 3.5 Effect of hydromethanolic fruit pulp extract of *Azanza garckeana* on the time spent in the male chambers by the female Wistar rats (proximity time).

The result obtained showed an increase in time spent in the male chamber in the groups administered the standard drug, 250 mg/kg, 500 mg/kg and 1000 mg/kg dose of extract when compared to the negative (normal) control group that received distilled water as shown in table 4. The time spent increased significantly in the 250 mg/kg and 1000 mg/kg dose of the extract when compared to positive control and normal control, with the increase at the 1000 mg/kg dose significantly higher than the dose of 250 mg/kg of the extract (Table 4).





**Table 4: Effect of hydromethanolic fruit pulp extract of *Azanza garckeana* on the time spent in the male chambers by the female Wistar rats**


Groups	Time spent in male chamber (Proximity time)
Negative control	171.60±4.72 <sup>a</sup>
Positive control	271.80±41.35 <sup>b</sup>
250mg/kg Extract	318.80±10.52 <sup>c</sup>
500mg/kg Extract	202.40±4.39 <sup>a</sup>
1000mg/kg Extract	357.60±6.58 <sup>d</sup>

Data represented mean ± SEM, values with different super superscript letters a,b,c,d =  $p < 0.05$  show the level of significance between the groups. Positive control= Combined estrogen and progesterone (Estradiol valerate 100µg and progesterone 50µg).

### **3.6 Effect of hydromethanolic fruit pulp extract of *Azanza garckeana* on histo-architecture of the ovary in female Wistar rats**

Plate I showed slight hyperplasia of the inflammatory cells (IH) i.e cumulus oophorus, 65% primary follicle (PF), 20% secondary follicle (SF) and 15% corpus luteum (CL) in the negative control group, Plate II showed slight IH, 40% PF, 35% SF and 15% CL in the positive control group, the group administered with 250 mg/kg dose of extract showed no hyperplasia of the inflammatory cells, 50% PF, 30% SF and 20% CL as shown on plate III, the 500mg/kg dose of extract group showed normal ovary with 60% PF, 25% SF and 25% CL on plate IV. The 1000mg/kg dose of extract group result showed a normal ovary with 55% PF, 35% SF and 10% CL as shown on plate V. The percentages signify the number of a type of cell compared to the total number of all the cells present in the ovary.





Photomicrograph sections of the ovary using H & E  
X250 magnification

Plate I: Normal control

Plate II: Positive control (combined estrogen and  
progesterone)

Plate III: 250mg/kg extract treated group

Plate IV: 500mg/kg extract treated group

Plate V: 1000mg/kg extract treated group

IH: hyperplasia of the inflammatory cells. SF: Secondary  
follicle. PF: primary follicle. CL: corpus luteum

#### 4. DISCUSSION

The phytochemical analysis of the hydromethanolic fruit pulp extract of *A. garckeana* revealed the presence of cardiac glycosides, saponins, phenolic compounds, tannins, steroids, carbohydrates, flavonoids and terpenoids. Similar result to this was reported by Ochokwu *et al.*, [18] but anthraquinones and alkaloids were not detected in this study as in contrast with the stated research.

The results on the effect of *A. garckenana* fruit pulp on the level of estradiol obtained from this study revealed that the level of estradiol in the groups treated with the extract was not significantly different from that of the positive control group, however it showed a significant increase in the level of estradiol in the three doses of the extract tested when compared to normal control. The significant increase in estradiol may be as a result of increase in gonadotropin released by direct stimulation of accessory sex organs by the male Wistar rats. Estrogen is known to increase sexual desire [25], and it also acts in a feedback mechanism, influencing the production of follicle stimulating hormones (FSH) from the pituitary gland. FSH in turn promotes the development of the immature ovarian follicles, which increases the production of estrogen from the ovary [26]. This result agrees with the work of Mishra *et al.* [27], who reported the estrogenic activity of aphrodisiac plant *Bougainvillea spectables* leaves being connected to folliculogenesis. Increase in estradiol levels could also be related to the saponin present in the extract which is believed to be an estrogen precursor [28].

The results on the effect of *A. garckenana* fruit pulp on the level of progesterone obtained from this study revealed a decrease in the level of progesterone in the groups administered 250mg/kg and 1000mg/kg doses of the extract compared to the normal control while there was a significant increase at the dose of 500mg/kg as compared to control, making the 500mg/kg dose more potent for increasing the level of progesterone hormone. However, the decrease at 250mg/kg and 1000mg/kg could be as a result of inhibition of luteinizing hormone of the anterior pituitary by the high level of estrogen in the extract treated groups. This finding agrees with the report of Yakubu *et al.* [26], who observed that extract of *C. aconitifolius* caused a decrease in serum progesterone level of female rats; this could be due to the flavanoid content present in both extracts. The increase in progesterone level observed at 500mg/kg at estrus phase could be due to potent positive feedback mechanism on the pituitary to release gonadotropic hormones which stimulate granulosa cells to increase production of progesterone. It could also be as a result of an increase in follicular numbers observed at that dose from the histological study in this work.

The results on the effect of *A. garckeana* fruit pulp extract on proceptive (paracopulatory) behavior shows a significant increase in the level of proceptive behavior in the positive control, 250mg/kg, 500mg/kg and

1000mg/kg treated group when compared to the normal control group. The level of proceptive behavior noted in the three extract treated groups as well as the positive control was not significantly different. Generally, elevated sex hormone level enhances sexual behavior in humans [29]. This finding may be as a result of the increase in the level of estrogen in the groups treated with the extract as compared to control obtained in this study. This in turn may be due to the steroidal saponin content of *A. garckeana*, the regulation of female sexual motivation in rats by ovarian hormones has been demonstrated previously in numerous experiments [29]. This result is in agreement with the study carried out by Khadabadi and Bhajipale [30] on the effect of *Abutilon indicum* extract on female libido in rats who reported a significant increase in proceptive behavior after 11 and 21 days pretreatment with 200 and 400mg/kg of the extract. The result is also in agreement with Atuadu *et al.* [31] on the effect of ethanolic seed extract of *citrullus lanatus* on hypothalamic-pituitary-gonadal (hpg) axis and sexual behavior of female Wistar rats who reported a significant increase in the proceptive behavior the female Wistar rats after week 1 and 2 treatment with high dose of the seed.

The results obtained from this study on the effect of *A. garckeana* extract on the female rat receptivity revealed a significant dose-dependent increase in receptivity in the three doses of the extract when compared to both negative and positive control groups, indicating that 1000mg/kg dose of the extract resulted in the highest level of sexual receptivity. This findings are similar with the work of Nwafor *et al.* [32] who reported that lordosis quotient significantly increased with a corresponding decrease in lordosis latency which might be as a result of the estrogenic property or presence of phytochemical constituents (flavonoids and saponin) in the methanolic extract of *smilax kraussiana*. Contrary-wise the study reported by Tijjani *et al.* [33] on the aphrodisiac effect of aqueous stem bark extract of *ficus sycomorus* on female Wistar rats revealed that lordosis quotient did not significantly change across the doses of the extract tested. The alternating hormonal levels gotten from the study could be the reason for the non-significant change in the lordosis quotient Tijjani *et al.* [33] reported in their research. The discrepancy could also be due to the different extracts used.

The time spent in the male chamber by the female rat increased significantly in the groups administered with 250 mg/kg and 1000 mg/kg doses of the extract when compared to normal and positive control. This

is as a result of the high level of the sex steroid hormone estrogen which increased the ability of the female rat to stay in close proximity to the male rats and subsequently promote the copulatory act. The increase in the time spent at 500mg/kg dose of the extract when compared to normal control was not statistically significant, this could be related to the increased progesterone hormone level noted in this group. This result is in-line with the work on the aphrodisiac effect of aqueous stem bark extract of *ficus sycomorus* on female Wistar rats by [33], they reported an increase in mean time spent by the female animal in the male compartment though not significant. This result is also in agreement with the study carried out by Khadabadi and Bhajipale [30], on the effect of *Abutilon indicum* extract on female libido in rats who reported that pretreatment with AI 100, 200 and 400 significantly improved the core proximity time for male target as compared to control rats.

The presence of flavonoid and saponin in the fruit extract could be the reason for the high aphrodisiac activity i.e increased sexual behavior and sex hormone levels observed in the extract. Steroidal saponins have been recorded to play a role in enhancing sexual behavior by either binding to hormone receptors, which may result in conformational change that will enhance the physiological function of the hormones or bind to enzymes that are involved in the synthesis of such hormones and thus enhance its production [34]. Saponins found primarily in the leaf *Tribulis terrestris* L. have been used as an aphrodisiac agent in rats [35]. These phytochemical contents of *A. garckeana* may justify the claim for their remarkable aphrodisiac activities, and its usage in treatment of diseases.

The results from the histological studies on the effect of hydromethanolic fruit pulp extract of *Azanza garckeana* on the histo-architecture of the ovary showed an increase in the proliferation and growth of the ovarian follicle from primary follicle to secondary follicle then subsequently to corpus luteum (shown by the percentages of each type of follicle) in all three doses of the extract when compared to control. This corroborates with the significant increase in estrogen level in the extract treated groups when compared to control. Estrogen acts in a feedback mechanism where it influences the production of follicle stimulating hormones (FSH) from the pituitary gland. FSH in turn promotes the development of the immature ovarian follicles which increases the production of estrogen from the ovary [26]. This result is similar to the result obtained by Tijjani *et al.* [33] on the aphrodisiac effect of aqueous stem bark extract of

*ficus sycomorus* on female Wistar rats, they reported an increase in the number of ovarian follicles in the high dose seed extract treated group when compared to control.

## CONCLUSION

This study has demonstrated that hydromethanolic fruit pulp extract of *A. garckeana* possesses significant aphrodisiac potentials in female Wistar rats evidenced by the increased sexual behavior, sex hormone levels, histological changes in the ovaries, and increased proximity time of the extract treated animals.

## ETHICAL APPROVAL

Ethical approval was sought and gotten from A.B.U committee on animal use and care with the approval number ABUCAUC/2021/121. The study was carried out in accordance with the principles of laboratory animal care and standard experimental procedure.

## REFERENCES

1. World Health Organization. Defining sexual health. 2020. Retrieved from [https://www.who.int/health-topics/sexual-health#tab=tab\\_1](https://www.who.int/health-topics/sexual-health#tab=tab_1).
2. World Health Organization. Defining Sexual Health: Report of a Technical Consultation on Sexual Health. 2006.
3. World Health Organization. working document on sexual and reproductive health. 2002.
4. Baulieu EE. (1996). Dehydroepiandrosterone (DHEA): a fountain of youth? Journal of Clinical Endocrinology and Metabolism, 81: 3147-3151. DOI: [10.1210/jcem.81.9.8784058](https://doi.org/10.1210/jcem.81.9.8784058)
5. NIH Consensus Conference. Impotence: NIH Consensus Development Panel on Impotence. Journal of the American Medical Association. 1993; 270:83–90. <https://doi.org/10.1001/jama.1993.03510010089036>
6. Laumann EO, Paik A and Rosen RC. Sexual dysfunction in the United States: prevalence and predictors. Journal of the American Medical Association. 1999; 281(5): [37-44](https://doi.org/10.1001/jama.281.6.537). [doi:10.1001/jama.281.6.537](https://doi.org/10.1001/jama.281.6.537)

7. Franjić S. Female Sexual Dysfunction. International Journal of Reproductive Medicine and Sexual Health. 2019;p: 24-29. DOI:[10.36811/ijrmsh.2019.110003](https://doi.org/10.36811/ijrmsh.2019.110003)
8. Gundidza GM, Mmbengwa VM, Magwa ML, Ramalivhana NJ, Mukweroh NT, Ndaradzi W and Samie A. Aphrodisiac properties of some Zimbabwean medicinal plants formulation. African Journal of Biotechnology. 2009;8(22):6402-6407. DOI: [10.5897/AJB09.430](https://doi.org/10.5897/AJB09.430)
9. Smith GE. Health and disease in ancient egypt. Papyrus Ebers Oxford University press. Chicago: Ares Publishers. 1974;Pp: 857-877. DOI:[10.30861/9781407305004](https://doi.org/10.30861/9781407305004)
10. Alok S, Retendra, K and Roman-deep S. Nature Aphrodisiac review of Current Scientific Literature. International Journal of Recent Advances in pharmaceutical Research. 2013;3(2):1-25. DOI:[10.4314/njbas.v26i1.8](https://doi.org/10.4314/njbas.v26i1.8)
11. Watcho P, Fabien Z, Telesphore BN, Silvere N, Phelix BT, Pierre K, Etienne T and Albert K. Effects of the aqueous and hexane extracts of *Mondia whitei* on the sexual behavior and some fertility parameters of sexually inexperienced male rats. African Journal of Traditional Complementary and Alternative Medicine. 2007;4(1):37-46. doi: [10.4314/ajtcam.v4i1.31190](https://doi.org/10.4314/ajtcam.v4i1.31190)
12. Orwa C, Mutua A, Kindt R, Jamnadass R and Anthony S. *Azanza garckeana* agroforestry database: A tree reference and selection guide version 4.0. 2009. Available: <http://www.worldagro-forestry.org/Af/tree>.
13. Okafor JC. Selection and improvement of indigenous tropical trees. Journal of Tropical Resources. 1985;2: 87-95.
14. Adamu HM, Ushie OA, Lawal DS and Oga IA. Phytochemical screening of fruit of *Azanza garckeana* and root of *Acacia macrothyrsa*. International Journal of Traditional and Natural Medicines. 2013;3(1): 19-25. DOI:[10.30574/gscbps.2020.11.3.0179](https://doi.org/10.30574/gscbps.2020.11.3.0179)
15. Masila VM, Midiwo, JO, Zhang J, Gisacho BM, Munayi R, Omosa LK, Wiggers FT, Jacob MR, Walker LA and Muhammad I. Antivancomycin-resistant *Enterococcus faecium* and *E. faecalis* activities of gossypol and derivatives from *Thespesia garckeana*. Natural Product Communications. 2015;10(4) 613-616. DOI:[10.1016/j.jep.2015.12.028](https://doi.org/10.1016/j.jep.2015.12.028)
16. Maroyi A. An ethnobotanical survey of medicinal plants used by the people in Nhema communal area, Zimbabwe. Journal of Ethnopharmacology. 2011;136: 347–354. DOI: [10.1016/j.jep.2011.05.003](https://doi.org/10.1016/j.jep.2011.05.003)

17. Mshelia EH, Watirahyel EM, Maigari AU, Yohanna, C. and Ismail F. Cytotoxicity and antioxidant activity of stem bark extracts of *Azanza garckeana* (kola of Tula). European Journal of Pure and Applied Chemistry. 2016;3(2): 1-11. DOI: <https://doi.org/10.53555/as.v7i6.4356>
18. Ochokwu IJ, Dasuki A. and Oshoke JO. *Azanza garckeana* (Goron Tula) as an edible indigenous fruit in North Eastern Part of Nigeria. Journal of Biology, Agriculture and Healthcare. 2015;5(15): 26-31.
19. Dikko YJ, Khan ME, Tor-Anyiin TA, Anyam JV and Linus UA. In vitro Antimicrobial Activity of Fruit Pulp Extracts of *Azanza garckeana* (F. Hoffm.) Exell and Hillc. And Isolation of One of its Active Principles, Betulinic Acid. British Journal of Pharmaceutical Research. 2016;14(1-10). DOI:[10.9734/BJPR/2016/30152](https://doi.org/10.9734/BJPR/2016/30152)
20. Long JA and Evans HM. The estrous cycle in rats and its associated phenomena. Memories of University of California. 1992;6:1-148.
21. Mokhtari M. and Zanboori M. The effects of lead acetate on sexual behavior and level of testosterone in adult male rats. International Journal of Fertility and Sterility. 2011;5(1):13-20). PMID: [PMC4040238](https://pubmed.ncbi.nlm.nih.gov/24040238/)
22. Tennent BJ, Smith ER and Davidson JM. The effects of estrogen and progesterone on female rat proceptive behavior. Hormones and behavior. 1980;14 65-75. [doi: 10.1016/0018-506x\(80\)90016-1](https://doi.org/10.1016/0018-506x(80)90016-1).
23. Adebayo EM. Effect oral administration of decoction on serum levels of levels of luteinizing hormone, progesterone and estrogen in female Dutch-white rabbits. Research Journal of Medicinal Plants. 2015;9:141-145. [doi=rjmp.2015.141.145](https://doi.org/10.141.145)
24. Tietz NW. Clinical guide to laboratory tests, 3rd edition WB Saunders, Philadelphia. 1995;1-997. <https://doi.org/10.1111/j.1537-2995.1995.tb03571.x>
25. Thornton JE. and Finn PD. Estrus in: Encyclopedia of Reproduction. Nigerian Journal of Pharmaceutical and Applied Science Research. 1999;1(2):1-14.
26. Yakubu MT, Akanji MA, Oladiji AT et al. Effect of *Cnidioscolousacontifolious* (Miller) I.M. Johnston leaf extract on reproductive hormones of female rats. Iranian Journal or Reproductive Medicine. 2008;6(3):149-155.
27. Mishra N, Joshi S, Tandon VL and Munjal A. Evaluation of Anti-fertility potential of Aqueous Extract of *Bougainvillea spectabilis* leaves in Swiss Albino mice. International Journal of Pharmaceutical Sciences and Drug Research. 2009;1(1):19-23. DOI <https://doi.org/10.25004/IJPSDR.2009.010105>

28. Gauthamen K, Adaikan PG, Prasan RNV and Ngaji SC. Proevectile Pharmacological effects of *Tribulus terrestris* extract on the rabbit Corpus cervanosum. *Annals of the Academy of Medicine Singapore*. 2000;29(1):22-26.
29. Wilson CA and Hunters AJ. Progesterone stimulates sexual behaviour in female rats by increasing 5-HT activity on 5-HT<sub>2</sub> receptors. *Brain*. 1985;p: 33. [https://doi.org/10.1016/0006-8993\(85\)91575-6](https://doi.org/10.1016/0006-8993(85)91575-6)
30. Khadabadi SS, Bhajipale NS. Effect of *Abutilon indicum* extract on Female Libido in Rats. *International Journal of PharmTech Research*, 2011;3(3):1652-1659.
31. Atuadu VO, Anyanwu EG and Ikele I. Effect of ethanolic seed extract of *Citrullus lanatus* on hypothalamic-pituitary-gonadal (hpg) axis and sexual behavior of female Wistar rats. *International Journal of Research and Analytical Reviews*. 2018;5(3): 588-601. DOI: <https://doi.org/10.22271/phyto.2021.v10.i4a.14182>
32. Nwafor PA, Idiong OJ and Davies K. Contraceptive and Sexual behavioral effects of methanol extract of *Smilax kraussiana* root in rodent. *African Journal of Pharmacology and Therapeutics*. 2013;2(3):94-100
33. Tijjani U, Mabrouk MI, Yarube IU, Muhammad M. Aphrodisiac effect of aqueous stem bark extract of *Ficus Sycomorus* on female Wistar rats. *Nigerian Journal of Basic and Applied Science*. 2018;26(1):70-79. DOI: [10.4314/njbas.v26i1.8](https://doi.org/10.4314/njbas.v26i1.8)
34. Gauthaman K. and Adaikan PG. The hormonal effect of *Tribulus Terrestris* and role its role in the management of erectile dysfunction- an evaluation using primates rabbit and rats. *Phytomedicine*, 2008;15(1):44- 54. doi: 10.1016/j.phymed.2007.11.011.
35. Goel B and Maurya NK. Aphrodisiac Herbal therapy for Erectile Dysfunction. *Archives of Pharmacy Practice*. 2020;11(1):1-6.