

## **Review Article**

**A review of COVID-19's role in pregnancy and its association with the risk of miscarriage in pregnant women**

**Comment [U1]:** The topic would be better as a review of COVID-19's impact on pregnancy. The miscarriage is also part of the impact. So no need for mentioning miscarriage in the title.

**Comment [U2]:** No need for mentioning pregnant women again. Everyone know that pregnancy is found only on pregnant women.

### **Abstract**

**Background and Aims:** Since the new Coronavirus started spreading in December 2019, the virus has had enormous consequences on people's health and way of life in every nation on earth has been touched. A pregnant lady is more susceptible to infection than a healthy one because she is immunocompromised throughout pregnancy. The purpose of this present paper was to review relevant published literature and also aims to determine whether COVID-19 infection increases the risk of miscarriage in pregnant women.

**Comment [U3]:** The content of the review is not only on miscarriage.

**Method:** In this narrative review all publications published in the databases PubMed, Scopus, Embase, Science Direct, and Web of Science from December 2019 to August 2022 by conducting a complete search with the required keywords, including COVID-19, coronavirus illness 2019, SARS-CoV-2, and miscarriage, abortion, or pregnancy outcomes. There are few accessible data on the effects of COVID-19 on pregnancy. Every day, new information emerges on the management of pregnancy during the COVID-19 era.

**Comment [U4]:** Is there no inclusion and exclusion criteria?

**Result:** In this review, we assess the most recent research regarding the effects of SARS-CoV-2 infection during pregnancy. COVID-19 can cause fetal distress, miscarriage, respiratory distress, and preterm delivery in pregnant women, according to the reviewed literature.

**Comment [U5]:** Why no results in the main work?

**Comment [U6]:** But you said all under method, and here you said most recent. Is this not contradiction?

**Comment [U7]:** That is why you should remove miscarriage from the title since it is not only miscarriage

**Conclusion:** Due to the absence of adequate data regarding the effects of COVID-19 on pregnancy, it is required to follow suspected pregnant women prior to and after delivery. For verified cases, both the mother and newborn should be thoroughly monitored.

**Keywords:** COVID-19, Pregnancy, miscarriages,

### 1. Introduction:

In December 2019, an unidentified pneumonia outbreak was reported in Wuhan, Hubei Province, China<sup>1</sup>. Those who have COVID-19 typically experience viral pneumonia, with the most common symptoms being fever, cough, sore throat, myalgia, and weariness<sup>2</sup>. COVID-19 has afflicted over 170 million individuals globally, resulting in over 3.5 million deaths by the end of May 2021. Several COVID-19 vaccinations were introduced to prevent the spread of COVID-19 during the pandemic, and treatment regimens were quickly established with the introduction of new pharmaceuticals, including anti-SARS-CoV-2 monoclonal antibodies and antiviral treatments<sup>3</sup>. Coronaviruses are enclosed, non-segmented, single-stranded ribonucleic acid (RNA) viruses that cause illnesses ranging from simple colds to serious prenatal disorders. SARS-CoV, which causes severe acute respiratory syndrome (SARS), and MERS-CoV, which causes Middle East respiratory syndrome (MERS), are the two most well-known fetal viruses<sup>4</sup>. The World Health Organization declared a pandemic on March 12, 2020, due to the global spread of SARS-CoV-2 and thousands of deaths caused by coronavirus illness<sup>1</sup>.

Due to physiological changes in the immunological and cardiopulmonary systems during pregnancy, pregnant women may be more likely to experience more severe symptoms after contracting respiratory viruses. MERS-CoV and SARS-CoV have also been linked to more

**Comment [U8]:** The conclusion is not tallying with the title and even with the results. Your study is not on prevention of impacts of covid 19 on pregnancy. Why are you concluding the following suspected pregnant women. Moreover, the study is on confirmed cases of covid 19 in pregnancy not suspected cases. In conclusion you should state the impacts of covid 19 on pregnancy in a form of super-summary. Make conclusion based on what is in your results. You said COVID-19 can cause fetal distress, miscarriage, respiratory distress, and preterm delivery in pregnant women. Your conclusion should be based on this findings.

**Comment [U9]:** This should be recommendations. Based on your findings you are recommending that ... in order to prevent the fetal distress, miscarriage, respiratory distress, and preterm delivery in pregnant women

**Comment [U10]:** There is no connection between first and second sentences. The first sentence has not indicated that the unidentified pneumonia outbreak was covid-19.

**Comment [U11]:** Source of this statement is necessary

**Comment [U12]:** This supposed to come before talking about vaccine and treatment of covid 19. Vaccine and treatment should come after description of the diseases.

**Comment [U13]:** This should form part of the first sentences

**Comment [U14]:** This should be the beginning of the paragraph

**Comment [U15]:** Citation is necessary here

serious pregnancy problems and higher case fatality rates. On the effect of COVID-19 during pregnancy, there are, however, few reports. Because of this, it is uncertain how COVID-19 treatment may affect fetal and neonatal outcomes, and research are urgently needed to address this issue <sup>5</sup>.

**Comment [U16]:** Citation is necessary here

**Comment [U17]:** It is better to state it as this: However, there are few reports on the effect of covid-19 on pregnancy.

**Comment [U18]:** This statement is irrelevant because the review is not on covid-19 treatment, and you are not doing a research to address the issue of covid-19 treatment in pregnancy.

Pregnancy is thought to be a risk factor for severe sickness from COVID-19, which is caused by SARS-CoV-2 infection. Furthermore, pregnant women with COVID-19 may be at a greater risk of other negative outcomes, such as preterm delivery and abortion. An increasing body of evidence suggest linking of COVID-19 to abortion <sup>6</sup>. The immune system, respiratory system, cardiovascular system, and coagulation are all significantly impacted by physiological changes during pregnancy. These could accelerate or halt the progression of COVID-19 disease <sup>7</sup>. SARS-CoV-2 primarily spreads through the respiratory route, suspended on droplets or, less frequently, aerosols <sup>8</sup>.

**Comment [U19]:** Please give citation here?

**Comment [U20]:** But you talk about physiological changes in pregnancy above. Why can't this sentence be together with that one above. This shows lack of logic in your writing. You write with flight of thought.

The chance of developing a serious COVID-19-related illness is increased in pregnant women. Compared to non-pregnant women, pregnant women are more likely to be referred to an intensive care unit (ICU) <sup>9</sup>. By conducting a systematic review that evaluated the prevalence of miscarriages in pregnant women with COVID-19, this paper is aimed to review potential causes of COVID-19 that may cause pregnancy losses based on the findings of human studies during the pregnancy, as well as the current evidence of the COVID-19-impact on pregnancy losses.

**Comment [U21]:** The causes of covid-19 will give another meaning different from intended meaning. The sentence need to be restated correctly.

**1.1 Impact of Covid-19 on pregnancy:** The risk of COVID-19 to expectant mothers is minimal generally. However, those who are pregnant or have recently given birth run a higher chance of developing a serious -illness from COVID-19. If the pregnant women have a serious sickness, they may need to be hospitalized and get critical care. Additionally, COVID-19-positive pregnant women are more likely to give birth before the 37th week of pregnancy (premature

**Comment [U22]:** Look at the first sentence of above paragraph where you said "The chance of developing a serious COVID-19-related illness is increased in pregnant women." Is this not a contradiction with this statement? Moreover, where is the citation for this statement?

**Comment [U23]:** Even this sentence contradict the previous sentence

birth). Furthermore, COVID-19-positive pregnant women may be more likely to experience issues including stillbirth and miscarriage<sup>10</sup>. Some of the main issues include whether pregnant women are more susceptible to COVID-19, whether infected pregnant women have worse outcomes in terms of morbidity and/or death compared to non-pregnant women of the same ages, whether infected pregnant women have more severe disease, or whether they have worse maternal-fetal outcomes compared with uninfected pregnant women of comparable ages<sup>11</sup>.

**Comment [U24]:** That means not assurance. But in the first sentence you mentioned that "The risk of COVID-19 to expectant mothers is minimal generally."

Compared to their contemporaries who are not pregnant, pregnant women with COVID-19 may be at higher risk for more serious sickness, according to the data that is currently available. These findings specifically show an increased likelihood of mechanical breathing and ICU hospitalizations. However, Table -1 shows some maternal symptoms which was caused in pregnant women during their gestational weeks. There hasn't been any evidence of a rise in mortality rates. Pregnant women with comorbidities like obesity and gestational diabetes may be at an even higher risk for serious illnesses than the general population with comparable comorbidities. It's significant that analyses performed thus far have been constrained by a significant quantity of missing data and the fact that many publications lack an adequate comparative control group that isn't pregnant<sup>12</sup>.

**Comment [U25]:** This word is not appropriate here

**1.2 COVID-19 infection and miscarriages:** A miscarriage occurs when a pregnancy ends on its own before the 20th week. The majority of miscarriages happen prior to the 12th week of pregnancy<sup>23</sup>. Research has revealed that the COVID-19 virus adheres to a number of bodily organs. In early placental tissue, SARS-Cov-ACE2 protein is expressed, which makes the early pregnancy vulnerable to the effects of COVID-infection<sup>24</sup>.

During the periconceptional period, miscarriage and unsuccessful embryo implantation are more likely in women with COVID-19. Unbalanced immune hormones causes "cytokine storm" results a hypercoagulable state that is harmful to normal fetal development <sup>25</sup>. In fact, anomalies in the mother's immune system during the peri-implantation and early stages of pregnancy have been associated to repeated implantation loss and miscarriage <sup>26</sup>.

## 2. Method

**2.1 Study Design:** This review intended to comprehensively analyze the risk of miscarriage in pregnant women who had COVID-19 during their pregnancy. This review period began in June 2022, with additional updates in July and August. The popular databases like PubMed, Scopus, Embase, Science Direct, and Web of Science during December 2019 to August 2022 were investigated using the following search terms: COVID-19, coronavirus illness 2019, SARS-CoV-2, and miscarriage, abortion, or pregnancy outcomes. The search was limited to English-language academic publications. EndNote X9 (Clarivate Analytics US LLC, Philadelphia, USA) was also utilized as reference management software. Additionally, a manual search of the references included in the chosen papers and reviews was done.

**2.2 Data extraction and quality assessment:** From the publications that were ultimately chosen, information about the place of origin, the number of pregnant patients with confirmed COVID-19, clinical symptoms, laboratory results, outcomes, and key findings were retrieved.

## 3. Discussion:

Pregnancy is a distinct immunological state in which complex physiological processes occur at the maternal-fetal contact. When this delicate equilibrium is troubled by infections, the system might collapse. Few data exist about the immunological response to SARS-CoV-2 infection

**Comment [U26]:** Please read PRISMA and get details information on method, results and discussion in systematic review

**Comment [U27]:** Where is the results? Including the flow diagram of the selected articles. Usually results contain search results, study range and characteristics, study quality etc.

**Comment [U28]:** You need to arrange your discussion logically based on categories of the findings such as fetal distress, miscarriage, respiratory distress, and preterm delivery. Also, discuss the limitations of the study, strengths and weaknesses of the review and implications for current practice.

during pregnancy, highlighting the significance of this review. Regarding the specific immunological features of COVID-19 in pregnant women, there is still no solid publication based on currently available scientific information. Regardless of study design, this review chose all studies pertaining to COVID-19, and pregnancy. All included papers are narrative reviews devoid of methodological and technical information regarding the selection of data.

Case reports and case studies from the pandemic have shown that moms who tested positive for SARS-CoV-2 have a higher risk of abortion. The majority of these research discussed how the virus affects the pregnant women and causes serious impact which slows down fetal growth and may cause abortion. A case report done by Fang, Nz in 2020 in USA resulted one miscarriages of a pregnant women with Covid 19<sup>27</sup>. In Malaysia another case report was done by Wong, T.C. and miscarriages occurred in two of the cases during the first trimester of pregnancy where both pregnant women were infected by SARS-Cov 2<sup>28</sup>. A study was done in 727 women made their first prenatal visit between June 29 and September 30, 2019, and 67 of them (9.2%) experienced a miscarriage. 542 women had their first prenatal appointment during the same time period in 2020, and 64 (11.8%) of them miscarried before 12 weeks' gestation. Prior to admission, SARS-CoV-2 was tested on all women who miscarried in 2020, and 4.7 percent (3/64) of the tests were positive<sup>29</sup>.

Comment [U29]: This is informal word

According to the results of the current study, further investigation into the potential for vertical transmission of COVID-19 from mother to fetus is considered essential. Results from studies indicate that infected or suspected mothers should be closely monitored before and after giving birth. The recommendation is for women who believe they may have been exposed to COVID-19 to hold off on nursing until they have received formal clearance<sup>30</sup>. A study done by Chen et al., 2020 found that COVID-19 can induce fetal discomfort in pregnant women but does not

infect infants<sup>31</sup>. A similar study was done by in China. They have found that there is no evidence of intrauterine COVID-19 infection caused by vertical transfer to the fetus<sup>32</sup>. According to a study in Iran, women who are suspected or know for sure that they have COVID-19 should not breastfeed their babies for at least two weeks after they are born. If COVID-19 infection is confirmed while a woman is pregnant, both the mother and the fetus should be closely watched<sup>33</sup>. Another study in China suggest that There is no proof that COVID-19 causes fetal distress or infections in babies. COVID-19 can be mild or severe during pregnancy, and it can cause the baby to come early which may sometimes cause critical issues in newborn<sup>34</sup>. According to a 2020 Chinese study, infection with COVID-19 during pregnancy might result in adverse consequences such miscarriage, fetal development impairment, early delivery, and even maternal fatality<sup>35</sup>. But another study shows that there is no indication that pregnant women can transmit COVID-19 vertically<sup>36</sup>.

According to a research paper done in the United States in 2020, some newborns showed signs of fetal distress and preterm delivery. After birth, SARS-CoV-2 testing was negative for all kids born to pregnant women with COVID-19<sup>37</sup>. According to a study done in China in 2020, prenatal COVID-19 infection may have detrimental effects on newborns. These effects may include fetal discomfort, early labor, respiratory distress, thrombocytopenia followed by impaired liver function, and even death<sup>38</sup>. As per some studies neither the mother nor the infant have shown any signs of having contracted SARSCoV-2 during pregnancy<sup>39-41</sup>. According to research, being affected with SARS during the perinatal period is associated with a high prevalence of potentially harmful maternal and neonatal side effects, including disseminated intravascular coagulopathy, spontaneous abortion preterm birth, intrauterine growth retardation, neonatal, intubation and the need for the newborn to be admitted to a neonatal intensive care unit (ICU)

Comment [U30]: By who?

and organ failure<sup>42</sup>. Because of physiological, immune, and anatomical changes, pregnant women are more susceptible to respiratory diseases. Some studies point to a significant shift to a T-helper lymphocyte type 2 (Th2) immune response during pregnancy as a potential contributor to the severity of COVID-19 cases. Furthermore, the cytokine storm that occurs in severe cases causes an increased inflammatory state, which may worsen the clinical prognosis in this population. As a result, pregnant women may be a vulnerable group to COVID-19 infection, owing to an immune imbalance at the maternal-fetal interface<sup>43</sup>. Therefore, there is a hypothesis that the hormone level and immunological condition differ at various phases of pregnancy. In early pregnancy, for instance, the author suggests that the immune balance is still unstable, which can lead to a severe immune system disorder and internal environment imbalance in the event of a viral infection, resulting in abortion or abnormal fetal growth, similar to that observed in other respiratory infections. According to this author, as the pregnancy proceeds, the mother is constantly changing this immunological balance, with presumably decreasing severity<sup>44</sup>. Recent research indicates that in severe cases of COVID-19 infection, a cytokine-storm occurs, which is characterized by increased concentrations of plasma mediators such as IL-2, IL-7, IL-10, granulocyte-colony stimulating factor, and tumor necrosis factor alpha, most of which are associated with an inflammatory response<sup>45</sup>. Dashraath et al. hypothesized that the physiological change to a Th2-anti-inflammatory environment during pregnancy and other undisclosed immune adaptations may function as the major immune response to SARS-CoV-2, resulting in a less severe manifestation of COVID-19 in the pregnant women. These results are consistent with the scant literature that asserts that, despite being put in high-risk groups mostly due to preventive measures, cases of COVID-19 among pregnant women may not be more severe or have worse clinical outcomes than those in the general population. Due to the lack of relevant

data about the effects of COVID-19 on the immune response during pregnancy, a number of issues remain unresolved until the publication of adequate, case-based, evidence-based studies <sup>46</sup>.

According to Li et al., 2020 four of seven pregnant women with COVID-19 who presented in the first trimester experienced spontaneous miscarriages, while two had voluntary terminations. The sole newborn survivor was born at full term, and no birth defects were noted <sup>47</sup>. Yan et al. hypothesized that pregnant women with COVID-19 had a higher rate of maternal mortality, intubation, and ICU hospitalization than non-pregnant women with COVID-19, but that the virus was not transmitted to the newborn. In addition, more problems such as miscarriage, premature birth, and small-for-gestational-age newborns have been documented. One study, however, found no increase in the risk of spontaneous abortion or spontaneous preterm birth among pregnant COVID-19 carriers. Numerous systemic infections and inflammatory conditions are related with premature birth, and preterm births have been observed in women infected with COVID-19. However, the majority were recorded in China, which has a distinct medical system, and it is still unclear whether SARS-CoV-2 caused these premature births or whether they were iatrogenic as a result of maternal distress or other indicators <sup>48</sup>. In table 2, this review study attempted to compile data on the number of miscarriages caused by COVID-19 in various countries.

### Conclusion

In summary, one of the mainstays of the healthcare system is the provision of safe care to expectant mothers who have COVID 19 infection. The lack of information about the epidemic among pregnant women is one of the study's inadequacies. This may be connected to pregnant moms' anxiety about receiving routine prenatal treatment in clinics during the COVID-19 epidemic. Additionally, because the mother's consent was not obtained, restricted sample of the

**Comment [U31]:** Please make conclusion based on your title and results. See my comments on conclusion under abstract

aborted fetus or placenta and absence of pathology reports have kept the current statistics on these abortions to the cases reported. There is little information available from the pregnant women who had abortions during the pandemic. Further investigation is therefore required to determine the causes of pregnancy loss in COVID-19 infected mothers and whether there is any evidence that viral transmission to the fetus causes miscarriage due to the virus's direct impact on the fetal organs.

## 6. References

- [1] Ciotti M, Ciccozzi M, Terrinoni A, Jiang W-C, Wang C-B, Bernardini SJCrcls. The COVID-19 pandemic. 2020; **57**(6): 365-88.
- [2] Chan JF-W, Yuan S, Kok K-H, *et al.* A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. 2020; **395**(10223): 514-23.
- [3] Cavalcante MB, Cavalcante CTdMB, Cavalcante ANM, Sarno M, Barini R, Kwak-Kim JJJorI. COVID-19 and miscarriage: From immunopathological mechanisms to actual clinical evidence. 2021; **148**: 103382.
- [4] Lu R, Zhao X, Li J, *et al.* Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. 2020; **395**(10224): 565-74.
- [5] Schwartz DA, Graham ALJV. Potential maternal and infant outcomes from coronavirus 2019-nCoV (SARS-CoV-2) infecting pregnant women: lessons from SARS, MERS, and other human coronavirus infections. 2020; **12**(2): 194.
- [6] Kazemi SN, Hajikhani B, Didar H, *et al.* COVID-19 and cause of pregnancy loss during the pandemic: A systematic review. 2021; **16**(8): e0255994.

- [7] Wastnedge EA, Reynolds RM, Van Boeckel SR, *et al.* Pregnancy and COVID-19. 2021; **101**(1): 303-18.
- [8] Meyerowitz EA, Richterman A, Gandhi RT, Sax PEJAoim. Transmission of SARS-CoV-2: a review of viral, host, and environmental factors. 2021; **174**(1): 69-79.
- [9] Pinar MH, Gibbins K, He M, Kostadinov S, Silver RJF, pathology p. Early pregnancy losses: review of nomenclature, histopathology, and possible etiologies. 2018; **37**(3): 191-209.
- [10] Figueiro-Filho EA, Yudin M, Farine DJJopm. COVID-19 during pregnancy: an overview of maternal characteristics, clinical symptoms, maternal and neonatal outcomes of 10,996 cases described in 15 countries. 2020; **48**(9): 900-11.
- [11] Elsaddig M, Khalil AJBP, Obstetrics RC, Gynaecology. Effects of the COVID pandemic on pregnancy outcomes. 2021; **73**: 125-36.
- [12] Mahtab NT, Chowdhury TSJBMJ. Clinical outcomes of women with COVID 19 during pregnancy: an update. 2020: 93-102.
- [13] Breslin N, Baptiste C, Miller R, *et al.* Coronavirus disease 2019 in pregnancy: early lessons. *American journal of obstetrics & gynecology MFM*. 2020; **2**(2): 100111.
- [14] Noelle B, Cynthia G-B, Russell M, Rebecca M, Kyra B, Laurence R. COVID-19 infection among asymptomatic and symptomatic pregnant women: two weeks of confirmed presentations to an affiliated pair of New York City hospitals. *Am J Obstet Gynecol MFM*. 2020; **100118**.
- [15] Baergen RN, Heller DS. Placental pathology in Covid-19 positive mothers: preliminary findings. *Pediatric and Developmental Pathology*. 2020; **23**(3): 177-80.
- [16] Fan C, Lei D, Fang C, *et al.* Perinatal transmission of COVID-19 associated SARS-CoV-2: should we worry? *Clinical infectious diseases*. 2020.

- [17] Khan S, Jun L, Siddique R, *et al.* Association of COVID-19 with pregnancy outcomes in health-care workers and general women. *Clinical microbiology and infection*. 2020; **26**(6): 788-90.
- [18] Cao D, Yin H, Chen J, *et al.* Clinical analysis of ten pregnant women with COVID-19 in Wuhan, China: A retrospective study. *International Journal of Infectious Diseases*. 2020; **95**: 294-300.
- [19] Huang W, Zhao Z, He Z, *et al.* Unfavorable outcomes in pregnant patients with COVID-19. *Journal of Infection*. 2020; **81**(2): e99-e101.
- [20] Ferrazzi E, Frigerio L, Savasi V, *et al.* Vaginal delivery in SARS-CoV-2 infected pregnant women in Northern Italy: a retrospective analysis. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2020; **127**(9): 1116-21.
- [21] Govind A, Essien S, Karthikeyan A, *et al.* Re: novel coronavirus COVID-19 in late pregnancy: outcomes of first nine cases in an inner city London hospital. *European Journal of Obstetrics and Gynecology and Reproductive Biology*. 2020; **251**: 272-4.
- [22] Gagliardi L, Danieli R, Suriano G, *et al.* Universal severe acute respiratory syndrome coronavirus 2 testing of pregnant women admitted for delivery in 2 Italian regions. *American Journal of Obstetrics & Gynecology*. 2020; **223**(2): 291-2.
- [23] Mily SJ, Akter KM, Jabin N, *et al.* COVID-19 Infection in Pregnancy: A Review. 2022; **22**(5): 12-21.
- [24] Overtoom EM, Rosman AN, Zwart JJ, *et al.* SARS-CoV-2 infection in pregnancy during the first wave of COVID-19 in the Netherlands: a prospective nationwide population-based cohort study (NethOSS). 2022; **129**(1): 91-100.

- [25] Sills ES, Wood SHJIoM, Medicine C. An experimental model for Peri-conceptual COVID-19 pregnancy loss and proposed interventions to optimize outcomes. 2020; **9**(3): 180.
- [26] Yang KM, Ntrivalas E, Cho HJ, *et al.* Women with multiple implantation failures and recurrent pregnancy losses have increased peripheral blood T cell activation. 2010; **63**(5): 370-8.
- [27] Fang NZ, Castaño PM, Davis AJC. A hospital-based COVID-19 abortion case in the early phase of the pandemic. 2020; **102**(2): 137-8.
- [28] Wong TC, Lee ZY, Sia TL, Chang AK, Chua HHJScm. Miscarriage risk in COVID-19 infection. 2020; **2**(9): 1449-52.
- [29] Sacinti K, Kalafat E, Sukur Y, Koc A. Increased incidence of first-trimester miscarriage during the COVID-19 pandemic. *Ultrasound in Obstetrics & Gynecology*. 2021; **57**(6): 1013.
- [30] Panahi L, Amiri M, Pouy S. Risks of novel coronavirus disease (COVID-19) in pregnancy; a narrative review. *Archives of academic emergency medicine*. 2020; **8**(1).
- [31] Chen H, Guo J, Wang C, *et al.* Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *The lancet*. 2020; **395**(10226): 809-15.
- [32] Msq C, Jcs L, Sulaiman S, Tan H. From the frontline of COVID-19-How prepared are we as obstetricians: a commentary. <https://obgyn.onlinelibrary.wiley.com/doi/pdfdirect/101111/1471-052816192>. 2020.
- [33] Mardani M, Pourkaveh B. Neoscriber Demo Publisher. *A Controversial Debate: Vertical Transmission of COVID-19 in Pregnancy*[Google Scholar]. 2020.
- [34] Wang X, Zhou Z, Zhang J, Zhu F, Tang Y, Shen X. A case of 2019 Novel Coronavirus in a pregnant woman with preterm delivery. *Clinical infectious diseases*. 2020.

- [35] Favre G, Pomar L, Musso D, Baud D. 2019-nCoV epidemic: what about pregnancies? *Lancet (London, England)*. 2020; **395**(10224): e40.
- [36] Liang H, Acharya G. Novel corona virus disease (COVID-19) in pregnancy: What clinical recommendations to follow? 2020.
- [37] Rasmussen SA, Smulian JC, Lednický JA, Wen TS, Jamieson DJ. Coronavirus disease 2019 (COVID-19) and pregnancy: what obstetricians need to know. *American journal of obstetrics and gynecology*. 2020; **222**(5): 415-26.
- [38] Zhu N, Zhang D, Wang W, *et al*. A novel coronavirus from patients with pneumonia in China, 2019. *New England journal of medicine*. 2020.
- [39] Liu D, Li L, Wu X, *et al*. Pregnancy and perinatal outcomes of women with COVID-19 pneumonia: a preliminary analysis. Available at SSRN 3548758. 2020.
- [40] Lu Q, Shi Y. Coronavirus disease (COVID-19) and neonate: What neonatologist need to know. *Journal of medical virology*. 2020; **92**(6): 564-7.
- [41] Liu W, Wang Q, Zhang Q, Chen L, Chen J, Zhang B. Coronavirus disease 2019 (COVID-19) during pregnancy: a case series. 2020. *Preprint*. 2020.
- [42] Lam CM, Wong SF, Leung TN, *et al*. A case-controlled study comparing clinical course and outcomes of pregnant and non-pregnant women with severe acute respiratory syndrome. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2004; **111**(8): 771-4.
- [43] Silva CRACd, Oliveira LVd, Lopes LP, Santos WAGd, Agra IKR. Immunological aspects of coronavirus disease during pregnancy: an integrative review. *Revista da Associação Médica Brasileira*. 2020; **66**: 696-700.
- [44] Jiao J. Under the epidemic situation of COVID-19, should special attention to pregnant women be given? *Journal of medical virology*. 2020.

- [45] Liu H, Wang L-L, Zhao S-J, Kwak-Kim J, Mor G, Liao A-H. Why are pregnant women susceptible to COVID-19? An immunological viewpoint. *Journal of reproductive immunology*. 2020; **139**: 103122.
- [46] Dashraath P, Wong JLJ, Lim MXK, *et al*. Coronavirus disease 2019 (COVID-19) pandemic and pregnancy. *American journal of obstetrics and gynecology*. 2020; **222**(6): 521-31.
- [47] Wong SF, Chow KM, Leung TN, *et al*. Pregnancy and perinatal outcomes of women with severe acute respiratory syndrome. *American journal of obstetrics and gynecology*. 2004; **191**(1): 292-7.
- [48] Yan J, Guo J, Fan C, *et al*. Coronavirus disease 2019 in pregnant women: a report based on 116 cases. *American journal of obstetrics and gynecology*. 2020; **223**(1): 111. e1-. e14.
- [49] Cosma S, Carosso AR, Cusato J, *et al*. Coronavirus disease 2019 and first-trimester spontaneous abortion: a case-control study of 225 pregnant patients. *American journal of obstetrics and gynecology*. 2021; **224**(4): 391. e1-. e7.
- [50] Sahin D, Tanacan A, Erol SA, *et al*. Updated experience of a tertiary pandemic center on 533 pregnant women with COVID-19 infection: a prospective cohort study from Turkey. *International Journal of Gynecology & Obstetrics*. 2021; **152**(3): 328-34.
- [51] Ayed A, Embaireeg A, Benawadh A, *et al*. Maternal and perinatal characteristics and outcomes of pregnancies complicated with COVID-19 in Kuwait. *BMC pregnancy and childbirth*. 2020; **20**(1): 1-9.
- [52] Wong TC, Lee ZY, Sia TL, Chang AK, Chua HH. Miscarriage risk in COVID-19 infection. *SN comprehensive clinical medicine*. 2020; **2**(9): 1449-52.
- [53] Rana MS, Usman M, Alam MM, *et al*. First trimester miscarriage in a pregnant woman infected with COVID-19 in Pakistan. *Journal of Infection*. 2021; **82**(1): e27-e8.



Table-2: Number of miscarriages due to COVID-19 in different countries

Location	No. of patients	No. of miscarriages	Gestational weeks	References
Italy	225	12	Approx. 12-13 weeks	<sup>49</sup>
China	116	1	38 weeks	<sup>48</sup>
Turkey	553	12	1 <sup>st</sup> trimester (130) 2 <sup>nd</sup> trimester (165), Third trimester (238)	<sup>50</sup>
Kuwait	185	3	29 weeks	<sup>51</sup>
Malaysia	2	2	10 weeks	<sup>52</sup>
Pakistan	1	1	10 weeks	<sup>53</sup>
Italy	7	1	37 Weeks	<sup>54</sup>
China	118	9 (3 spontaneous abortion)	Mostly 3 <sup>rd</sup> trimester	<sup>55</sup>
France	54	1	37 weeks	<sup>56</sup>
22 countries	887	11	Approx. 20-20 weeks	<sup>57</sup>
Switzerland	1	1	19 weeks	<sup>58</sup>