

Methanol Extract of *Dryopteris dilatata* Alleviates streptozotocin-induced Hyperglycaemia and Liver Injury in Male Wistar Rats in Type 2 Diabetes Mellitus

ABSTRACT

Dysfunctional pancreatic beta cell and impaired glucose metabolism in the liver has been implicated in streptozotocin (STZ)-type 2 diabetes mellitus and this has been linked with increased oxidative stress. Whether methanol extract of *Dryopteris dilatata* (MEDd), a flavonoid-rich plant can ameliorate STZ- type 2 diabetes liver damage remains an issue. Hence, this study investigated the effect of methanol extract of *Dryopteris dilatata* on STZ- type 2 diabetes mellitus in male Wistar rat. Animals were randomly selected into five groups (n=5) and were treated as follows; group 1 received distilled water (10ml/kg), Group 2 received only STZ (60mg/kg), Group 3 and 4 received STZ then 400 and 800mg/kg of MEDd respectively while Group 5 received STZ then pioglitazone (10mg/kg). Following 14 days of treatment, animals were euthanized and blood as well as spleen, liver, pancreas and kidney tissues were collected for further studies. Our results revealed that MEDd significantly reduced STZ-induced hyperglycaemia in Diabetic rats. Markers of oxidative injury (MDA, NO and GSH) were also significantly ameliorated in the pancreas and liver of the diabetic rats following treatment with MEDd. However, liver injury markers (ALT, AST and ALP) were significantly attenuated with marked decreased in organ weight in the diabetic rats after treatment with MEDd. We found that methanol extract of *Dryopteris dilatata* demonstrated anti-diabetogenic and hepato-protective potential by enhancing *in vivo* hepato-pancreatic antioxidant defence system.

Keywords: *Dryopteris dilatata*; streptozotocin; diabetes mellitus; antioxidants; liver injury.

1. INTRODUCTION

“Diabetes mellitus (DM) is one of the most well-known chronic metabolic illnesses. It has been reported that it poses a hazard to human health and has had a significant negative impact in the twenty-first century. According to predictions, DM cases will continue to rise globally, perhaps leading to an increase in mortality rates if not properly controlled” [1,2]. This metabolic disease is caused by the pancreas inability to make insulin (insulin deficit) or properly use the insulin released (insulin resistance), or both, which can be caused by oxidative and inflammatory stress, among other things [3-5]. However, this condition resulted in elevated blood sugar levels, which are common in diabetic patients. The glycoproteins family, which includes glucose transporters (GLUTs), is responsible for glucose uptake in the cell (Simmons 2017). GLUT-2 regulates the

liver's glucose transporter, which supports glucose metabolism while also supplying molecules that promote the transcription of glucose-sensitive genes [6,7]. “Furthermore, insulin is responsible for glucose metabolism regulation and blocking insulin production resulted in increased hepatic glucose release and decreased glucose absorption in muscle cells” [8,7]. “Chemically active diabetogenic drugs like streptozotocin have been employed in laboratory animal models to study type 1 diabetes. The GLUT-2 glucose uniporter transports this drug through the plasma membrane of pancreatic beta-cells, specifically impairing them” [9-12]. “Hyperglycemia, hyperlipidaemia, hypertension, nephropathy, neuropathy, polyuria, polyphagia, stroke, ketosis, and other diabetic complications resulted from changes in glucose uptake” [13,14]. “There have been numerous advancements in the prevention,

treatment, and management of glycemic changes that contribute to DM. However, the incidence of DM-related complications is still a major concern around the world" [15]. The preferred anti-diabetic drugs used in its management or treatments has been reported to cause adverse reactions in the body after administration; therefore, finding a low-toxic and effective functional food product or less-adversely reactive drug for the prevention and management of DM is extremely important.

Nutraceuticals or medicinal plants, according to ethnopharmacology and therapeutic studies, may help manage diabetes and its consequences [16-20]. "Phytochemicals found in various plant-based components, such as vitamins, polyphenols, tannins, flavonoids, saponins, terpenes, and sterols, can boost the body's antioxidant defence mechanism" [21,18]. "However, a large number of studies have suggested that phyto-bioactive compounds with high antioxidant activity may have a variety of biological effects, including influencing glucose absorption and uptake, modulating insulin secretion, and managing or reducing the risk of developing DM complications" [4,14,22]. "*Dryopteris dilatata* is one of the plants used to treat health problems and the *Dryopteris dilatata*, popularly known as the 'Broad buckler fern,' is a member of the *Dryopteridaceae* family. It has dark green tripinnate fronds with brown scales on the ribs, and its ribs are covered in brown scales" (Runk *et al.*, 2012). "It's known as Okpomie and is found primarily in Nigeria's tropical region" (Mordi *et al.*, 2016; Akpotu *et al.*, 2018). The roots and leaves of the plant have a variety of medical uses, including treating dandruff on the scalp and removing worms from the body [23]. "We also discovered that an ethanol extract of the plant's leaves has anti-hyperlipidemic properties" [24]. Our research also demonstrated that plant leaves contain a large number of phytochemicals and active substances [24]. The effect of a methanol extract of *Dryopteris dilatata* (MEDd) on type 1 diabetes after streptozotocin administration was studied in this study using a mechanism involving endogenous antioxidant activities and liver function indicators.

2. MATERIALS AND METHODS

2.1 Chemicals and Drugs

Santa-cruz provided the streptozotocin (USA). In Port-Harcourt, Nigeria, pioglitazone was acquired at a community pharmacy. Sigma Aldrich produced thiobarbituric acid (TBA), Griess, and Ellman reagents (Germany). Burgoyne

Burbidges & Co. (Mumbai, India) provided the trichloroacetic acid (TCA), and Immunometrics Limited provided the insulin ELISA kit, AST, ALT, and ALP kits (UK). The rest of the reagents and solvents were of analytical quality.

2.2 Source of Experimental Animals

Twenty-five mature male Wistar rats weighing 120-150 g were purchased from the Department of Pharmacology and Therapeutics, PAMO University of Medical Sciences, Port Harcourt, Nigeria's central animal house. The animals were kept in regular laboratory conditions, as per the University's ethical guidelines, which adhere to the "Principle of Laboratory Animal Care" (NIH Publication No. 85-23). The rats were given free access to the regular rat meal (Ladokun feeds) and water for one week (12–12 h light–dark cycle, 28 °C).

2.3 Experimental Design

The animals were divided into five groups of six animals in each group

- Group 1 (Normal control)- Fed with normal feed and water ad libitum
- Group 2 (Diabetic control)- Fed with normal feed after induction of diabetes 60 mg/kg of STZ
- Group 3 (Treated group)- Fed with normal feed and water ad libitum after induction of diabetes with daily treatment with 800 mg/kg of Dd
- Group 4 (Treated group)- Fed with normal feed and water ad libitum after induction of diabetes with daily treatment with 400 mg/kg of Dd
- Group 5 (Treated group)- Fed with normal feed and water ad libitum after induction of diabetes with daily treatment with 500 mg/kg of pioglitazone

2.4 Plant Collection and Identification and Extract Preparation

Dryopteris dilatata fresh leaves were obtained from the Olomoro community in Isoko South, Delta State, Nigeria. The leaves were authenticated for herbarium numbering, FHI 110338, at the Forestry Research Institute of Nigeria (FRIN), Ibadan. Before extraction, the leaves were washed, air dried, and macerated into powder form. Furthermore, the Kemelayefa and Kagbo plant crude extraction methods were modified (2018). 70 percent methanol was used to extract the mixed powder, which was then filtered using Whatman No. 2 filter paper. A

rotary evaporator was then used to vacuum concentrate the solvent to dryness. The extract was then dried at 40°C in an incubator before being kept at 0-4°C. The yield of the extract was 13%. Finally, the dried extract was diluted in normal saline to get the needed dose of 800 mg/kg.

2.5 Diabetic Induction in Rats

“Fasted male Wistar rats were given a single intraperitoneal injection of streptozotocin (STZ; 60 mg/kg) in sterile citrate buffer (0.1 M, pH 4.5) to develop Type 2 Diabetes mellitus, as described” by Asiwe et al. [25]. The rats' diabetic status was determined after 72 hours using a glucometer (ACCU-CHEK® Active) and appropriate blood glucose test strips, and animals with a fasting blood glucose level of more than or equal to 200 mg/dl were chosen for the study [25].

2.6 Preparation of Blood Samples and Tissue Homogenates for Biochemical Analysis

2.7 Estimation of Levels of Lipid Peroxidation

“The amount of oxidative stress in diabetic rats caused by streptozotocin was calculated. The lipid peroxidation end product marker malondialdehyde was measured in the pancreas and liver supernatants using the thiobarbituric reactive assay (TBARS) as previously described” [26,27]. The amounts of TBARS in the tissues were measured in gmol MDA/mg tissue protein.

2.8 Measurement of Tissue Nitrite

Spectrophotometric methods with Griess reagent were used to quantify nitrite in the pancreas and liver. Griess reagent was made in a 1:1 ratio from reagents A (1 percent sulfanilamide in 5% phosphoric acid) and B (0.1 percent N-1-naphthyl ethylenediamine dihydrochloride). Griess reagent was used to incubate the samples, which were then examined at 540 nm in a spectrophotometer. A standard curve of sodium nitrite (0–100 M) was used to estimate the nitrite content.

2.9 Estimation of Glutathione Reduction

The amount of reduced glutathione (GSH) in pancreas and liver supernatants was determined using Ellman's reagent in a modified approach

At the end of the treatment duration, the rats were subjected to deep ether anaesthesia before euthanasia through cervical dislocation. Cardiac puncture was used for blood collection; the plasma was obtained by centrifugation at 3,000 rpm at room temperature using a bench top centrifuge (Bosch, UK). The pancreas, spleen and liver were excised, rinsed in iced-cold Tris-KCl buffer (0.1 M, pH 7.4), and then blotted with adsorbent paper. Furthermore, the liver and pancreas were homogenized in sodium phosphate buffer (0.1 M, pH 7.4), centrifuged at 10,000 rpm for 10 min at 4 °C. For biochemical analysis, the tissue supernatant (pancreas and liver) was collected and used for the estimation of malondialdehyde (MDA), Nitrite (NO) and reduced Glutathion (GSH). Liver function tests were also estimated from the plasma. For biochemical analysis, the tissue supernatant (pancreas and liver) was collected and used for the estimation of malondialdehyde (MDA), Nitrite (NO) and reduced Glutathion (GSH). Liver function tests were also estimated from the plasma.

[28]. 0.1 mL cell-free exudate supernatant was diluted 10 times and deproteinized with 1 mL Trichloroacetic acid (20%) before centrifugation at 10,000 rpm for 10 minutes at 4°C. The supernatant was then combined with 0.75 mL sodium phosphate buffer (0.1 M, pH 7.4) and 2 mL 5, 5-Dithios-nitrobenzoic acid (0.0006 M) (DTNB). In a UV/Vis spectrophotometer, the absorbance was measured at 412 nm in less than 5 minutes (752N INESA, China). The glutathione concentration, which was determined using a standard curve generated with standard glutathione (0–200 µM), was expressed as a function of the volume of tissue supernatant (µM GSH/mL of supernatant).

2.10 The Activity of the Liver Function Indicators

“Alanine aminotransferase (ALT), Aspartate transaminase (AST), and Alkaline Phosphatase (ALP) for liver damage mediators in the plasma were evaluated using a Randox test kit, as described by Reitman and Frankel” [29].

2.11 Serum Insulin Estimation

Following the manufacturer's instructions, the concentration of insulin in the blood was determined using an ELISA kit (Elabscience, UK).

2.12 Data Analysis

All data collected were represented as Mean \pm SEM (standard Error of Mean) and analysed using one-way and two-way analysis of variance (ANOVA) followed by Bonferroni post hoc multiple comparison tests. $P < 0.05$ was considered statistically significant. Graphs and statistical analyses were performed in GraphPad Prism software version 5.01 (GraphPad Software, Inc. La Jolla, CA 92037 USA).

3. RESULTS

3.1 Change in Body Weight

There was no significant change in body weight of animals at week 0. However, at week 1 and 2, diabetes control animals showed a reduced body weight but following treatment with MEDd and pioglitazone at week 1 and 2, the body weight were significantly normalized as presented in Fig. 1.




Fig. 1. Change in body weight (g)

All values are expressed as Mean \pm standard error of mean, (n=5), * $p < 0.05$ when compared with the diabetes control while # $p < 0.05$ when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 2. Organ weight of spleen, liver and kidney.

3.2 Relative Weight of Spleen, Liver and Kidney

Fig. 2 shows that there was no significant difference in the weight of spleen across the experimental groups. However, the liver showed a marked increase in diabetes control group. Following treatment with MEDd (400 and 800mg/kg) and pioglitazone (10mg/kg), the liver weight was significantly reduced. Similar data was observed in the weight of kidney though MEDd (400mg/kg) did not have any significant effect on the weight of the kidney when compared with the control.

3.3 Fasting Blood Glucose (mg/dl)

Following the induction of diabetes with STZ, the fasting glucose level was observed to be above 200mg/dl after 72hrs of induction. However, treatment with MEDd (400 and 800mg/kg) and pioglitazone significantly reduced the glucose level in a duration dependent manner as presented in Fig. 3.

All values are expressed as Mean \pm standard error of mean, (n=5), *p < 0.05 when compared with the diabetes control while #p < 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 3. Estimation of glucose

All values are expressed as Mean \pm standard error of mean, (n=5), *p < 0.05 when compared with the diabetes control while #p < 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 4. Oral Glucose Tolerance (OGT)

All values are expressed as Mean \pm standard error of mean, (n=5), *p < 0.05 when compared with the diabetes control while #p < 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 5. Estimation of serum insulin

All values are expressed as Mean \pm standard error of mean, (n=5), * $p < 0.05$ when compared with the diabetes control while # $p < 0.05$ was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone

3.4 Glucose Tolerance Test

Following oral administration of glucose, the glucose level in diabetes control group was significantly increased in 30, 60, 90, 120 and 150 minutes when compared with the normal control. However, treatment with MEDd (400 and 800mg/dl) and pioglitazone (10mg/kg) significantly reduce the glucose level in 30, 60, 90, 120, and 150 minutes when compared with diabetes control but was significantly elevated when compared with normal control animals as shown in Fig. 4.

3.5 Estimation of Serum Insulin

Fig. 5 represent the serum insulin concentration. The serum insulin concentration was significantly reduced in diabetes control animal following the STZ induction. However, treatment with MEDd (400 and 800mg/kg) and pioglitazone significantly increase the serum concentration of insulin.

3.6 Liver Enzymes

Figs. 6-8 represents the markers of liver injury. Streptozotocin induction significantly increases serum ALP, AST and ALT in diabetes control rats when compared with normal control rats. However, the treatment of MEDd (400 and 800mg/dl) and pioglitazone significantly reduce AST and ALT when compared with diabetes control animals (Figs. 6 and Fig. 7). ALP was not significant different when MEDd (400mg/kg) was compared with the diabetes control animals (Fig. 8).

3.7 Pancreatic Oxidative Stress Markers

Presented in Figs. 9-11 is the pancreatic oxidative stress marker. There was significant increase in pancreatic MDA and Nitrite level in diabetes control animals while GSH level was significantly reduced in diabetes control animal. However, following treatment with MEDd (400 and 800mg/dl) and pioglitazone (10mg/kg) significantly reduces the MDA as well as elevated the pancreatic GSH level when compared to diabetes control animal. Treatment

with MEDd (400 and 800mg/kg) did not show any significant difference in pancreatic Nitrite level when compared with diabetes control animal.

3.8 Liver Antioxidants

Liver MDA and Nitrite level were significantly elevated in diabetes control animals. However, treatment with MEDd (400 and 800mg/kg) and pioglitazone significantly down-regulated MDA and Nitrite levels when compared with diabetes control animals. Liver GSH level was significantly reduced following STZ induction as shown in Fig. 12 and Fig. 13. However, MEDd (800mg/kg) and pioglitazone significantly upregulated liver GSH level in comparison with diabetes control while MEDd (400mg/kg) did not show any significant difference in liver GSH level when compared with diabetes control animals (Fig. 14).

4. DISCUSSION

“Retrospective research have showed that diabetes is a major worry that causes a variety of negative effects in the body, particularly in the brain, liver, kidneys, cardiovascular system, and musculoskeletal system. This effect could be due to uncontrollable reactive oxygen species (ROS) generation, which alters gene morphology, triggering gene mutations, and resulting in the synthesis of undesired proteins” [30]. Furthermore, flavonoids and bioactive substances have been found to reduce metabolic dysregulation in animal models of diabetes and other problems in the past [13,15]. “The findings of this study on the methanol extract of *Dryopteris dilatata* (MEDd) corroborated previous reports on its ability to treat diabetes and associated consequences” [31,24,32]. “Our research found evidence that *Dryopteris dilatata* therapy can reduce the risk of diabetic complications such as hyperglycemia and hyperlipidaemia. *Dryopteris dilatata* includes bioactive phytochemicals/compounds discovered in a recent study” [24] that can help to improve health by avoiding oxidative damage to cells [31,24].




Fig. 6. Aspartate transaminase (AST)

All values are expressed as Mean \pm standard error of mean, (n=5), *p < 0.05 when compared with the diabetes control while [#]p < 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 7. Alanine amino transferase (ALT)

All values are expressed as Mean \pm standard error of mean, (n=5), *p < 0.05 when compared with the diabetes control while [#]p < 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 8. Alkaline phosphatase (ALP)

All values are expressed as Mean \pm standard error of mean, (n=5), *p< 0.05 when compared with the diabetes control while #p< 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 9. Pancreatic MDA level

All values are expressed as Mean \pm standard error of mean, (n=5), *p< 0.05 when compared with the diabetes control while #p< 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 10. Pancreatic Nitrite level

All values are expressed as Mean \pm standard error of mean, (n=5), *p< 0.05 when compared with the diabetes control while #p< 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 11. Pancreas antioxidant, GSH level

All values are expressed as Mean \pm standard error of mean, (n=5), *p< 0.05 when compared with the diabetes control while #p< 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 12. Liver MDA level

All values are expressed as Mean \pm standard error of mean, (n=5), *p< 0.05 when compared with the diabetes control while #p< 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 13. Liver nitrite level

All values are expressed as Mean \pm standard error of mean, (n=5), *p< 0.05 when compared with the diabetes control while #p< 0.05 was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone




Fig. 14. Liver GSH level

All values are expressed as Mean \pm standard error of mean, (n=5), * $p < 0.05$ when compared with the diabetes control while # $p < 0.05$ was significant when compared with the normal control. D = streptozotocin induction, MEDd = methanol extract of *Dryopteris dilatata*, Pio = Pioglitazone

Several researches on tropical plants have discovered a strong link between antioxidant activity and polyphenol levels [33-35,25]. Repeated oral treatment with MEDd, on the other hand, was found to reduce the effects of streptozotocin (STZ)- type 2 diabetes induced oxidative stress and hyperglycemia in this investigation. In STZ-induced diabetic rats, the glucose-lowering impact of repeated oral therapy with MEDd was considerable. In the meantime, a single dose of MEDd did not significantly reduce STZ-induced hyperglycemia in diabetic rats. We also discovered that after putting diabetic rats through an oral glucose test for 30, 60, 90, 120, and 150 minutes, MEDd was able to reverse the hyperglycemic condition. In addition, the STZ-induced diabetic rats' glucose levels were favourably modulated after week 1 and 2 of treatment compared to baseline and 72 hours after streptozotocin injection. Normoglycemia is a major aim in diabetes that must be met in order to avoid organ damage and inflammation caused by persistently high blood sugar levels [36,15,25,37] Streptozotocin is an antibiotic that inhibits the production of insulin, a peptide hormone responsible for the conversion of glucose to glycogen in the liver, by targeting the beta cells of the pancreas. Reduced insulin production or insulin receptor insensitivity has been linked to impaired glucose utilization [25,37,22]. In line with prior research, our investigation found a drop in serum insulin in the diabetes control group, resulting in the diabetes phenotype seen in this study. Treatment with MEDd and Pioglitazone, on the other hand, increased serum insulin levels, which could be attributable to *Dryopteris dilatata's* antioxidant

impact on pancreatic beta cells [32]. "Poor blood glucose levels have also been connected to the uncontrollable development of diabetes complications such as neuropathies, cardiovascular and renal dysfunction, and cognitive and behavioral impairment, according to reports" [38,36,15]. "Several plant items have been utilized as nutraceuticals to help diabetics control their blood glucose levels" [39,31,40,24,42]. As a result, our findings suggest that *Dryopteris dilatata* could be utilized as a nutraceutical in diabetes patients to help them control their blood sugar levels.

"The etiology of diabetic vascular disease has been linked to hyperglycemia-induced oxidative stress" [42]. "However, in diabetes, the production of reactive oxygen species (ROS) due to hyperglycemia causes oxidative and inflammatory stress in a variety of organs and tissues" [15]. "The unbalanced production of free radicals and the diminished endogenous antioxidant defence system are the most common causes of oxidative and inflammatory stress" [43,44]. Increased reactive oxygen species (ROS) combined with a decrease in the body's natural antioxidants increase cell and tissue damage, accelerating the development of diabetic complications [45]. "The use of STZ to induce diabetes in rats resulted in significant alterations in oxidative stress markers indicators in the rats' pancreas and liver. Oxidative stress markers, on the other hand, disrupts oxidative and antioxidative system functions, which is linked to cellular lipid peroxidation" [27]. Organ toxicity is linked to an increase in ROS production that exceeds the cellular capacity to

eliminate toxicants (Rashid et al. 2013). The increased nitrite and malondialdehyde (MDA) levels in the pancreas and liver of STZ- type 2 diabetes induced diabetic mice were dramatically lowered by MEDd in this investigation. In our work, glutathione, an oxidative stress markers, was shown to be dramatically reduced in the pancreas and liver of STZ-induced diabetic rats, implying a large increase in oxidative stress markers caused by ROS. The inactivation produced by excessive free radical production in hyperglycemic rats may explain the lower amount of GSH reported in diabetic control rats [15]. Additionally, due to exacerbated O_2^- and H_2O_2 levels, this GSH reduction increased nitrite and malondialdehyde levels in the pancreas and liver. Meanwhile, MEDd treatment reduced GSH loss in diabetic rats' pancreatic and liver. This demonstrated that the extract had free radical scavenging and antioxidant capabilities. When these effects are added together, the results confirm the antioxidant potentials of *Dryopteris dilatata* ethanol extract previously reported [24]. Streptozotocin- type 2 diabetes induced changes in metabolic activities and functions in rats on biomarkers of hepatic functions were reversed in animals treated with MEDd. Streptozotocin- type 2 diabetes induced significant liver damage in diabetic rats, as evidenced by increased Alanine amino transferase (ALT), Aspartate transaminase (AST), and Alkaline Phosphatase (ALP) activity. Treatment with MEDd, on the other hand, reduced the levels of liver damage indicators in the diabetic rats' plasma. One of the most common diabetic consequences is hepatotoxicity. Diabetes caused hepatic damage, which resulted in uncontrolled glucogenesis and elevated protein levels. In diabetics, the enzymes ALT, AST, and ALP have been linked to the conversion of amino acids to keto acids [46,47]. However, the hepato-protective effect of *Dryopteris dilatata* methanol extract seen in this investigation could be due to its antiradical scavenging potential, as evidenced by reduced hepato-pancreatic lipid peroxidation (NO and MDA levels) and enhanced hepato-pancreatic GSH levels. After STZ administration, this action also reduced the increased relative liver and kidney weight produced by hyperglycemia in diabetic rats. Furthermore, increasing liver and kidney weight in patients and animal models is widely considered as an indicator of diabetes [15,48,25]. However, scavenging reactive oxygen species in diabetic-induced rats is likely to be the most effective defence mechanism offered by this therapeutic plant (methanol extract of *Dryopteris dilatata*). *Dryopteris dilatata* is capable of alleviating hyperglycemia and lowering liver

toxicity in diabetic patients due to all of these effects.

5. CONCLUSION

In conclusion, the current investigation proved *Dryopteris dilatata's* medicinal potential in the treatment of blood glucose changes. However, increased *in-vivo* antioxidant activity in the pancreas and liver facilitate this anti-diabetogenic and hepatoprotective action. The phytochemicals found in the *Dryopteris dilatata* leaf may also be responsible for these activities.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The PAMO University of Medical Sciences Animal Research Ethics Committee, which agreed with the "Guide to the Care and Use of Laboratory Animals in Research and Teaching" as prescribed in NIH publications volume 25 No.28 revised in 1996, approved the use of animals for this study with approval number PUMS-AREC/056.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Giovannini P, Howes MJR, Edwards SE. Medicinal plants used in the traditional management of diabetes and its sequelae in Central America: A review. *Journal of Ethnopharmacology*. 2016; 184:58-71.
2. Liu C, Ge HM, Liu BH, Dong R, Shan K, Chen X, Yan B. Targeting pericyte-endothelial cell crosstalk by circular RNA-cPWWP2A inhibition aggravates diabetes-induced microvascular dysfunction. *Proceedings of the National Academy of Sciences*. 2019;116(15):7455-7464.
3. Baynes HW. Classification, pathophysiology, diagnosis and management of diabetes mellitus. *Journal of Diabetes and Metabolism*. 2015;6(5):1-9.
4. Miaffo D, Guessom Kamgue O, Ledang Tebou N, Maa Maa Temhoul C, Kamanyi A. Antidiabetic and antioxidant potentials of *Vitellaria paradoxa* barks in alloxan-induced diabetic rats. *Clinical Phytoscience*. 2019;5(1):1-8.

5. Islam Z. Japan Epidemiology Collaboration on Occupational Health Study Group, et al. Prediabetes, diabetes, and the risk of all-cause and cause-specific mortality in a Japanese working population: Japan epidemiology collaboration on occupational health study. *Diabetes Care*. 2021;44(3):757–764.
6. Leturque A, Brot-Laroche E, Le Gall M, Stolarczyk E, Tobin V. The role of GLUT2 in dietary sugar handling. *Journal of Physiology and Biochemistry*. 2005;61(4):529-537.
7. Chunudom L, Thongsom M, Karim N, Rahman MA, Rana MN, Tangpong J. *Tithonia diversifolia* aqueous fraction plays a protective role against alloxan-induced diabetic mice via modulating GLUT2 expression. *South African Journal of Botany*. 2020;133:118-123.
8. Al-Goblan AS, Al-Alfi MA, Khan MZ. Mechanism linking diabetes mellitus and obesity. *Diabetes, Metabolic Syndrome and Obesity Targets and Therapy*. 2014;7:587–591.
9. Gorus FK, Malaisse WJ, Pipeleers DG. Selective uptake of alloxan by pancreatic B-cells. *The Biochemical Journal*. 1982;208(2):513–515.
10. Elsner M, Tiedge M, Guldbakke B, Munday R, Lenzen S. Importance of the GLUT2 glucose transporter for pancreatic beta cell toxicity of alloxan. *Diabetologia*. 2002;45(11):1542-1549.
11. Asiwe JN, Anachuna KK, Moke EG, Sanusi KO, Okonofua DE, Omeru O, Fasanmade AA. High dietary salt intake alleviates fasting blood glucose in streptozotocin-induced diabetic male Wistar rats. *Thai J Pharm Sci*. 2021; 45(3):172-177.
12. Haddad D, Al Madhoun A, Nizam R, Al-Mulla F. Role of caveolin-1 in diabetes and its complications. *Oxidative Medicine and Cellular Longevity*; 2020.
13. Anachuna KK, Oyem CJ, Nwoguzie BC, Asiwe JN. Glucose lowering effects and histomorphological changes of *Vernonia amygdalina* on pancreatic compromised wistar rats using alloxan monohydrate. *Trop. J. of Health Sciences*. 2018;25(2):27-31.
14. Malfa GA, Tomasello B, Acquaviva R, Mantia AL, Pappalardo F, Ragusa M, Di Giacomo C. The antioxidant activities of *Betula etnensis* Rafin. ethanolic extract exert protective and anti-diabetic effects on streptozotocin-induced diabetes in rats. *Antioxidants*. 2020;9(9):847.
15. Ajayi AM, Adedapo ADA, Badaki VB, Oyagbemi AA, Adedapo AA. *Chrysophyllum albidum* fruit ethanol extract ameliorates hyperglycaemia and elevated blood pressure in streptozotocin-induced diabetic rats through modulation of oxidative stress, NF-κB and PPAR-γ. *Biomed Pharmacother*. 2021;2:141:111879. DOI: 10.1016/j.biopha.2021.111879. Epub ahead of print. PMID: 34225016.
16. Roy B. Biomolecular basis of the role of diabetes mellitus in osteoporosis and bone fractures. *World Journal of Diabetes*. 2013;4(4):101–113.
17. Samarghandian S, Azimi-Nezhad M, Farkhondeh T. Immunomodulatory and antioxidant effects of saffron aqueous extract (*Crocus sativus* L.) on streptozotocin-induced diabetes in rats. *Indian Heart Journal*. 2017;69 (2):151–159.
18. El Rabey HA, Al-Seeni MN, Bakhshwain AS. The antidiabetic activity of *Nigella sativa* and propolis on streptozotocin-induced diabetes and diabetic nephropathy in male rats. *Evidence-Based Complementary and Alternative Medicine: eCAM*. 2017;5439645.
19. Alawode DI, Asiwe JN, Moke EG, Okonofua DE, Sanusi KO, Adagbada EO, Yusuf MO, Fasanmade AA. The effect of ethanol leaf extract of *Cnidiosculus aconitifolius* on cardiorenal functions in hypertensive and normotensive male wistar rats. *Int J Nutr Sci*. 2021;6(3):155-160. DOI: 10.30476/IJNS.2021.92067.1145.
20. Alawode DI, Asiwe JN, Moke EG, Okonofua DE, Sanusi KO, Adagbada EO, Yusuf MO, Fasanmade AA. The Effect of Ethanol Leaf Extract of *Cnidiosculus Aconitifolius* on Cardiorenal Functions in Hypertensive and Normotensive Male Wistar Rats. *Int J Nutr Sci*. 2021;6(3):155-160. DOI: 10.30476/IJNS.2021.92067.1145.
21. Miguel-Carrasco JL, Zambrano S, Blanca AJ, Mate A, Vázquez CM. Captopril reduces cardiac inflammatory markers in spontaneously hypertensive rats by inactivation of NF-κB. *Journal of Inflammation*. 2010;7(1):1-9.
22. Okonofua, DE, Asiwe JN, Anachuna KK, Moke EG, Sanusi KO, Adagbada EO, Yusuf MO, Alawode DI, Fasanmade AA. Effect of diabetes mellitus and

- hypertension on osmotic fragility and hemorheological factors in male wistar rats. *Biology, Medicine, & Natural Product Chemistry*. 2021;10 (2):73-79.
23. Rünk K, Zobel M, Zobel K. Biological Flora of the British Isles: *Dryopteris carthusiana*, *D. dilatata* and *D. expansa*. *J. Ecol*. 2012;100 (4):1039 –1063.
 24. Akpotu Ajirioghene, Celestine Ani, Choice Nworgu, Okorie Pamela, Igwe Uzoma, Jide Uzoigwe, Adeyemo Mercy, Nwaeme Ogochukwu, Obinna Onwujekwe. Antidiabetic and anti- hyperlipidemic effects of ethanolic extract of *Dryopteris dilatata* leaves. *Journal of Diabetes and Endocrinology*; 2018.
 25. Asiwe JN, Kolawole TA, Anachuna KK, Ebuwe EI, Nwogweze BC, Eruotor H, Igbokwe V. Cabbage juice protect against Lead-induced liver and kidney damage in male Wistar rat. *Biomarkers*. 2022;27(2):151–158. DOI:10.1080/1354X.2021.2022210
 26. Adam-vizi V, Seregi M. Receptor dependent stimulatory effect of noradrenaline on Na-K ATPase in rat brain homogenate: Role of lipid peroxidation. *Biochem. Pharmacol.* 1982;31:2231-2236.
 27. Bakre A, Odusanya S, Olowoparija S, Ojo O, Olayemi J, Aderibigbe A. Behavioral and biochemical evidences for antidepressant activity of ethanol extract of *Jatropha curcas* in mice subjected to chronic unpredictable mild stress. *Journal of Biology and Nature*. 2020;1-10.
 28. Sin YM, Pook SH, Tan TM, Petterssoon A, Kara AU, Teh WF. Changes in glutathione and its associated enzymes during carrageenan-induced acute inflammation in mice. *Comp Biochem s Physiol*. 1997;116C:191–5.
 29. Reitman S, Frankel S. Glutamic-pyruvate transaminase assay by colorimetric method. *Am. J. Clin. Path.* 1957;28:56.
 30. Klil-Drori AJ, Azoulay L, Pollak MN. Cancer, obesity, diabetes, and antidiabetic drugs: Is the fog clearing? *Nature reviews. Clinical Oncology*. 2017;14 (2):85–99.
 31. Mordi JC, Ewhre OL, Ojebah C. Aqueous leaf extract of *Dryopteris dilatata* on STZ – induced diabetic wistar rats with associated hyperlipidemic; 2016.
 32. Akpotu EA, Ghasi SI, Ewhre LO, Adebayo OG, Asiwe JN. Anti-diabetogenic and in vivo antioxidant activity of ethanol extract of *Dryopteris dilatata* in alloxan-induced male Wistar rats. *Biomarkers*. 2021; 26:8:718-725.
 33. Bayili RG, Abdoul-Latif F, Kone OH, Diao M, Bassole IH, Dicko MH. Phenolic compounds and antioxidant activities in some fruits and vegetables from Burkina Faso. *African Journal of Biotechnology*. 2011;10(62):13543-13547.
 34. Oboh G, Ademosun AO, Akinleye M, Omojokun OS, Boligon AA, Athayde ML. Starch composition, glycemic indices, phenolic constituents, and antioxidative and antidiabetic properties of some common tropical fruits. *Journal of Ethnic Foods*. 2015;2(2):64-73.
 35. Adebayo OG, Onasanwo SA, Ajayi AM, Aduema W, Oyebanjo OT, Nicodemus OU. *Cnidioscolus aconitifolius*-supplemented diet enhanced neurocognition, endogenous antioxidants and cholinergic system and maintains hippocampal neuronal integrity in male Wistar rats. *Drug Metabolism and Personalized Therapy*; 2021.
 36. Afroz A, Ali L, Karim M, Alramadan MJ, Alam K, Magliano DJ, Billah B. Glycaemic control for people with type 2 diabetes mellitus in Bangladesh-an urgent need for optimization of management plan. *Scientific Reports*. 2019;9(1):1-10.
 37. David UE, Asiwe JN, Fasanmade AA. Maternal hypothyroidism prolongs gestation period and impairs glucose tolerance in offspring of wistar rats. *Hormone Molecular Biology and Clinical Investigation*, PMID: 34907695 (ahead of print); 2021.
 38. Al-Badri A, Hashmath Z, Oldland GH, Miller R, Javaid K, Syed AA, Chirinos JA. Poor glycemic control is associated with increased extracellular volume fraction in diabetes. *Diabetes Care*. 2018;41(9):2019-2025.
 39. Christensen AS, Viggers L, Hasselström K, Gregersen S. Effect of fruit restriction on glycemic control in patients with type 2 diabetes—a randomized trial. *Nutrition Journal*. 2013;12(1):1-6.
 40. Li Y, Wang DD, Ley SH, Vasanti M, Howard AG, He Y, Hu FB. Time trends of dietary and lifestyle factors and their potential impact on diabetes burden in China. *Diabetes Care*. 2017; 40(12):1685-1694.
 41. Sadiya A, Mnla R. Impact of food pattern on glycemic control among type 2 diabetic patients: a cross-sectional study in the United Arab Emirates. *Diabetes, metabolic*

- syndrome and obesity: Targets and Therapy. 2019;12:1143–1150.
42. Ahmadi S, Awliaei H, Haidarizadeh M, Rostamzadeh J. The effect of ethanolic extract of *urtica dioica* leaves on high levels of blood glucose and gene expression of glucose transporter 2 (Glut2) in liver of alloxan-induced diabetic mice. *Gene Cell Tissue*. 2015;2(3):e30355.
 43. Pantoja PKD, Colmenares DAJ, Isaza MJH. New caffeic acid derivative from *Tithonia diversifolia* (Hemsl.) A. gray butanolic extract and its antioxidant activity. *Food and Chemical Toxicology*. 2017;109:1079–1085.
 44. Karim N, Rahman A, Chanudom L, Thongsom M, Tangpong J. Mangosteen vinegar rind from *Garcinia mangostana* prevents high-fat diet and streptozotocin-induced type II diabetes nephropathy and apoptosis. *Journal of Food Science*. 2019;84(5):1208-1215.
 45. Danilova IG, Bulavintceva TS, Gette IF, Medvedeva SY, Emelyanov VV, Abidov MT. Partial recovery from alloxan-induced diabetes by sodium phthalhydrazide in rats. *Biomedicine & Pharmacotherapy*. 2017;95:103-110.
 46. Qian K, Zhong S, Xie K, Yu D, Yang R, Gong DW. Hepatic ALT isoenzymes are elevated in gluconeogenic conditions including diabetes and suppressed by insulin at the protein level. *Diabetes/Metabolism Research and Reviews*. 2015;31(6):562-571.
 47. Safhi MM, Alam MF, Sivakumar SM, Anwer T. Hepatoprotective potential of *Sargassum muticum* against STZ-induced diabetic liver damage in wistar rats by inhibiting cytokines and the apoptosis pathway. *Analytical Cellular Pathology*; 2019.
 48. Mobasher MA, Germoush MO, Galal El-Tantawi H, Samy El-Said K. Metformin improves biochemical and pathophysiological changes in hepatocellular carcinoma with pre-existed diabetes mellitus rats. *Pathogens*. 2021; 10(1):59.

© 2022 Ajrioghene et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.