

1 **Phenolic compounds** from the immature fruits of *Solanum nigrum* L.

2 **Abstract**

3 *Solanum nigrum* L. has been traditionally used as traditional Chinese medicine and
4 food all over the world. Phytochemical investigations of the immature fruits of *S.*
5 *nigrum* have been carried out and five coumarins (**1-5**) and five phenolic compounds
6 (**6-11**) were obtained. Their structures were elucidated on the basis of spectroscopic
7 and chemical methods, including IR, MS, ¹H and ¹³C-NMR. They were identified as
8 6,7-dimethoxycoumarin(**1**), 6-methoxy-7-hydroxycoumarin(**2**),
9 6,8-dimethoxycoumarin(**3**), 6,7-dimethoxycoumarin(**4**), umbelliferone(**5**), vanillin(**6**),
10 vanillic acid(**7**), syringic aldehyde(**8**), isovanillic acid(**9**), ethyl
11 3,4-dihydroxybenzoate(**10**), and ethyl 3,4-dihydroxyphenylacetate(**11**).

12 **Keywords:** *Solanum nigrum* L., immature fruits, chemical investigation, coumarin,
13 phenolic compound

14 **1. Introduction**

15 “*Solanum nigrum* L. belongs to the family of Solanaceae and is widely distributed
16 all over the world” [1]. “In China, *S. nigrum* has been used as a common traditional
17 Chinese medicine, which has the effects of clearing heat and detoxification, activating
18 blood circulation and removing blood stasis, promoting water and swelling, and is
19 mainly used to treat cold and fever, toothache, and cancers. Modern pharmacological
20 studies indicated that *S. nigrum* exhibited a variety of biological activities including
21 antiproliferative [2], anti-inflammatory [3], antiviral [4], hepatoprotective [5], and
22 antioxidative activities” [6].

23 “*S. nigrum* contains steroidal saponins, steroidal alkaloid glycosides and phenolic
24 compounds” [7-10], anthocyanin and polysaccharides [11-13] in the previous
25 chemical studies. “Steroidal alkaloid glycosides are the main chemical components of
26 *Solanum* species, which possess various pharmacological activities such as
27 antiproliferative and anti-inflammatory properties” [14, 15]. Currently, most of the
28 chemical investigations on *S. nigrum* were focused on the aerial parts, while the
29 bioactive components of its unripe berries are still unclear.

30 In this study, the phytochemicals from the immature fruits of *S. nigrum* has been
31 carried out to investigate its phytochemical profile, and eleven compounds were
32 obtained, including five coumarins (1-5) and five phenolic compounds (6-11).

33

34 **2. MATERIALS AND METHODS**

35 **2.1. General experimental procedures**

36 “IR spectra ($4000-450\text{ cm}^{-1}$) were recorded with a Spectrum 100 FT-IR
37 spectrometer (Perkin Elmer Inc., Waltham, MA). MS spectra were obtained on an
38 Acquity UPLC-Q-TOF Microsystem (Waters Co., Milford, MA). NMR spectra were

39 taken on a Bruker Avance III 500 MHz spectrometer (Bruker, Switzerland)” [10].
40 ODS packed column (40–60 μm, Merck KGaA, Darastadt, Germany) and column
41 chromatography was employed on silica gel (Anhui Liangchen Silicon Source
42 Material Co. Ltd, Lu’ an, China). All other analytical chemicals and reagents were
43 purchased from Sinopharm Chemical Reagent Co. Ltd. (Shanghai, China).

44

45 **2.2. Plant materials**

46 The immature fruits of *S. nigrum* were purchased from Haerbin (Heilongjiang
47 province, China) and dried at room temperature in the shade. The plant was
48 authenticated by Prof. X. J. He and the voucher specimen has been deposited in the
49 School of Pharmacy, Guangdong Pharmaceutical University, China.

50

51 **2.3. Extraction and isolation**

52 Dried immature fruits of *S. nigrum* (2.5 kg) were extracted with 70% EtOH (v/v, 15
53 L × 3) for 2 h. The ethanol-free suspension was subjected to a D101 macroporus resin
54 column (80 × 1100 mm), and eluted with H₂O, 10% MeOH, 30% MeOH, 50% MeOH,
55 70% MeOH and MeOH to give six fractions (I to VI). The 30% MeOH elution
56 (fraction IV, 51.0 g, yield 2.04%) was separated by a silica gel chromatography
57 column (200-300 mesh, 1500 g) into 12 fractions (IV-1 to IV-12) with a
58 CHCl₃-MeOH gradient (10:1 to 0:1, v/v).


59 Compounds **6** (15.1 mg, purity 95.1%), **7** (7.9 mg, purity 96.2%), and **10** (13.7 mg,
60 purity 90.5%) were obtained from fraction IV-6 followed by an ODS column, eluted
61 with MeOH-H₂O gradient (1:9 to 10:0, v/v). Fraction IV-8 was applied to an ODS
62 column eluted with a gradient of MeOH-H₂O (1:9 to 10:0, v/v) to afford five
63 subfractions (IV-8-1 to IV-8-5). Subfraction IV-8-3 was further purified by an

64 Octadecylsilyl gel (ODS) medium pressure liquid chromatography (MPLC) and
 65 semi-preparative HPLC to obtain compounds **1** (9.6 mg, purity 97.0%), **2** (7.3 mg,
 66 purity 94.6%), **4** (11.2 mg, purity 98.1%), and **8** (6.1 mg, purity 97.3%). Fraction
 67 IV-10 was subjected to an ODS MPLC and eluted with MeOH-H₂O gradient (1:9 to
 68 10:0, v/v) to afford nine fractions (IV-10-1 to IV-10-9). Subfraction IV-10-5 was
 69 further separated by an ODS column and semi-preparative HPLC to obtain
 70 compounds **3** (20.1 mg, purity 92.6%), **5** (10.5 mg, purity 97.0%), **9** (7.1 mg, purity
 71 97.2%), and **11** (6.0 mg, purity 98.0%), respectively.

72

73 3. RESULTS AND DISCUSSION

74 The 70% ethanol extract of the immature fruits of *S. nigrum* was separated
 75 successively by column chromatography on D101 macroporus resin, silica gel, ODS
 76 MPLC, and preparative HPLC, to afford five coumarins (**1-5**) and five phenolic
 77 compounds (**6-11**) (Fig. 1). Their structures were elucidated on the basis of
 78 spectroscopic data, including IR, MS, and ¹H and ¹³C-NMR.



79
80

81 **Fig. 1** Structures of the coumarins and phenolic compounds **1-11**

82 Compound **1** was white amorphous powder and easily soluble in chloroform and

83 methanol. There was a blue fluorescence under 365 nm after TLC development. Its
84 molecular formula was determined as $C_{11}H_{10}O_4$ according to the ion m/z 207.0654
85 $[M+H]^+$ (calcd for $C_{11}H_{11}O_4$, 207.0652) in the positive HR-ESI-MS. In the 1H -NMR
86 (Table 1), δ_H 6.26 (1H, d, $J=9.3$ Hz) and 7.61 (1H, d, $J=9.3$ Hz) were the characteristic
87 hydrogen signals of coumarins. δ_H 3.90 (3H, s) and 3.92 (3H, s) were two hydroxyl
88 groups. In the ^{13}C -NMR, a total of 11 carbon signals were given, including an ester
89 carbonyl carbon at δ_C 161.4. Signals at δ_C 113.6 and 143.3 were carbon signals at C-3
90 and C-4 of coumarin, and signals at δ_C 108.0, 146.5, 150.3, 100.2, 153.0, 111.5 were
91 the carbon signals on the benzene ring. δ_C 56.5 is a methoxy group. Based on above
92 analysis and the literature [16], compound **1** was identified as
93 6,7-dimethoxycoumarin.

94 Compound **2** was yellowish needle crystal (MeOH) and there was a blue
95 fluorescence under 365 nm after TLC development. Its molecular formula was
96 determined as $C_{10}H_8O_4$ according to the ion m/z 191.0342 $[M+H]^+$ (calcd for $C_{10}H_7O_4$,
97 191.0350) in the positive HR-ESI-MS. In the 1H -NMR (Table 1), δ_H 6.23 (1H, d,
98 $J=9.4$ Hz) and 7.68 (1H, d, $J=9.4$ Hz) were the characteristic signals of coumarins. δ_H
99 3.82 (3H, s) was a hydroxyl group. In the ^{13}C -NMR, a total of 10 carbon signals were
100 given, including an ester carbonyl carbon at δ_C 160.7. Signals at δ_C 109.5 and 144.5
101 were carbon signals at C-3 and C-4 of coumarin, and signals at δ_C 111.6, 145.3, 151.3,
102 102.8, 149.5, 110.5 were the carbon signals on the benzene ring. δ_C 56.0 is a methoxy
103 group. Based on above analysis and the literature [17], compound **2** was identified as
104 6-methoxy-7-hydroxycoumarin.

105 Compound **3** was white amorphous powder and easily soluble in methanol. There
 106 was a blue fluorescence under 365 nm after TLC development. Its molecular formula
 107 was determined as C₁₁H₁₀O₅ according to the ion *m/z* 221.0446 [M-H]⁻ (calcd for
 108 C₁₁H₉O₅, 221.0455) in the negative HR-ESI-MS. In the ¹H-NMR (Table 1), δ_H 6.24
 109 (1H, d, J=9.3 Hz) and 7.88 (1H, d, J=9.3 Hz) were the characteristic hydrogen signals
 110 of coumarins. δ_H 3.82 (3H, s) and 3.83 (3H, s) were two hydroxyl groups. In the
 111 ¹³C-NMR, a total of 11 carbon signals were given, including an ester carbonyl carbon
 112 at δ_C 160.3. Signals at δ_C 112.0 and 144.6 were carbon signals at C-3 and C-4 of
 113 coumarin, and signals at δ_C 104.7, 145.5, 144.5, 134.2, 143.2, 110.1 were the carbon
 114 signals on the benzene ring. δ_C 56.2 and 60.6 is a methoxy group. Based on above
 115 analysis and the literature [18], compound **3** was identified as
 116 6,8-dimethoxycoumarin.

117 **Table 1** ¹H and ¹³C NMR data of compounds **1-5** (δ in ppm and *J* in Hz)^a

No	1		2		3		4		5	
	δ _H	δ _C	δ _H	δ _C	δ _H	δ _C	δ _H	δ _C	δ _H	δ _C
2		161.4		160.7		160.3		160.2		160.2
3	6.26 d, (9.3)	113.6	6.23 d, (9.4)	109.5	6.24 d, (9.3)	112.0	6.30 d, (9.4)	114.0	6.21 d, (9.4)	111.5
4	7.61 d, (9.3)	143.3	7.68 d, (9.4)	144.5	7.88 d, (9.3)	144.6	7.90 d, (9.4)	144.2	7.92 d, (9.4)	144.3
5	6.81, s	108.0	7.02, s	111.6	7.01, s	104.7	6.82, s	100.5	7.50 d, (8.4)	129.8
6		146.5		145.3		145.5		149.5	6.76 dd, (8.4, 2.3)	113.3
7		150.3		151.3		144.5		140.0		161.5
8	6.80, s	100.2	6.77, s	102.8		134.2		138.3	6.70 d, (2.3)	102.0
9		153.0		149.5		143.2		138.1		155.3
10		111.5		110.5		110.1		114.1		111.1
6-OCH ₃	3.90, s	56.5	3.82, s	56.0	3.83, s	56.2	3.83, s	56.1		
7-OCH ₃	3.92, s	56.5					3.80, s	60.5		

118 a: 500 MHz for ¹H and 126 MHz for ¹³C in in DMSO-*d*₆

119 Compound **4** was colorless needle crystal (MeOH). There was a blue
120 fluorescence under 365 nm after TLC development. Its molecular formula was
121 determined as C₁₁H₁₀O₅ according to the ion *m/z* 245.0412 [M+H]⁺ (calcd for
122 C₁₁H₁₀O₅Na, 245.0420) in the positive HR-ESI-MS. In the ¹H-NMR (Table 1), δ_{H}
123 6.30 (1H, d, J=9.4 Hz) and 7.90 (1H, d, J=9.4 Hz) were the characteristic hydrogen
124 signals of coumarins. δ_{H} 3.80 (3H, s) and 3.83 (3H, s) were two hydroxyl groups. In
125 the ¹³C-NMR, a total of 11 carbon signals were given, including an ester carbonyl
126 carbon at δ_{C} 160.2. Signals at δ_{C} 114.0 and 144.2 were carbon signals at C-3 and C-4
127 of coumarin, and signals at δ_{C} 100.5, 149.5, 140.0, 138.1, 138.3, 114.1 were the
128 carbon signals on the benzene ring. δ_{C} 56.1 and 60.5 is a methoxy group. The NMR
129 data of **4** is very close to that of **3**, and the difference in chemical shift values between
130 the two is due to the different positions of hydroxyl and methoxy groups. Based on
131 above analysis and the literature [19], compound **4** was identified as
132 6,7-dimethoxycoumarin.

133 Compound **5** was white amorphous powder and there was a blue fluorescence
134 under 365 nm after TLC development. Its molecular formula was determined as
135 C₉H₆O₃ according to the ion *m/z* 161.0240 [M-H]⁻ (calcd for C₉H₅O₃, 161.0244) in
136 the negative HR-ESI-MS. In the ¹H-NMR (Table 1), δ_{H} 6.21 (1H, d, J=9.4 Hz) and
137 7.92 (1H, d, J=9.4 Hz) were the characteristic hydrogen signals of coumarins. In the
138 ¹³C-NMR, a total of 9 carbon signals were given, including an ester carbonyl carbon

139 at δ_C 160.2. Signals at δ_C 111.5 and 144.3 were carbon signals at C-3 and C-4 of
140 coumarin, and signals at δ_C 161.5, 155.3, 129.8, 113.3, 111.1, 102.0 were the carbon
141 signals on the benzene ring. Based on above analysis and the literature [20],
142 compound **5** was identified as umbelliferone.

143 Compounds **6** to **11** were identified as vanillin (**6**) [21], vanillic acid (**7**) [22],
144 syringic aldehyde (**8**) [23], isovanillic acid (**9**) [24], ethyl 3,4-dihydroxybenzoate (**10**)
145 [25], and ethyl 3,4-dihydroxyphenylacetate (**11**) [26], respectively, based on their
146 spectroscopic analysis.

147

148 **4. Conclusions**

149 Further chemical investigation of the immature fruits of *S. nigrum* led to the
150 isolation of five coumarins and five phenolic compounds were obtained. Their
151 structures were elucidated on the basis of spectroscopic and chemical methods. They
152 were identifies as 6,7-dimethoxycoumarin, 6-methoxy-7-hydroxycoumarin,
153 6,8-dimethoxycoumarin, 6,7-dimethoxycoumarin, umbelliferone, vanillin (**6**), vanillic
154 acid, syringic aldehyde, isovanillic acid, ethyl 3,4-dihydroxybenzoate, and ethyl
155 3,4-dihydroxyphenylacetate.

156

157

158 **ACKNOWLEDGEMENTS**

159 This work was financially supported by Guangzhou Science and technology
160 planning project of Guangzhou Municipal Science and Technology Bureau
161 (202002020091).

162 **COMPETING INTERESTS**

163 Authors have declared that no competing interests exist.

164 **Supplementary data**

165 The original spectra of compounds **1** to **11** including MS, ¹H and ¹³C NMR were
166 given as supplementary data.

167

168 **REFERENCES**

- 169 [1] F.Z. Wang, J. Tang, Flora of China, Science Publishing House, Beijing, China, 1980, pp. 76-79.
- 170 [2] X. Ding, F.S. Zhu, M. Li, S.G. Gao, Induction of apoptosis in human hepatoma SMMC-7721 cells
171 by solamargine from Solanum nigrum L. J. Ethnopharmacol. 2012, 139:599-604.
- 172 [3] Z.A. Zakaria, H.K. Gopalan, H. Zainal, N.H.M. Pojan, N.A. Morsid, A. Aris, M.R. Sulaiman,
173 Antinociceptive, anti-inflammatory and antipyretic effects of Solanum nigrum chloroform extract in
174 animal models. Yakugaku Zasshi, 2006, 126:1171-1178.
- 175 [4] T. Javed, U.A. Ashfaq, S. Riaz, S. Rehman, S. Riazuddin, In-vitro antiviral activity of Solanum
176 nigrum against Hepatitis C virus. Virol. J. 2011, 8:26.
- 177 [5] C.C. Hsieh, H.L. Fang, W.C. Lina, Inhibitory effect of Solanum nigrum on thioacetamide-induced
178 liver fibrosis in mice. J. Ethnopharmacol. 2008, 119:117-121.
- 179 [6] Y.S. Aly, M.A. Shallan, Antioxidant properties of wild Solanum nigrum ripe fruit. Planta Med. 2011,
180 77:1367-1367.
- 181 [7] X.L. Zhou, X.J. He, G.H. Wang, H. Gao, G.X. Zhou, W.C. Ye, X.S. Yao, Steroidal saponins from
182 Solanum nigrum, J. Nat. Prod. 2006, 69: 1158-1163.
- 183 [8] Y. Wang, L. Xiang, X. Yi, X. He, Potential anti-inflammatory steroidal saponins from the berries of

- 184 Solanum nigrum L. (European Black Nightshade). J. Agric. Food Chem. 2017, 65:4262-4272.
- 185 [9] L. Xiang, Y. Wang, X. Yi, X. He, Anti-inflammatory steroidal glycosides from the berries of
- 186 Solanum nigrum L. (European black nightshade). Phytochemistry, 2018, 148:87-96.
- 187 [10] L. Xiang, Y. Wang, X. Yi, X. He. Steroidal alkaloid glycosides and phenolics from the immature
- 188 fruits of Solanum nigrum. Fitoterapia, 2019, 137:104268.
- 189 [11] E.A. Eltayeb, A.S. AlAnsari, J.G. Roddick. 1997. Changes in the steroidal alkaloid solasodine
- 190 during development of *Solanum nigrum* and *Solanum incanum*. Phytochemistry, 1997, 46:489-494.
- 191 [12] J. Li, Q.W. Li, D.W. Gao, Z.S. Han, W.Z. Lu. Antitumor and immunomodulating effects of
- 192 polysaccharides isolated from *Solanum nigrum* Linne. Phytother. Res. 2009, 23:1524-1530.
- 193 [13] J. Li, Q. Li, Y. Peng, R. Zhao, Z. Han, D. Gao. Protective effects of fraction 1a of polysaccharides
- 194 isolated from *Solanum nigrum* Linne on thymus in tumor-bearing mice. J. Ethnopharmacol. 2010,
- 195 129:350-356.
- 196 [14] Y. Wang, J. Xu, Y. Wang, L. Xiang, X. He. S-20, a steroidal saponin from the berries of black
- 197 nightshade, exerts anti-multidrug resistance activity in K562/ADR cells through autophagic cell death
- 198 and ERK activation. Food Funct. 2022, 13:2200-2215.
- 199 [15] S. Emmanuel, S. Ignacimuthu, R. Perumalsamy, T. Amalraj, Antiinflammatory activity of Solanum
- 200 trilobatum. Fitoterapia 2006, 77:611-612.
- 201 [16] X. Xu, J. Yang. Study on aromatic chemical constituents of *Aralia subcapitata* Hoo. *China J. Chin.*
- 202 *Mat. Medica* 2005, 30:398-399.
- 203 [17] Z. Dawa, Y. Zhou, Y. Bai, S. Sangge, P. Xie, L. Ding. Studies on chemical constituents of
- 204 *Saussurealaniceps*. *China J. Chin. Mat. Medica* 2008, 33:1032-1035.
- 205 [18] F. Tian, L. Zhang, J. Tian, W. Zhou. Chemical constituents of *Artemisia anomala* S.Moore. *Chin.*

206 *Med. Chem.* 2008, 18:362-365.

207 [19] T. Yasuda, M. Fukui, T. Nakazawa T, et al. Metabolic fate of fraxin administered orally to rats. *J.*
208 *Nat. Prod.* 2006, 69:755-757.

209 [20] L. Kong, Z. Min. Studies on chemical constituents of *Euphorbia pekinensi*. *Acta Pharm. Sin.* 1996,
210 31:524-529.

211 [21] E. Kolehmainen, K. Laihia, J. Knuutinen, et al. ^1H , ^{13}C and ^{17}O NMR study of Chlorovanillins and
212 some related compounds. *Magnet. Reson. Chem.* 1992, 30:253-258

213 [22] L. Chen, W. Zhou. Chemical constituents from *Aeschynanthus longicaullis*. *China J. Chin. Mat.*
214 *Medica* 2009, 34:2758-2760.

215 [23] X. Yang, M. Wong, N. Wang, X. Yao. Effect of phenolic acids isolated from *Sambucus williamsii*
216 on proliferation and differentiation of rat osteoblastic UMR106 cells. *Chin. Trad. Herb. Drugs*, 2005,
217 36:1604-1607.

218 [24] J. Han, M. Wang. Chemical constituents of *Pyrrhosia sheareri*. *J. Nanjing Coll. Pharm.* 1984, 15:
219 40-43.

220 [25] T. Yang, C. Wang, H. Liu, et al. A new antioxidant compound from *Capparis spinosa*. *Pharm. Biol.*
221 2010, 48:589-594.

222 [26] K. Kanie, M. Nishii, T. Yasuda, et al. Self-assembly of thermotropic liquid-crystalline folic acid
223 derivatives: hydrogen-bonded complexes forming layers and columns. *J. Mat. Chem.* 2001, 11:
224 2875-2886.