

FACTORS AFFECTING RECURRENCE RATE OF URETHRAL STRICTURE FOLLOWING INTERNAL URETHROTOMY

ABSTRACT

BACKGROUND

Urethral stricture (post-scarring contraction in the urethral mucosa) is among the most challenging clinical conditions in urology. In Pakistan, Urethral stricture is a high-volume disease and constitutes 4-5% of urology workload (majority of them being in the 3rd to 5th decade of life). Different urethral strictures have different management options depending upon site, size and number of recurrences. OIU is indicated for urethral stricture for small (1.5 cm) bulbar and membranous urethral stricture. However, recurrence of strictures after OIU is quite common (4 to 5%). It has been proposed that duration and size of catheter after OIU affects the recurrence rate. The foremost management choice is internal urethrotomy, but recurrences are common, and many believe that the duration and size of catheterization post internal urethrotomy may affect the recurrence rate.

OBJECTIVE

To determine the factors affecting recurrence rates of urethral stricture after internal urethrotomy.

METHODOLOGY

It was Quasi Experimental Study conducted in Urology ward of Dr. Ziauddin University Hospital, Karachi from 10th March 2020 to 31 december 2021. Total 60 patients were included; all of the patients were divided into four equal groups. All the patients were followed quarterly for 12 months and data was recorded for urinary flow, retention, burning sensation during urination. UFM, Cystoscopy and urethrogram were done once indicated. Symptomatic patients or patients with UFM 15ml/sec were checked for recurrence followed by cystoscopy if needed. The data analysis was obtained using SPSS v. 21.0.

RESULTS

The mean age of the sample stood at 27 (SD \pm 4) years (range: 20 to 40) years were enrolled in the study. Urethral stricture disease recurred cumulatively in 24 (40%) patients within 12 months. Mean duration of catheterization was 8.71 ± 1.1 and 2.97 ± 0.91 days in patients with and without recurrences, respectively ($p=0.001$). Catheter dwell times for ≥ 7 days was associated with increased recurrence ($p=0.039$). Of the patients with recurrent strictures, increased catheter diameter was also associated with higher recurrence rates ($p=0.041$).

Conclusion

After careful consideration, it can be concluded that longer catheter in-dwell time and greater catheter diameter lead to a greater probability of stricture recurrence and hence the indwell time should be limited to ≤ 3 days and the catheter size chosen to be no more than 16Fr.

KEYWORDS

Urethral Stricture, Stricture Recurrence, Internal Urethrotomy, Catheter Diameter, & Catheter Indwell Time.

INTRODUCTION

A urethral stricture manifests as a contraction cause by scaring and fibrosis of mucosa of urethra and or adjacent tissue of spongiosum in the urethral cavity. (1) Urethral strictures are still the most challenging problem for urologist.(2) The urethral strictures greatly ruin the patient's QOL (quality of life) by causing urinary disorders and also destroy the entire urinary tract leading to loss of function of kidneys.(3)

Urethral strictures prevalence in industrial countries is predicted to be around 0.9% with an estimated incidence of 0.6%.(4) Approximately more than 5,000 medical consultations per year are affected by urethral strictures.(5) Almost all strictures that can determine the cause can be obtained. A recent meta-analysis of etiology found that most common causes are idiopathic (33%) and iatrogenic (33%), followed by post-traumatic (19%) and inflammatory (15%).(6) The management of urethral stricture depends upon the location, length and characteristics of the stricture due to its complexity.(7) It also depends on the anatomy, pathophysiology and specific reconstruction technique for each patient.(8)

Cochrane review carried out in 2012 reported that there are four main procedures accessible and convenient for the management of urethral stricture: periodic dilatation, endoscopic urethrotomy (Optical Internal Urethrotomy), indwelling ureteral stent or urethroplasty each having its own indications.(9) The technique of Opening stricture through transurethral incision with cold knife under direct visualization-DVIU (direct visualization internal urethrotomy) is known as internal urethrotomy.(10) For a urethral stricture which is single, short (< 1 cm) , flimsy and not treated previously, optical internal urethrotomy remains first line of treatment.(11)

Leaving indwelling transurethral catheters after optical internal urethrotomy is a routine practice. However, the duration and French(fr) size of indwelling urethral catheter after OIU is debatable.(12)

Several studies have proven its efficacy in smooth recovery post OIU and decreasing recurrence rate.(13) No evidence has been reported to have difference in the recurrence rate with varying indwelling catheter size 14fr-24fr.(14) Various authors suggest that the time duration of catheterization should be modified according to the features of the stricture.(15) A US urologist survey found that 51% of patients placed the catheter for 2-7 days.(16) Dutch urologists reported that catheters were removed only one day after DVIU in 81% of patients who underwent OIU in Holland.(17)

Major limitations of OIU are high recurrence rate. Wang et al, in their study reported that recurrence rate after OIU within 1 year is around 50%. Multiple techniques have been used by different surgeons to minimize the fibrosis and subsequent contraction of wound in order to prevent the stricture.(18) Unfortunately no technique has been proven successful in preventing the urethral stricture recurrence.(19) Since there is no consensus on optimum duration and size of post OIU indwelling catheter internationally as well as locally. We designed this study to reach a conclusion that what duration and size of post OIU indwelling per urethral catheterization is optimum in order to minimize the rate of recurrence of urethral strictures.

OBJECTIVE

To determine the factors affecting recurrence rates of urethral stricture after internal urethrotomy.

METHODS AND MATERIALS

This quasi-experimental study was conducted upon a sample of 60 patients (after taking written informed consent), divided into 4 groups of 15 each (aged 20 to 40 years, presenting with a urethral stricture from _____ to _____ at Dept. of Urology – Dr. Ziauddin University Hospital, Karachi. Primary endpoint (success) was achieved if patient was asymptomatic and the flow rate > 15 mL / sec on UFM. Symptomatic patients were checked for recurrence. The data obtained was analyzed using SPSS v. 21.0.

Inclusion Criteria:

- ❖ Patients aged 20-40 years
- ❖ Stricture less than 1.5 cm
- ❖ Patients presenting with single stricture

Exclusion Criteria:

- ❖ Post Radiation strictures or failure of Direct visualization internal urethrotomy surgery, and use of urethral stent instead of a catheter
- ❖ Patients with prostate enlargement on urethroscopic examination or medication
- ❖ Patients' post-hypospadias surgery

All cases were performed under the supervision of 2 consultant urologists, a two way silicon urethral catheters (16 Fr-18Fr) was placed for at the end of DVIUs. Patients were grouped as:

Group A had post op indwelling catheter of 16 Fr size for < 3days

Group B had post op indwelling catheter of 16 Fr size for > 7days

Group C had post op indwelling catheter of 18 Fr size for < 3days

Group D had post op indwelling catheter of 18 Fr size for >7 days

Patients with urethral catheters were sent home on their first postoperative day. After removal of the catheter, on the operating day depending on the groups in which they are allocated, patients were followed up on 12 weekly interval at 1, 3, 6, 9, and 12 months postoperatively. Symptoms associated with recurrence were noted, including reduced urinary flow, urinary retention, and burning sensation during urination. On followup if the patient is asymptomatic and the flow rate is > 15 mL / sec on UFM, the procedure considered successful. Patient with cystoscopy and Retrograde urethrography were followed and checked for recurrence if the patient was symptomatic or experienced a significant decrease in flow rate.

RESULTS

The mean age of the sample stood at 27 (SD ± 4) years (range: 20 to 40) years were enrolled in the study. Urethral stricture disease recurred cumulatively in 24 (40%) patients within 12 months. Mean duration of catheterization was 8.71 ± 1.1 and 2.97 ± 0.91 days in patients with and without recurrences, respectively (p=0.001). Catheter dwell times for ≥7 days was associated with increased recurrence (p=0.039). Of the patients with recurrent strictures, increased catheter diameter was also associated with higher recurrence rates (p=0.041).

Table 01: Summary of Descriptive Statistics

Variable		Statistic – n (%)
Age (Years)	Mean	27 (SD ± 05)
	Range	20 to 40
Duration of post operative catheterization (Days)	Mean in Recurrence	8.71 ± 1.1
	Mean in Without Recurrence	2.97 ± 0.91
Age Groups categorized according to 5 year bracket	Up to 25	9 (15%)
	26 to 30	17 (28.3%)
	31 to 35	20 (33.3%)
	36 to 40	14 (23.4%)
	Yes	24 (40%)

Recurrence	No	36 (60%)
Mean duration of urethral stricture (Months)	3 Months	7 (11.6%)
	6 Months	9 (15%)
	9 Months	5 (8.3%)
	12 Months	3 (5%)
Etiology of urethral stricture	Known	24 (40%)
	Idiopathic	36 (60%)
Size of urethral stricture (cm)	Mean	0.62± (SD=0.34)
Site of Urethral Stricture	Penile urethra	6 (10%)
	Bulbar urethra	11 (18.3%)
	Membranous Junction	7 (11.6%)

Table 02: Recurrence of Urethral Stricture in all Groups

Variable		Group I		Group II	
		≤ 3 days & 16Fr	≤ 3 days & 18Fr	≥ 7 days & 16Fr	≥ 7 days & 18Fr
Time of recurrence in months	3 Months	2 (8.3%)	2 (8.3%)	2 (8.3%)	1 (4.1%)
	6 Months	1 (4.1%)	2 (8.3%)	2 (8.3%)	4 (16.7%)
	9 Months	0 (0%)	1 (4.1%)	1 (4.1%)	3 (12.5%)
	12 Months	0 (0%)	1 (4.1%)	0 (0%)	2 (8.3%)

Variable		Group I		Group II		P Value
		≤ 3 days & 16Fr	≤ 3 days & 18Fr	≥ 7 days & 16Fr	≥ 7 days & 18Fr	
Age in Years	Mean	31.1 (SD ±12.2)	31.4 (SD ±9.8)	32.4 (SD ±13.2)	37.1 (SD ±10.8)	0.097*
	Range	18 to 40	18 to 40	18 to 40	18 to 40	

Table 03: Group v/s Site of Stricture

<i>Variable</i>		<i>Group I</i>		<i>Group II</i>		<i>P Value</i>
		≤ 3 days & 16Fr	≤ 3 days & 18Fr	≥ 7 days & 16Fr	≥ 7 days & 18Fr	
Site of Stricture	Penile urethra	2 (8.3%)	1 (4.1%)	2 (8.3%)	1 (4.1%)	0.021 ^a
	Bulbar urethra	1 (4.1%)	3 (12.5%)	2 (8.3%)	5 (20.8%)	
	Membranous Junction	0 (0%)	2 (8.3%)	1 (4.1%)	4 (16.7%)	

DISCUSSION

The third most frequent urological clinical problem in men is urethral stricture disease. In brief anterior urethral strictures less than 1cm, optical internal urethrotomy (OIU) is a well-accepted therapy of choice, however the main concern with OIU is a high recurrence rate of urethral stricture till date, there is no consensus on optimum size and duration of post-operative catheter to reduce rate of recurrence of urethral stricture. So the goal of this study was to see how the length and size of catheterization after an internal urethrotomy affected the rate of urethral stricture recurrence. Regardless of the approach used, an indwelling urethral catheter was inserted at the end of the procedure. Our findings revealed that post-operative urethral stricture recurrence is more likely in patients who have been catheterized for a long time (7 days) or who have big (18Fr) catheters.

Our study's mean age was 27 (SD±4) years, with participants ranging in age from 20 to 40 years old. Same findings were documented by Santucci et al. who discovered that urethral stricture disease is common in the younger and middle aged population, with a considerable increase around the age of years. (20) In addition, the findings of Islam et al. (21) and Ali et al. (22) have opposite the study's findings. Patients over the age of 40 are more likely to develop urethral stricture, according to both of them. This can be explained as current study included the patients' up to the 40 years of age. Due to this limitation mean age differs with the other studies mean.

In the current study, we found that 24 (40%) of patients in both groups had urethral stricture recurrence at multiple follow-up intervals within 12 months with the mean duration of catheterization was 8.71 ± 1.1 and 2.97 ± 0.91 days in patients with and without recurrences, respectively. Our results are strengthened by study done in 2016, Emrah Yuruk found the similar result that urethral stricture disease

recurred in 45 (23.31 percent) of patients during the first year after DVIU. According to him patients with and without recurrences had a mean catheterization duration of 7.47 ± 4.03 and 4.79 ± 1.94 days, respectively ($p=0.0001$). Catheter stay durations of less than 5 days were likewise linked to higher recurrence ($p=0.0001$). 16.18, 22Fr, and 22Fr catheters were inserted in 22.22 percent, 20 percent, and 57.78 percent of the patients with recurrent strictures, respectively. Higher recurrence rates were also linked to larger catheter diameters ($p=0.004$).[\(23\)](#)

Because the observation intervals and recurrence criteria were not constant, comparing stricture series was problematic. Recurrence rates range from 10% to 58% of the time. [\(24\)](#) Because most patients treated with optical urethrotomy still have major abnormalities following urethrotomy, this reflects the nature of the urethral strictures, which certainly do not cure fully but stable after urethrotomy. [\(25\)](#) Even after several years, some of them begin to improve. Nonetheless, as previously documented, the majority of recurrences occurred during the first year, emphasizing the significance of vigilant follow-up during this time. [\(26\)](#)

Our research also discovered that membranous and bulbar urethral strictures were the most common types of strictures in those who had been catheterized for more than 7 days, with the membranous urethra coming in second. A substantial relationship between the location and recurrent urethral strictures was seen in all groups (p -value 0.021).

Cheng Li, in 2018 discovered in his research that the bulbar urethra is responsible for 46.9% of urethral strictures, followed by penile urethral strictures (30.5%).[\(27\)](#) The likelihood of sigmoid curvature and reactive fixity of the urethra are the most common causes of urethral stricture at the bulbar level. [\(28\)](#) This is because of the anatomical curve, which predisposes to bulbar urethral injury in, iatrogenic and straddle urethral injury. [\(29\)](#)

In addition, we discovered that urethral stricture recurrence occurred between 6 and 9 months following the procedure, and that the rate of recurrence declined with time until no new instances were reported at 1 year. Naude and Heyns found almost identical findings, stating that if recurrence occurs, it usually happens 3 to 12 months following OIU. [\(30\)](#) This could be due to the use of toxic latex catheters for prolonged periods of time likely aggravated the local mechanical injuries to the urethra by speeding up local cell loss and inflammation while delaying the healing process in the affected areas. [\(31\)](#)

Furthermore, in our study we found a statistically significant relationship between the size and length of catheterization kept in all groups. The size and duration (or length) of the catheter remained inside increases the chances of recurring stricture. Desmond et al. suggested that patients with few previous treatments keep the urethral catheter for at least three days, and that patients with poor detrusor function keep it for up to four weeks. [\(32\)](#) Lipsky and Hubmer published the results of a study involving 32 individuals who underwent DVIU. They used a silicon catheter for about 7 days and suggested a 6-month hydraulic self-dilation protocol. After the operation, 25 (83%) of these individuals voided well and required no additional therapy. [\(33\)](#) One reason for the higher recurrence rate of the strictures noticed over time could be the use of "toxic" catheters during optical urethrotomy. [\(34\)](#)

Another retrospective study evaluated the recurrence rates of 798 strictures in relation to the postoperative catheter drainage. (35) In 89.7% of the instances, a silicone catheter was employed, with a mean length of 5.5 days. With 1–3, 4–7, and more than 7 days of catheterization, 34 percent, 43 percent, and 65 percent of patients, respectively, suffered recurrence. According to the authors, postoperative catheter drainage for fewer than 3 days appears to reduce the likelihood of recurrence. They didn't say how they came up with the duration of catheterization or the size of the silicone catheter they utilized.(30)

There is a scarcity of information on the implications of catheter size on the outcome of IU. In most cases of recurrent urethral stricture, however, ischemia is recognized to have a role in the establishment of the stricture. (36) Larger bore catheters may reduce blood flow and obstruct the re-epithelization process, which will eventually result in the healing of the incised urethra with fibrosis, due to the pressure they exert on the urethral wall.(37) Karakus et al. evaluated meatal stenosis rates in patients who had tabularized incised plate urethroplasty and found that tubularization of the urethral plate over a small-sized (6Fr) catheter prevented meatal stenosis regardless of the patients' age. (38)

CONCLUSIONS

After careful consideration, it can be concluded that longer catheter in-dwell time and greater catheter diameter lead to a greater probability of stricture recurrence and hence the indwell time should be limited to ≤ 3 days and the catheter size chosen to be no more than 16Fr.

REFERENCES

1. Lal A, Singh P, Bhujade H. 12 Urinary Bladder and Urethral Diseases. Comprehensive Textbook of Diagnostic Radiology: Four Volume Set. 2021:199.
2. Sterling J, Policastro C, Nikolavsky D. Pathophysiology of radiation-induced urethral strictures and therapeutic strategies optimizing outcomes of surgical repair. Scientific Advances in Reconstructive Urology and Tissue Engineering: Elsevier; 2022. p. 51-80.
3. Sekido N, Igawa Y, Kakizaki H, Kitta T, Sengoku A, Takahashi S, et al. Clinical guidelines for the diagnosis and treatment of lower urinary tract dysfunction in patients with spinal cord injury. International Journal of Urology. 2020;27(4):276-88.
4. Ademi I, Ademi M. INTERNAL URETHROTOMY AS A CHOICE OF TREATMENT ON A RECURRENT URETHRAL STRICTURE. KNOWLEDGE-International Journal. 2020;40(4):695-9.
5. Mangera A, Osman N, Chapple C. Evaluation and management of anterior urethral stricture disease. F1000Research. 2016;5.

6. Partin AW, Peters CA, Kavoussi LR, Dmochowski RR, Wein AJ. Campbell-Walsh-Wein Urology Twelfth Edition Review E-Book. 2020.
7. Wessells H, Angermeier KW, Elliott S, Gonzalez CM, Kodama R, Peterson AC, et al. Male urethral stricture: American urological association guideline. *The Journal of urology*. 2017;197(1):182-90.
8. Xiong S, Wang J, Zhu W, Yang K, Ding G, Li X, et al. Onlay repair technique for the management of ureteral strictures: A comprehensive review. *BioMed Research International*. 2020;2020.
9. Wong SS, Aboumarzouk OM, Narahari R, O'Riordan A, Pickard R. Simple urethral dilatation, endoscopic urethrotomy, and urethroplasty for urethral stricture disease in adult men. *Cochrane Database of Systematic Reviews*. 2012(12).
10. Halpern J, Cordon B, Armenakas N, Brandes S. MP36-16 SALVAGE INTERNAL URETHROTOMY FOR MANAGEMENT OF URETHRAL STRICTURE FOLLOWING URETHROPLASTY. *The Journal of urology*. 2017;197(4S):e472-e.
11. Lee YJ, Kim SW. Current management of urethral stricture. *Korean Journal of Urology*. 2013;54(9):561-9.
12. Poelaert F, Oosterlinck W, Spinoit A-F, Lumen N. Duration of urethral catheterization after urethroplasty: How long is enough? *Minerva Urologica e Nefrologica*. 2017;69(4):372-6.
13. Redon-Galvez L, Molina-Escudero R, Álvarez-Ardura M, Otaola-Arca H, Parra RA, Páez-Borda Á. Predictors of urethral stricture recurrence after endoscopic urethrotomy. *Actas Urológicas Españolas (English Edition)*. 2016;40(8):529-33.
14. Peng Y, Li X, Li X, Zhang C, Yang K, Xiong G, et al. Fluoroscopy-free minimally invasive ureteral stricture balloon dilatation: a retrospective safety and efficacy cohort study. *Translational Andrology and Urology*. 2021;10(7):2962.
15. Kumar V, Dharwadkar S, Doshi C, Batta A. Role of intermittent self catheterization in prevention of recurrence of stricture urethra following visual internal urethrotomy: a prospective single centre study. *International Surgery Journal*. 2019;6(4):1093-6.
16. Awad MA, Osterberg EC, Chang H, Gaither TW, Alwaal A, Fox R, et al. Urethral catheters and medical malpractice: a legal database review from 1965 to 2015. *Translational Andrology and Urology*. 2016;5(5):762.
17. Djordjevic ML. Treatment of urethral stricture disease by internal urethrotomy, dilation, or stenting. *European Urology Supplements*. 2016;15(1):7-12.
18. Wang L, Wang Z, Yang B, Yang Q, Sun Y. Thulium laser urethrotomy for urethral stricture: a preliminary report. *Lasers in Surgery and Medicine*. 2010;42(7):620-3.
19. Hughes M, Blakely S, Nikolavsky D. Advancements in transurethral management of urethral stricture disease. *Current Opinion in Urology*. 2021;31(5):504-10.
20. Santucci RA, Mario LA, Aninch JWMc. Anastomotic urethroplasty for bulbar urethral stricture: analysis of 168 patients. *The Journal of urology*. 2002;167(4):1715-9.
21. 97. Shams-UI-Islam BW, Ali Shahid M, Ansari I, Haider M. Role of mitomycin C in anterior urethral stricture recurrence after internal urethrotomy. *Pak J Med Health Sci*. 2019;13(2):497-9.
22. 98. Ali L, Shahzad M, Orakzai N, Khan I, Ahmad M. Efficacy of mitomycin C in reducing recurrence of anterior urethral stricture after internal optical urethrotomy. *Korean Journal of Urology*. 2015;56(9):650-5.
23. 99. Yürük E, Yentur S, Çakır ÖO, Ertaş K, Şerefoğlu EC, Semerciöz A. Catheter dwell time and diameter affect the recurrence rates after internal urethrotomy. *Turkish Journal of Urology*. 2016;42(3):184.
24. 100. Perdzyński W, Adamek MJCEJoU. Calibration of urethra with anti-scar gel as a new preventive method for stricture recurrence after urethroplasty performed due to refractory anterior urethral stricture disease. 2020;73(1):80.
25. 101. Thirugnanasundralingam V, Tasevski R, Lawrentschuk N. *Benign Surgical Conditions. Men's Health and Wellbeing: Springer; 2022. p. 205-95.*

26. Hussain M, Khan MS, Lal M, Hashmi A, Naqvi SAA, Rizvi SAH. Stricture of urethra: patterns and outcomes of management from a single centre in Pakistan over 7 years. *J Coll Physicians Surg Pak*. 2020;30(1):79-84.
27. Cheng L, Li S, Wang Z, Huang B, Lin J. A brief review on anterior urethral strictures. *Asian Journal of Urology*. 2018;5(2):88-93.
28. Rourke KF, Welk B, Kodama R, Bailly G, Davies T, Santesso N, et al. Canadian Urological Association guideline on male urethral stricture. *Canadian Urological Association Journal*. 2020;14(10):305.
29. Ali L, Shahzad M, Orakzai N, Khan I, Ahmad MJKJoU. Efficacy of mitomycin C in reducing recurrence of anterior urethral stricture after internal optical urethrotomy. 2015;56(9):650-5.
30. Naudé AM, Heyns CF. What is the place of internal urethrotomy in the treatment of urethral stricture disease? *Nature clinical practice Urology*. 2005;2(11):538-45.
31. Pang KH, Chapple CR, Chatters R, Downey AP, Harding CK, Hind D, et al. A systematic review and meta-analysis of adjuncts to minimally invasive treatment of urethral stricture in men. 2021;80(4):467-79.
32. Desmond A, Evans C, Jameson R, Woolfenden K, Gibbon N. Critical evaluation of direct vision urethrotomy by urine flow measurement. *British Journal of Urology*. 1981;53(6):630-3.
33. Lipsky H, Hubner G. Direct vision urethrotomy in the management of urethral strictures. *British Journal of Urology*. 1977;49(7):725-8.
34. Abbas A, Qazi SM, Altaf AJJoUM, College D. COMPARISON OF END TO END ANASTOMOTIC URETHROPLASTY AND OPTICAL URETHROTOMY IN THE MANAGEMENT OF PATIENTS PRESENTING WITH SHORT SEGMENT INCOMPLETE URETHRAL STRICTURE. 2018;9(2):8-11.
35. Pansadoro V, Emiliozzi P. Internal urethrotomy in the management of anterior urethral strictures: long-term followup. *The Journal of urology*. 1996;156(1):73-5.
36. Tian Y, Wazir R, Yue X, Wang K-J, Li H. Prevention of stricture recurrence following urethral endoscopic management: what do we have? *Journal of Endourology*. 2014;28(5):502-8.
37. LOPES JF, SCHNED A, ELLSWORTH PI, CENDRON M. Histological analysis of urethral healing after tubularized incised plate urethroplasty. *The Journal of urology*. 2001;166(3):1014-7.
38. Karakus SC, Koku N, Parmaksiz ME, Ertaskin I, Kilincaslan H, Deliaga H. The effect of urethral catheter size on meatal stenosis formation in children undergoing tubularized incised plate urethroplasty. *Urology Journal*. 2014;10(4):1095-8