

Viral Load and asymptomatic virus status of the SARS-CoV-2 variants towards influencing the transmission

ABSTRACT

The Covid-19 virus (CoV) has created a worsened situation due to its transmissibility worldwide. Several millions of people have been infected since 2019 when it was first detected. Unpredictable impact and disruption created by the pandemic influence global financial or economic crisis. This motivated governments to encourage more vaccination to manage the spread and counter the pandemic, thereby breaking the chain. However, despite the level of vaccination against CoV, that has not been found to prevent an individual from being infected. Many transmission of SARS-CoV-2 occurs at the pre-symptomatic infection stage, which thus makes it difficult to contain.

This study aims to identify how critical the viral loads and asymptomatic virus status of the SARS-CoV-s influence the transmission of Omicron and how the quantitative SARS-CoV-s viral load assay corresponds to the clinical outcome while managing Covid-19 infected patients.

Keywords: Covid-19, SARS-CoV-2, Covid variants, asymptomatic status, Transmissibility

1. INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infections have resulted in an unprecedented global health pandemic of the 21st century. The Covid-19 virus (CoV) has thus created a worsened situation due to its transmissibility worldwide. Several millions of people have been infected since 2019 when it was first detected.

Since the evolution of since the virus was first discovered in the Wuhan region of China, various variants have thus emerged in terms of the characteristics and catastrophic impact they exhibit on infected individuals. SARS-CoV-2 is now diagnosed using the real-time reverse transcriptase polymerase chain reaction technique and reported using the binary assessments model for the test to indicate whether it is positive or negative.

Unpredictable impact and disruption created by the pandemic influence global financial or economic crisis. This motivated governments to encourage more vaccination to manage the spread and counter the pandemic, thereby breaking the chain. However, despite the level of vaccination against CoV, that has not been found to prevent an individual from being infected.

Many transmission of SARS-CoV-2 occurs at the pre-symptomatic infection stage, which thus makes it difficult to contain (1) (2). Unlike influenza, most SARS-CoV-2 usually do not transmit the virus, while just a small percentage of those transmit and infect a large population (1).

Several bodies of data have been provided to study the characteristics of those emerging CoV variants that have remained a major threat or concern to different regions worldwide. For instance, those emerging 'variants of concern' (VOC) possess unique viral characteristics that are being studied.

The goal is to understand the relationship with viral transmissibility, immune evasion, viral load, disease severity and lethality among the patients that are already infected—getting to understand the available scientific Evidence from both the clinical studies and animal models needed to be reviewed to understand the relationship between the viral load and covid-19 prognosis with vaccine efficacy.

In addition, the viral load dynamics (low Ct value indicates a high viral load) and kinetics among both the vaccinated and unvaccinated populations have been further studied using longitudinal follow-up data. However, because of the scarcity of such studies, many researchers have found this approach limited.

2. Methods

A literature search was undertaken using PubMed/Medline, Viruses, Frontiers, and PLOS ONE journal databases. A combination of search tags or keywords were used, including the SARS-CoVs viral load, Omicron viral load, COVID-19 viral load, quantitative viral loads, viral dynamics, Viral kinetics, associated with severity, transmissibility, household contacts, and asymptomatic covid-19 infection.

In addition, 22 publications relating to the research questions were collected and reviewed. Data on viral load, viral shedding, duration, variants, age, gender, and household contacts were all reviewed. A careful review of Omicron variants, sublineage, and other main variants was also done to achieve the objective.

3. RESULTS AND DISCUSSION

Isolation policies vital to managing Covid-19 were formulated based on the viral load dynamics and the dose-response model for the virus previously studied before the emergence of Omicron variants (3) (4). These might have influenced the number of studies currently published on the variant. Regardless, current data on the viral load, viral load shedding, the duration of viral load shedding, and several other co-founding factors have to be considered to evaluate the transmissibility and management of the variants (4).

The emerging variants of SARS-CoV-2 demonstrate diverse characteristics that major groups across the world have found to influence the viral transmissibility, viral load, disease severity, lethality and impact on vaccination or management of the patients. Most of the findings are results of the clinical studies or animal models focusing on linking the viral load to the prognosis and vaccine efficacy, especially among those already vaccinated, intending to compare to those not vaccinated. Therefore, it is understandable that longitudinal data on this topic is limited to viral load dynamics.

Variants of Concern (VOC) and viral load

A review publication by Silva et al. (5) emphasized that the emerging SARS-CoV-2 Variants of significant concern (VOC) have become a major issue. The role the RNA-dependent RNA polymerase played in introducing mutations in the viral genome, which eventually resulted in the emerging variant, was also highlighted (5). This was attributed to the average of 0.5 mutations which tend to accumulate over time during the infection cycle of each individual (5). As a result of the high transmission rate of the virus over the globe, it can be concluded that the generation of mutations and new variants tends to rise (5).

The eventual result of such is the continuity of the pandemic, given the evolution of the SARS-CoV-2. Currently, more than five variants have been identified and categorized according to their transmissibility or severity. Among the identified variants, Omicron (B.1.1.529) was the fifth to be discovered, and it was first reported in Botswana and immediately followed by South Africa, both in November 2021 (5) (6). To show how dangerous the new variants were to the world, the rate at which they spread to 6 continents within 30 days of the initial discovery. The rapid rate at which those emergent variants spread around the population accumulates widespread attention with Evidence for the transmission or clinical implications (6).

According to the randomized controlled trial study by Marc et al. (7), a relationship between the viral load and studied variants on disease transmission was projected. Considering the current knowledge and findings, it was shown that the VOCs' viral load tends to increase by 2-8 fold but with no major changes in terms of the pattern of contacts across the variants. It was also noted that the viral load level among the variants could be linked with a relative increase in the 'probability of transmission of about 24-58% for household contacts but 15-39% for non-household contacts (7).

Researchers found Omicron to be heavily mutated variants with 30 amino acid substitutions, some deletions involving six residues, and insertions of 3 residues of the spike protein. Those were found to be concentrated around the receptor binding motif. In addition, some insertions were not identified when the variants were first identified but have been later identified among the mutations to be the insertion (ins214EPE) in spike. For the Omicron, scientists have also hypothesized that the identified insertion could easily be acquired as a form of template switching, which entails the genome of a lower pathogenic coronavirus. Especially those coronaviruses found to have the capability to cause the common cold (HCoV-229E). Omicron sublineages have emerged with increasingly higher replication

advantages over the other previously identified predominant sub lineages. The first Omicron identified was the BA.1. sub lineage while others followed (6). Several Omicron sub lineages have been found to have replication advantages over the other Delta variants. They also evade the infection and vaccine induced humeral immunity to a greater extent than what has been noticed with prior variants (6).

However, in terms of association with severe diseases, the omicron sublineages appear to be associated less. Observational data have shown that the risk of severe illness or death with Omicron is lower than with other variants (6).

In a quest to get appropriate data, Van Der Veer et al. (8) implemented infection prevention guidelines for their healthcare workers and necessary testing conducted to evaluate the Ct-value and variant determination via whole genome sequencing with the required genotyping (8). As a result, the researchers in the longitudinal study found the healthcare workers' viral load dynamics to be different, especially between the three main variants studied (Delta, BA 1. and BA 2. variants).

However, the BA 2. Variants were found with the highest viral load within the first 10 days after the initial PCR was collected, although the infection duration among the three variants appears not to be different (8). The findings further provide a better understanding of the dynamics of BA 2, especially concerning the transmission and current quarantine rules, which are majorly formulated based on the earlier understanding of the other variants (8). For instance, quarantine rules are based on viral load found during the infection duration, which targets the individual from the dimension that such an individual will have to be asymptomatic for at least one day during the quarantine days is 5 and 7 days for healthcare workers (8).

To replicate previous findings from earlier studies on Omicron and subsequent sublineages, Dewald et al. (9) determined the viral dynamics of breakthrough omicron infections; both the viral load of the vaccinated and bolstered individuals were measured (9). The first 3 days of onset of symptoms showed significantly higher viral load Ct 21.76 for the vaccinated and 23.14 for boosted individuals (9).

Longitudinal studies were further conducted on the group of patients for 14 days to understand the difference in the viral load trajectory (9). Findings indicate that booster immunization reduced the detectable viral loads without changing the main viral load dynamics for the period, unlike in vaccinated individuals (9). However, the findings of this study appear to be totally different from what Bouton et al. (10) found from a 10-day longitudinal study that has shown that booster vaccination does not have any impact on the virus kinetics but against infection because it reduces transmission (10).

Van Der Veer et al. (8) also identified the various viral loads level of the three studied variants for the five days after the first PCR was done. For instance, BA 2 viral loads (5.7 log₁₀ copies/mL [Ct-value 21]) appears to be highest among the three followed by that of BA1 (4.8 log₁₀ copies/mL [Ct-value 24]) and then Delta variants (4.3 log₁₀ copies/mL [Ct-value 26]). However, based on their findings, the viral load changes occur with BA 2 having a Ct value of 26, equal to Delta on day 5. The detected loads were considered to be representing the intact viral particles in both the vaccinated and unvaccinated persons. Based on the findings, Delta, BA1, and BA 2 were noted to have different viral load dynamics. BA 2 has the highest of the 3 in terms of viral loads. This is considered a strong influencer of the transmission potential despite following the suggested quarantine period (8).

SARS-Cove infections begin in the respiratory tract before spreading to other organs. Therefore, the isolation of viral nucleic acids from those tissues, blood and body secretions could easily help analyse the systemic spread and determine the severity of the infection (11). In addition, a persistent increased viral load from reviewed data explains the correlations between the increased infectivity and rapid transmission, making it easier to assess viral load from the respiratory tracts to determine the risk of severity (11).

According to scientists, high viral load strongly correlates with elevated cytokine and lymphopenia, which are the main markers of inflammation and portrays a poor prognosis (11). When comparing the duration of SARS-CoV-2 within the stool, serum and sputum sample. It was found that virus stay longer in stool samples compared to others. This means that management of stool needs to be thoroughly considered while planning for prevention and control of the epidemic. However, in the respiratory tissues of patient with more severe diseases, It was found that the virus could persist longer (6).

Infectious Viral Load, Age, Sex and community transmission

SAR-CoV infections occur almost anywhere. Household contact exposure remains a point studied by Singanayagam et al. (12) to understand community transmission and viral load kinetics; The authors focused on the delta variants concerning vaccinated and unvaccinated individuals (12). Researchers noted peak viral loads do not differ in terms of vaccination status or variant type but tend to increase modestly with age, with a difference of 0.39 in peak \log_{10} viral load per mL between those at the age of 10 years to those at 50 years (12).

Another group of research, Puhach et al. (13), tested the correlations between the infectious viral loads, age and sex among vaccinated and unvaccinated groups of individuals. It was evident that no correlation existed between age and infectious viral load among the four groups studied. Similarly, no correlation existed in terms of the infectious viral loads between the male and female patients for those infected with pre-VOC, Delta (which are fully vaccinated or unvaccinated) or Omicron BA 1, which is fully vaccinated only. This is unlike what was founded by Singanayagam et al., (2022), which observed a modest difference in correlations.

It was also observed that for these delta variants, the viral load had a much faster decline mean rate in fully vaccinated individuals compared to those unvaccinated individuals with pre-alpha (0.69), alpha (0.82), or Delta (0.79) variant infections (12). The dynamics with regards to the viral growth show that individuals with faster viral load growth tend to have higher peak viral load correlation, i.e. correlation 0.42 [95% credible interval 0.13-0.65] but with a much slower decline rate (-0.44 [-0.67 to -0.18]) (12). Those results show the impact of vaccination on the viral load, reduced risks of infection of the delta variants and helped in viral clearance (12).

However, there are still several instances where fully vaccinated individuals with breakthrough infections have peak viral loads similar to that seen in unvaccinated ones, which indicate that such group of individuals can transmit infection within their homes or community is high. The understanding provided by the study by Singanayagam et al. (12) shows that the type of host-virus interactions that occur during the early period of infections strongly influences the entire viral trajectory through the course of infection (12).

The relationship between viral load and transmissibility using epidemiological and clinical data was tested by Kawasuji et al. (14). The index patient (the patient that first transmits the disease to at least another) and the non-index (the patient that causes secondary transmission). High Nasopharyngeal Viral load data at the onset provides a better understanding of why some patients easily transmit infection, and others do not (15). This

attributes to the findings regarding the median viral load relating to the initial sample in a symptomatic patient, which is higher than that in asymptomatic patients or higher in adults compared to children (15). Viral load at onset was higher among the symptomatic patients considered to be index than those non-index patients (15), (7). Regardless of whether either symptomatic or asymptomatic, median viral loads maintain a high value in adult patients in the index compared to non-index patients (15).

In addition to those findings, Meyerowitz et al. (16) also noted that having a high viral load associated with the index case needed to be considered vital to super-spreading events, especially during transmission. However, the importance of other specific host factors' contribution to the events is yet unknown. Further study also highlighted how the secondary attack rate is associated with the increased severity of the index case and related symptoms of fever and expectoration, which are known with secondary infections(16).

A study by Darlene Bhavnani et al. (17) found that 77% of the transmission events were associated with the case viral loads of $\geq 1 \times 10^5$ copies of viral RNA per mL of saliva (17). In view of the role played by viral shedding in transmission, if cough actually serve as a consequence of higher viral loads was questioned by Darlene Bhavnani et al. (17) if it actually perpetuate the transmission independent of viral load. However, their study still tries to demonstrate the existence of relationship between the viral load at the time of test with presence of cough on the day of exposure or the risk that is present in terms of transmission if cough occur on the day of exposure (17). These are major questions that is yet to be answered but still linked to the viral loads.

In the study by Cathinka et al. (18), focusing on the household transmission, Secondary attack rate was found to be higher in the Alpha variant (78%) when compared to the non-VOC variants (43%). In addition, those cases were found to be associated with higher viral load (18). Unlike the study by Meyerowitz (16), Cathinka et al (18) found that children had lower viral loads and were usually more asymptomatic than adults. However, secondary attack rate was noted to be higher among household contacts that are above 40 years of age (69%) compared to younger contacts (18).

Pre-symptomatic transmission

Tindale et al., (2020) described the rate of novel coronavirus transmission as 40-80%, which usually happens 2-4 days before the onset of symptoms in an infected patient. This type of transmission has made it challenging to manage the pandemic because of the pre-symptomatic infection's difficulty in containing the spread (1). In addition, studies have shown viral load to be at a high level during the pre-symptomatic period and then peak during the onset of symptoms but later declines over time (2), (7), (11). According to Tindale et al. (19), most of those transmissions which occur in healthy individuals depict that home lockdown or isolation when an individual shows symptoms is not enough to control the spread of SARS-CoV-2 (19).

Several other measures are needed, such as broad-scale social distancing, which has proven to help prevent asymptomatic transmission of Covid-19. The over-dispersion is usually described as an instance whereby most infected individuals don't transmit at all, and only a few per cent of the infected population remains to be the super-spreader. Another study by Goyal et al. (1), it was described that the viral load in the upper respiratory airway to help determine the likelihood of transmission. This was more of a reason because viral load at such upper respiratory tract was found to be highest during the early phase of the disease but later increases downward to the lower respiratory tract (sputum) (16). This also suggests that more viral replications occur at such regions or sites and thus shift to the lower region (16).

The findings by Goyal et al. (1) indicate that exposure to the airway viral load that is shedding determines the probability of transmission of the infection (1). This other study by Ra et al. (20) indicated that 1/5th of persons without severe symptoms of Covid-19 were asymptomatic. The viral load of the population is comparable to those found in symptomatic patients. It was also highlighted by the notable findings in a large proportion of mildly symptomatic individuals with Covid-19 or those that are asymptomatic with SARS-CoV-2, which is the existence of persistent positive upper respiratory tract RT-PCR results at follow-up (20). This asymptomatic population were indicated, according to the authors, to exhibit the viral load in the nasopharynx of the patients was comparable to what was found in those with mild symptoms. They emphasized the dangers of such asymptomatic individuals with SARS-CoV-2 infection, which are considered a major contributor to the ongoing community Covid-19 pandemic.

A study by Puhach et al. (13) quantified the infectious viral load in those infected with the covid-19 during the first five days of showing symptoms utilizing the in-vitro culturability assay in both that are either vaccinated or unvaccinated. The identified individuals are infected with the pre-variant of concern (pre-VOC), including the SARS-CoV-2, Delta or Omicron BA1 (13). Unvaccinated individuals with the pre-VOC SARS-CoV-2 were found to have lower infectious viral load compared to those unvaccinated individuals infected with Delta variants (13).

On the other hand, Omicron BA 1 in these breakthrough cases was found to have reduced infectious viral load only in individuals with boosted and not full vaccinated individuals compared to other groups. Those other groups include the fully vaccinated Omicron BA. 1-infected persons show lower infectious viral load compared to fully vaccinated Delta variants (13). The implication is that increased infectious viral load determines the infectiousness of the SARS-CoV-2 Omicron BA.1. variants (13). However, vaccination has its role in lowering the transmission risks, which is pertinent for the public health protection of citizens or healthcare workers (13).

The relationship between the SARS-CoV-2 viral load and infectiousness, which is poorly known in the academic world, was explored by Marc et al. (7). The viral load and probability of infection were analysed to understand the effect of the viral load (7). The noted transmission probability was found to be large, around 48% among the household contacts compared to non-household contacts, especially when the viral load was found to be more significant than 10^{10} copies/mL (7). Similarly to the viral load level at the symptom onset noted by many scientists, the transmission probability was found to peaked at that exact point (7) but with a mean probability of transmission rate of 29%, although with large individual variations (7).

Another cohort study on the SARS-CoV-2 G614 variant infection found a form of rapid viral load peak occurring at Covid-19 symptom onset, followed by a slower decline but correlated positively with the severity of the symptoms (21). The longitudinal evaluation conducted by Stankiewicz Karita et al. (21) on the SARS-CoV-2 G614 utilizing the molecular testing mode allows such to serve as the point of reference for future comparison for the emergent viral lineage that could be discovered. This is vital for future clinical trials and the development of public health strategies or policies to contain the virus (21). It was noted that patient with omicron Infections are more likely to be vaccinated compared to those with Delta. However, they were less likely to be admitted or succumb to infection regardless of the vaccination status. A booster dose, however, influences the recovery of infectious virus in the cell culture compared to the fully vaccinated that has no booster or unvaccinated that is infected with delta variants (22).

Apart from the viral load level based on gender or sex which no correlates were found, the relative probability of transmission and the relationship with age and gender is still yet unclear. However, Meyerowitz et al. (16) found the replication-competent virus to be readily isolated from children compared to adults, especially when the onset of symptoms is not a key determinant of the viral load (16). This is simply because the study findings indicate the higher viral load to be more linked with the increased likelihood of infectious viral isolates. About eight days is considered to be the median after symptoms onset for such isolation to be possible. However, the probability for the isolation declined to about less than 5% post 15.2 days while it further decreased over time.

4. CONCLUSION

At present, reviewed literature has no unified agreement on the implication of quantitative SARS-CoV-2 viral load on the severity of infection. Therefore, the study aims to identify how critical the viral loads and asymptomatic virus status is to SARS-CoV transmission with a focus on Omicron or the importance of the viral load assay to the corresponding clinical outcome of the infection. Several current studies have been critically reviewed to understand viral load dynamics and transmission probability better. However, very few studies focused on Omicron. Emphasis on the VOC's viral load relationship on the transmission probability was considered to differ between household and non-household contacts. The viral loads were on the rise with about 2-8 folds, although the contact pattern has not differed across the variants. Despite the difference in viral load noted by some researchers among the variants and their sublineage, most findings do not note any noticeable difference in terms of the infection duration. Regardless, quarantine rules are still formulated based on the viral load found during the infection duration.

Further studies have explored the impact of vaccination and booster immunizations' on viral loads and implications on the transmission rate. Most of the findings show this to be a positive correlation. However, several exceptions had no relationship between the factors and considered the viral load between unvaccinated and vaccinated to be no significant difference. Based on the reviews, viral load appears to peak at the onset of infection; detected loads represent the intact viral particles regardless of vaccination status. The viral load has a degree of influence on the transmission potential irrespective of the quarantine period, and an increase in viral load positively correlates with the rate of infectivity and rapid transmission regardless of gender. There is a further correlation between the viral load level and index case and the existence of super-spreading events, although not yet agreed upon by a lot of researchers; hence still needs further investigation.

REFERENCES

1. Goyal A, Reeves DB, Fabian Cardozo-Ojeda E, Schiffer JT, Mayer BT. Viral load and contact heterogeneity predict sars-cov-2 transmission and super-spreading events. *Elife*. 2021 Feb 1;10:1–63.
2. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med* [Internet]. 2020 May 1 [cited 2022 Oct 11];26(5):672–5. Available from: <https://pubmed.ncbi.nlm.nih.gov/32296168/>
3. Watanabe T, Bartrand TA, Weir MH, Omura T, Haas CN. Development of a Dose-Response Model for SARS Coronavirus. *Risk Anal* [Internet]. 2010 Jul 1 [cited 2022 Oct 11];30(7):1129–38. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1539-6924.2010.01427.x>
4. Wu C, Chen X, Cai Y, Jia'an Xia ;, Zhou X, Xu S, et al. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China Supplemental content. *JAMA Intern Med* [Internet]. 2020;180(7):934–43. Available from: <https://jamanetwork.com/>
5. Silva SJR da, Lima SC de, Silva RC da, Kohl A, Pena L. Viral Load in COVID-19 Patients: Implications for Prognosis and Vaccine Efficacy in the Context of Emerging SARS-CoV-2 Variants. *Front Med*. 2022 Jan 31;8:3111.
6. Kenneth McIntosh, Martin Hirsch, Allyson Bloom. COVID-19: Epidemiology, virology, and prevention [Internet]. *UpToDate*. 2022 [cited 2022 Oct 12]. Available from: <https://www.uptodate.com/contents/covid-19-epidemiology-virology-and-prevention#H2895385127>
7. Marc A, Kerioui M, Blanquart F, Bertrand J, Mitjà O, Corbacho-Monné M, et al. Quantifying the relationship between sars-cov-2 viral load and infectiousness. *Elife*. 2021 Sep 1;10.
8. Van Der Veer BMJW, Dingemans J, Bank LEA, Von Wintersdorff CJH, Van Loo IHM, Savelkoul PHM. Viral load dynamics in healthcare workers with COVID-19 during Delta and Omicron era. 2022 [cited 2022 Oct 4]; Available from: <https://orcid.org/0000-0003-4409-4611>
9. Dewald F, Detmer S, Pirkl M, Hellmich M, Heger E, Herrmann M, et al. Viral Load Dynamics in SARS-CoV-2 Omicron Breakthrough Infections. *J Infect Dis* [Internet]. 2022 Jul 30 [cited 2022 Oct 4]; Available from: <https://academic.oup.com/jid/advance-article/doi/10.1093/infdis/jiac290/6652280>
10. Bouton TC, Atarere J, Turcinovic J, Seitz S, Sher-Jan C, Gilbert M, et al. Viral dynamics of Omicron and Delta SARS-CoV-2 variants with implications for timing of release from isolation: a longitudinal cohort study. *medRxiv* [Internet]. 2022 Apr 5 [cited 2022 Oct 4]; Available from: [/pmc/articles/PMC8996632/](https://pmc/articles/PMC8996632/)
11. Shenoy S. SARS-CoV-2 (COVID-19), viral load and clinical outcomes; lessons learned one year into the pandemic: A systematic review. *World J Crit Care Med* [Internet]. 2021 Jul 7 [cited 2022 Oct 4];10(4):132. Available from: [/pmc/articles/PMC8291003/](https://pmc/articles/PMC8291003/)

12. Singanayagam A, Hakki S, Dunning J, Madon KJ, Crone MA, Koycheva A, et al. Community transmission and viral load kinetics of the SARS-CoV-2 delta (B.1.617.2) variant in vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study. *Lancet Infect Dis* [Internet]. 2022 Feb 1 [cited 2022 Oct 4];22(2):183–95. Available from: <http://www.thelancet.com/article/S1473309921006484/fulltext>
13. Puhach O, Adea K, Hulo N, Sattoune P, Genecand C, Iten A, et al. Infectious viral load in unvaccinated and vaccinated individuals infected with ancestral, Delta or Omicron SARS-CoV-2. *Nat Med* 2022 287 [Internet]. 2022 Apr 8 [cited 2022 Oct 4];28(7):1491–500. Available from: <https://www.nature.com/articles/s41591-022-01816-0>
14. Kawasuji H, Takegoshi Y, Kaneda M, Ueno A, Miyajima Y, Kawago K, et al. Transmissibility of COVID-19 depends on the viral load around onset in adult and symptomatic patients. *PLoS One* [Internet]. 2020 Dec 1 [cited 2022 Oct 4];15(12):e0243597. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0243597>
15. Kawasuji H, Takegoshi Y, Kaneda M, Ueno A, Miyajima Y, Kawago K, et al. Transmissibility of COVID-19 depends on the viral load around onset in adult and symptomatic patients. *PLoS One* [Internet]. 2020 Dec 1 [cited 2022 Oct 4];15(12). Available from: <https://pubmed.ncbi.nlm.nih.gov/33296437/>
16. Meyerowitz EA, Richterman A, Gandhi RT, Sax PE. Transmission of SARS-CoV-2: A Review of Viral, Host, and Environmental Factors. <https://doi.org/10.7326/M20-5008> [Internet]. 2020 Sep 17 [cited 2022 Oct 4];174(1):69–79. Available from: <https://www.acpjournals.org/doi/10.7326/M20-5008>
17. Bhavnani D, James ER, Johnson KE, Beaudenon-Huibregtse S, Chang P, Rathouz PJ, et al. SARS-CoV-2 viral load is associated with risk of transmission to household and community contacts. *BMC Infect Dis* [Internet]. 2022 Dec 1 [cited 2022 Oct 4];22(1):1–11. Available from: <https://bmcinfectdis.biomedcentral.com/articles/10.1186/s12879-022-07663-1>
18. Julin CH, Robertson AH, Hungnes O, Tunheim G, Bekkevold T, Laake I, et al. Household transmission of sars-cov-2: A prospective longitudinal study showing higher viral load and increased transmissibility of the alpha variant compared to previous strains. *Microorganisms* [Internet]. 2021 Nov 1 [cited 2022 Oct 4];9(11):2371. Available from: <https://www.mdpi.com/2076-2607/9/11/2371/htm>
19. Tindale LC, Stockdale JE, Coombe M, Garlock ES, Lau WYV, Saraswat M, et al. Evidence for transmission of COVID-19 prior to symptom onset. *Elife* [Internet]. 2020 Jun 1 [cited 2022 Oct 11];9:1–34. Available from: <https://pubmed.ncbi.nlm.nih.gov/32568070/>
20. Ra SH, Lim JS, Kim GU, Kim MJ, Jung J, Kim SH. Upper respiratory viral load in asymptomatic individuals and mildly symptomatic patients with SARS-CoV-2 infection. *Thorax* [Internet]. 2021 Jan 1 [cited 2022 Oct 4];76(1):61–3. Available from: <https://thorax.bmj.com/content/76/1/61>
21. Stankiewicz Karita HC, Dong TQ, Johnston C, Neuzil KM, Paasche-Orlow MK, Kissinger PJ, et al. Trajectory of Viral RNA Load Among Persons With Incident SARS-CoV-2 G614 Infection (Wuhan Strain) in Association With COVID-19 Symptom Onset and Severity. *JAMA Netw Open* [Internet]. 2022 Jan 4 [cited 2022 Oct 4];5(1):e2142796–e2142796. Available from: <https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2787768>
22. Fall A, Eldesouki RE, Sachithanandham J, Morris CP, Norton JM, Gaston DC, et al. The displacement of the SARS-CoV-2 variant Delta with Omicron: An investigation of

hospital admissions and upper respiratory viral loads. eBioMedicine [Internet]. 2022 May 1
[cited 2022 Oct 4];79:104008. Available from:
<http://www.thelancet.com/article/S235239642200192X/fulltext>

UNDER PEER REVIEW