

# Susceptibility and Resistance Patterns of Bacteria Isolated from Infected Wounds, Burn, Medical Device Tips and Blood to Doxycycline and Septrin from Patients in the city of Benghazi

## Abstract:

Wound infections and septicemia with drug-resistant bacteria lead to higher mortality and morbidity and increased healthcare costs. We aimed to evaluate the prevalence rate of nosocomial bacterial agents with antimicrobial susceptibility patterns in hospitalized and non-hospitalized patients. Samples are collected from the wound pus, burn; medical device tip and blood of male and female patients and are sent for culture and sensitivity to the Microbiology Department of Al Saleem medical laboratory, Benghazi, Libya, from October 2021 to January 2022. Collected blood samples were directly inoculated into brain heart infusion (BHI) broth, while various swabs were collected by Transport medium and both are cultured on Blood agar, Chocolate agar, MacConkey - aerobic and anaerobic blood agar at 35°C, with CO<sub>2</sub>. The wound pus was the most samples from which bacteria were isolated, followed by medical device tip swabs, and the least isolation was from a blood specimen. 47 of the specimens were culture-positive. Of these *Staph aureus*, *Strep pneumonia* and *E. coli* were the most abundant bacterial identified in clinical samples. According to susceptibility testing results, antibiotic resistance patterns of Gram-negative bacteria showed that the highest resistance rate was against Septrin (74.4%) and the lowest rate, 17.8%, was related to Doxycycline (44.6%) respectively. The isolates from the nosocomial infection in these patients are resistant to Septrin and Doxycycline antibiotics. We recommend that whenever the diagnosis of nosocomial infection is made, a more effective antibiotic treatment be instituted until the susceptibility of the strain is identified.

**Keywords:** Doxycycline, Nosocomial infection, Pus swab, *Staph aureus*.

## Introduction

Antibiotic resistance, and the extent to which bacteria are able to resist drugs manufactured to kill them, is one of the biggest challenges to public health today. Antibiotics are among the most importance tools that used in the fight against life-threatening infections. Nosocomial infections have called (hospital-acquired infections) and refer to infections that occur within 48 hours of a patient's admission to hospital, three days of a patient's discharge from hospital, or within 30 days of an operation (Inweregbu *et al.*, 2005; Khan *et al.*, 2015).

Moreover, this infection includes occupational injuries among medical staff in medical facilities (WHO, 2016), as well as invasive devices such as ventilators used in modern health care and catheters are associated with this infection (CDC, 2016). A wound can be defined as any damage or injury to the skin that limits its protective function. This acute damage generally caused by external injuries to the skin, such as sharp cuts, minor cuts, insect and animal bites, burns caused by heat, friction, cold, chemicals, lacerations, puncture wounds, and surgical incisions (Siddiqi and Bernstein, 2010) as well as the presence of necrotic tissue in the skin. Wound - this can aid the growth of bacteria, especially anaerobic (Hess, 2011; Inweregbu *et al.*, 2005) All exposed

skin wounds are colonized by bacteria, but this does not mean that all wounds are infected.

As infections may occur in wounds during healing, regardless of whether they are infected or not, a certain level of swelling and increased warmth at the wound site is normal and should not be confused with a clinical infection. When the skin cracks, its protective defense mechanisms become weak, and the environment prepared to become more conducive to infection with bacteria and thus their number increases.

These bacteria come from three essential sources; the environment (such as dirt, dust, foreign bodies, and bacteria on clothing, hands, and medical equipment). The surrounding skin (where normal skin contains bacteria, referred to as symbionts) and the mucous membranes (the digestive system, mouth, and urogenital system) (Siddiqi and Bernstein. 2011). Septicemia occurs when the infection spreads to all areas of the body via the bloodstreams and causes symptoms such as chills, fever, and tachycardia (Edwards and Harding. 2004; Siddiqui and Bernstein. 2011).

Several studies have indicated that adult bloodstream infection belongs to microorganisms of the family Enterobacteriales (*Klebsiella* spp and *Escherichia coli*) and *Enterococcus* spp (Daneman *et al.*, 2018).

Other studies have also indicated that most bacterial infections are caused by Gram-positive bacteria coagulase-negative *staphylococci* (CoNS), *Staphylococcus aureus* and *Enterococcus faecalis* (13%) (Lapeña, Jose. 2013). Thus, the study aims to determine the association of factors Bacteremia in hospitals with a pattern of antimicrobial susceptibility in inpatients and non-resident patients.

## **Materials and Methods**

### **2.1 Study area**

This cross-sectional study was conducted at Microbiology Department of Al-Saleem laboratory, Benghazi, Libya, from October 2021 to January 2022. The samples assembled from the wounds pus, burn; medical devices, tips and blood from male and female patients in the age group of 3 to 86 years with clinical characteristics and septicemia suspected to be infected.

### **2.2 Sampling**

All swabs specimens were collected aseptically and carried from patients by Stuart's, Amies. Collected blood samples were directly inoculated into brain heart infusion (BHI) broth with the blood to broth ratio of 1:10 for up to 5 days and were immediately transported to the microbiology department. Age, sex, health status and use of medicines, infection type were recorded for all included patients.

### **2.3 Blood cultures**

Positive culture samples were directly inoculated onto blood agar, chocolate agar, and MacConkey agar and incubated overnight at 37°C aerobically. Blood agar and Chocolate plates were incubated at 35 °C in microaerophilic conditions (containing 5% CO<sub>2</sub>).

Moreover, 0.5 to 1 mL of the swab specimen was inoculated into a tube of enriched thioglycolate broth (Lapeña, 2013; Noor-alhoda *et al.*, 2019).

## **2.4 Determination of bacterial isolates**

Visual investigation of the bacterial colonies shows culture positivity. The supposed organism is then defined by Gram's stain and a series of biochemical tests, including an oxidase test, catalase test, Carbohydrate utilization test, Citrate utilization test, DNAase test, urease test, coagulase test, Analytical profile index (api) 20e test and Mannitol Salt Agar. (Alawkally *et al.*, 2022; Alawkally *et al.*, 2019; Fakron *et al.*, 2022; Lapeña and Jose, 2013; Sweileh, 2009).

### **2.4.1 Direct Vitek 2 AST**

The supposed strains' morphologically similar colonies were determined by biochemical assays utilising the Vitek system (bioMerieux, Hazelwood, MO). In brief, the cell density of the remaining bacterial suspension was modified to a density of 0.5 McFarland after dilution in 0.45% saline; 145µL of the bacterial suspension was drawn into 3 mL of 0.45% saline solution to further adjust the bacterial cell density. Vitek cards were inoculated with suspension vials and loaded into the Vitek 2 automated reader-incubator. Vitek cards AST-GN13, AST-Gp67, and AST-Gp68 were utilized for gram-negative bacteria, *staphylococci/enterococci/ streptococci* and *Strep pneumoniae*, respectively (Tian *et al.*, 2016).

## **2.5 Antimicrobial susceptibility testing**

Susceptibility of bacterial isolates to antibacterial drugs was carried out by the disk diffusion method to the following antibiotics: Doxycycline- DO (30 µg) and Seprine-SXT (25 µg). Analysis of the results was carried out following the recommendations of CLSI. 2018; Kassam *et al.*, 2017. In brief, 4–6 morphologically similar bacteria colonies from clinical samples were collected with an inoculating loop and relocated into a tube including 5 mL of broth culture, gradually and carefully mixed until a relatively homogenous suspension was obtained, and incubated at 37°C. A sterile non - toxic dry cotton swab was utilized to streak a sample of the centralized inoculation on the Mueller–Hinton agar plate. With the lid on, the inoculums were allowed to dry for 5–15 minutes. The antibiotic disks were distributed to the plates with sterile forceps at a distance of 15 mm from the edge and 24 mm apart from each other. After 24 hours of incubation at 37°C, the diameters of the zones of bacterial growth inhibition around the disks were measured to the nearest millimetre. The isolates were classified as sensitive (S), or resistant (R) utilising a standardized table. The inhibition zone of a ruler was measured in millimetres (CLSI, 2010).

The antibiotics tested were Septrin (300 µg; Resistant ≤12; Sensitive ≥17) and Doxycycline- DO (30 µg; Resistant ≤12 ; Sensitive ≥16).

## **2.6 Data Analysis**

The data was analyzed by SPSS programs version 20.

### 3. Results

#### 3.1 Distribution table of different Wound pus and blood Infection by genders

The gender distributions of the patients with swab infection are reported in Table. 1 Concerning gender, the swab infection rate was higher in females than in males.

**Table. 1** Distribution table of different Wound pus and blood Infections by genders

Sex	Frequency	Percent
Female	25	53.2
Male	22	46.8
Total	47	100.0

#### 3.2 Distribution of positive cases by age groups.

Regarding age distribution, most of the positive patients were identified in the 45-65 age groups, followed by the 24-44 age groups.

**Table. 2** Distribution of positive cases by age groups.

Age	Frequency	Percent
3-23 years	11	23.4
24-44 years	14	29.8
45-65 years	15	31.9
66-86 years	7	14.9
Total	47	100.0

#### 3.3 Distributions of bacterial infection in wound pus, burn, medical device tips and blood specimens Collected from Patients.

Overall, Gram-positive organisms *Staphylococcus aureus*, *Streptococcus pneumonia*, *Staphylococcus epidermidis*, *Enterococcus faecalis* and *Strep agalactia* equally and *staph saprophyticus* and *Step pyogen* equally and accounted for most of the isolates 26 (55.3%). Gram-negative bacteria constituted about 21 (44.6%) of isolates. Among the Gram-negative isolates identified, *Klebsiella* spp (*klebsiella oxycota* and *Klebsiella pneumonia* ) was the predominant isolate, followed by *E. coli* with an isolation rate of 19.1% and 10.6% respectively.

**Table. 3** Distributions of bacterial infection in wound pus, burn, medical device tips and blood specimens Collected from Patients.

Bacteria	Frequency	Percent
<i>Acinetobacter baumannii</i>	2	4.3
<i>Citrobacter freundii</i>	1	2.1
<i>E. coli</i>	5	10.6
<i>Enterococcus faecalis</i>	2	4.3

<i>Serratia marcescens</i>	1	2.1
<i>Klebseilla pneumoniae</i>	3	6.4
<i>klebseilla oxycota</i>	6	12.8
<i>proteus mirabilis</i>	2	4.3
<i>pseudomonas aeruginosa</i>	1	2.1
<i>staph aureus</i>	10	21.3
<i>Staph epidermidis</i>	3	6.4
<i>staph saprophyticus</i>	1	2.1
<i>Step pyogen</i>	1	2.1
<i>strep pneumoniae</i>	7	14.9
<i>Strep agalactia</i>	2	4.3
<b>Total</b>	47	100.0

### 3.4 Frequency of isolated pathogens from each nosocomial infection.

The wound pus was the most samples from which bacteria were isolated, followed by medical device tip swab, and least isolation was from blood specimen. *klebseilla* spp and *Staph aureus* were the most prevalent pathogens of wound pus contagion respectively. A total of 7 patients with suspected septicemia were studied. The most frequent isolates were gram positive cocci (12.7%). *Pseudomonas aeruginosa* spp 1 (2.1%) was also fairly frequently identified.

**Table 4.** Frequency of isolated pathogens from each nosocomial infection.

Isolates	specimen	Frequency	Percent
<i>Acinetobacter baumannii</i>	Wound pus	2	4.3
<i>Citrobacter freundii</i>	Wound pus	1	2.1
<i>E. coli</i>	Medical device tips	2	4.3
<i>E. coli</i>	Burn	1	2.1
<i>E. coli</i>	Burn	2	4.3
<i>Enterococcus faecalis</i>	Wound pus	2	4.3
<i>Serratia marcescens</i>	Wound pus	1	2.1
<i>Klebseilla pneumoniae</i>	Medical device tips	2	4.3
<i>Klebseilla pneumoniae</i>	Burn	1	2.1
<i>klebseilla oxycota</i>	Wound pus	5	10.6
<i>Klebseilla oxycota</i>	Wound pus	1	2.1
<i>Proteus mirabilis</i>	Wound pus	2	4.3
<i>pseudomonas aeruginosa</i>	Blood	1	2.1
<i>staph aureus</i>	Blood	2	4.3
<i>staph aureus</i>	Burn	1	2.1
<i>staph aureus</i>	Wound pus	5	10.6
<i>staph aureus</i>	Medical device tips	2	4.3
<i>Staph epidermidis</i>	Blood	2	4.3
<i>staph saprophyticus</i>	Wound pus	1	2.1

<i>Staph epidermidis</i>	Wound pus	1	2.1
<i>Step pyogen</i>	Wound pus	1	2.1
<i>Strep pyogen</i>	Medical device tips	1	2.1
<i>strep pneumonia</i>	Burn	1	2.1
<i>strep pneumonia</i>	Blood	2	4.3
<i>Strep pneumonia</i>	Wound pus	4	8.5
<i>Strep agalactia</i>	Wound pus	1	2.1
<b>Total</b>	<b>47</b>	<b>47</b>	<b>100.0</b>

### 3.5 Antibiotic sensitivity, resistance and intermediate sensitivity of bacteria isolated from wound pus, medical device tip, burn and blood culture to Septrin and Doxycycline

In the present study, the antimicrobial resistance profile of isolates had been analyzed. Almost isolated strains showed a high ratio of resistance to the Septrin than Doxycycline.

**Table. 5** Antibiotic sensitivity, resistance and intermediate sensitivity of bacteria isolated from wound pus, medical device tip, burn and blood culture to Septrin

SXT			DO	
Susceptibility patterns	Frequency	Percent	Frequency	Percent
<b>R</b>	35	74.5	21	44.7
<b>S</b>	12	25.5	26	55.3
<b>Total</b>	<b>47</b>	<b>100.0</b>	<b>47</b>	<b>100.0</b>

**Note:** R- resistant; S-Sensitive

### 3.6 Resistance pattern of isolated pathogens to evaluated antibiotics.

All the bacterial isolates were tested for susceptibility against Septrin. Out of 47 bacterial isolates, 12 (25.5%) of the isolates remain susceptible to Septrin. While 35 (74.4%) were found to be resistant to Septrin. The susceptibility forms of the bacterial isolates to Gentamycin antibiotics are offered in Table 7. From the outcomes, 26 (5.3%) isolates were most sensitive to Doxycycline. The rate of resistance of isolates to Doxycycline antibiotics was 44.6%. The action of Doxycycline against the isolates was somewhat acceptable.

**Table. 6** Resistance pattern of isolated pathogens to evaluated antibiotics

Isolates	SXT Susceptibility		Total	Do Susceptibility		Total
	R	S		R	S	
<i>Acinetobacter baumannii</i>	2	0	2	1	1	2
<i>Citrobacter freundii</i>	0	1	1	0	1	1
<i>E. Coli</i>	4	1	5	1	4	5

<i>Enterococcus faecalis</i>	2	0	2	1	1	2
<i>Serratia marcescens</i>	1	0	1	0	1	1
<i>Klebseilla pneumoniae</i>	1	2	3	2	1	3
<i>Staph epidermidis</i>	1	2	3	1	2	3
<i>Step pyogen</i>	1	0	1	0	1	1
<i>Strep agalactia</i>	2	0	2	2	0	2
<i>klebseilla oxycota</i>	5	1	6	3	3	6
<i>proteus mirabilis</i>	2	0	2	1	1	2
<i>pseudomonas aeruginosa</i>	1	0	1	0	1	1
<i>staph aureus</i>	7	3	10	5	5	10
<i>staph saprophyticus</i>	1	0	1	0	1	1
<i>strep pneumonia</i>	5	2	7	4	3	7
<b>Total</b>	35 (74.4%)	12 (25.5%)	47	21 (44.65)	26 (5.3%)	47

### 3.7 Susceptibility pattern of each nosocomial infection to evaluated antibiotics

Table 7 illustrates the susceptibility character of nosocomial infections to antibiotics. As shown, Wound pus displayed the most sensitivity to Doxycycline antibiotic than antibiotics.

**Table. 7** Susceptibility pattern of each nosocomial infection to evaluated antibiotics

SXT Susceptibility			Total	DO Susceptibility		Total
Sample	R	S		R	S	
Blood	4	3	7	1	6	7
Wound pus	25	4	29	14	15	29
Medical Device Tip	3	4	7	5	2	7
Burn	3	1	4	1	3	7
<b>Total</b>	35	12	47	21	26	47

### 3.8 Comparison of sensitivity patterns to Septrin, Doxycycline antibiotics and age groups of patients

Patients between 3 to 65 years old (n=30) had a somewhat high resistance to Septrin. In the age group 66 to 86 years (n=5), the resistance to seprin is somewhat reduced.

**Table. 8** Comparison of sensitivity patterns to Septrin, Doxycycline antibiotic and age groups of patients

Age	SXT Susceptibility		Total	DO Susceptibility		
	R	S		R	S	Total
3-23	10	1	11	6	5	11
24-44	10	4	14	4	10	14
45-65	10	5	15	9	6	15
66-86	5	2	7	2	5	7

SXT Susceptibility			Total	DO Susceptibility		
Age	R	S		R	S	Total
3-23	10	1	11	6	5	11
24-44	10	4	14	4	10	14
45-65	10	5	15	9	6	15
66-86	5	2	7	2	5	7
<b>Total</b>	35	12	47	21	26	47

### 3.9 Distribution of Isolates by Ages

In females, the most predominant wound and blood stream infection was *Staph aureus* among 45-65. While *Strep pneumonia* is considered as the second type isolated from the age group of 24-44.

**Table. 9** Distributions of Isolates by Ages

Isolates	Age				Total
	3-23	24-44	45-65	66-86	
<i>Acinetobacter baumannii</i>	1	0	1	0	2
<i>Citrobacter freundii</i>	0	0	1	0	1
<i>E. Coli</i>	1	3	0	1	5
<i>Enterococcus faecalis</i>	0	2	0	0	2
<i>Serratia marcescens</i>	1	0	0	0	1
<i>Klebseilla pneumoniae</i>	0	0	1	2	3
<i>Staph epidermidis</i>	0	3	0	0	3
<i>Step pyogen</i>	0	0	1	0	1
<i>Strep agalactia</i>	1	0	1	0	2
<i>klebseilla oxycota</i>	3	2	1	0	6
<i>proteus mirabilis</i>	1	0	1	0	2
<i>pseudomonas aeruginosa</i>	0	0	0	1	1
<i>staph aureus</i>	1	1	6	2	10
<i>staph saprophyticus</i>	0	0	1	0	1
<i>strep pneumonia</i>	2	3	1	1	7
<b>Total</b>	11	14	15	7	47

### 3.10 Prevalence of different bacterial growth among female and male patients.

A total of 47 specimen culture results of patients with suspected infection during twelve months period were studied. The isolates and sex distribution of the patients are shown in Table 9. Bacteria isolation rate was in females 25 (53.1%) than males 22 (46.8%) had a positive bacterial culture result. *Staph aureus* was the most frequent isolate from female patients.

**Table. 10** Prevalence of different bacterial growth among female and male patients.

Bacteria	Sex		Total
	Female	Male	
<i>Acinetobacter baumannii</i>	1	1	2
<i>Citrobacter freundii</i>	0	1	1
<i>E. coli</i>	3	2	5
<i>Enterococcus faecalis</i>	2	0	2
<i>Serratia marcescens</i>	1	0	1
<i>Klebsiella pneumoniae</i>	2	1	3
<i>Staph epidermidis</i>	1	2	3
<i>Strep pyogen</i>	1	0	1
<i>Strep agalactia</i>	1	1	2
<i>klebsiella oxycota</i>	4	2	6
<i>Proteus mirabilis</i>	0	2	2
<i>pseudomonas aeruginosa</i>	0	1	1
<i>staph aureus</i>	6	4	10
<i>staph saprophyticus</i>	0	1	1
<i>strep pneumonia</i>	3	4	7
<b>Total</b>	25	22	47

### 3.11 Distribution of Susceptibility Patterns of Isolates to Septrin, Doxycycline by Sex.

In this study, a total of 47 isolates from diverse swabs and blood samples were experimented with in vitro by the disk diffusion test to define the susceptibility of these bacteria to Septrin. Isolates that were resistant to Septrin were shown in females more than males. Isolates that were resistant to Doxycycline was also shown in females more than males.

**Table. 11** Distributions of Susceptibility Patterns of Isolates to Septrin, Doxycycline by Sex.

Gender	SXT			DO		Total
	R	S	Total	R	S	
Female	21	4	25	11	14	25
Male	14	8	22	10	12	22
<b>Total</b>	35	12	47	21	26	47

## 4. Discussion

In the present research, *Staph aureus* was the most frequently isolated pathogen in our study followed by *strep pneumonia* and *klebsiella oxycota*. This is agreeing with the statements of several studies that expressed *S. aureus* as the most familiar bacteria rescued from wound infection (Aung *et al.*, 2016; Kassam *et al.*, 2017).

Bloodstream infection is a demanding issue, and occasionally, it may be life-threatening; consequently, timely detection, identification, and antimicrobial susceptibility testing of blood-borne. Culture positivity was noticed in 7 (14.8%) of adult

patients. In this study, males (n=6) had high culture positivity with septicemia as compared with females (n=1).

The result was consistent with the investigations done by Hussein et al., 2005; Kaur and Singh, 2014) who registered high culture positivity in men. The maximum number of positive blood cultures was from age of 75, and the minimum was from the age of 27 years.

Gram-positive cocci were the most ordinary organism isolated. These results studies are analogous to a study done by Kokku *et al.*, 2020, but not consistent with Henry *et al.*, 2018, were noticing Enterobacteriaceae was the most commonly isolated gram-negative organisms, constituting 67% (122/184) of all gram-negative bacilli (Emerging Infectious Diseases, 2018).

Several studies have been accomplished to determine the sources of bacterial infections in hospitals. Clearly, bacterial colonies have been recognised at various hospital environment sites including beddings, stethoscopes (Lapeña and Jose, 2013), computers, catheters (Alawkally *et al.*, 2019) and other small electronic devices and instruments utilized by health workers.

There was 7 medical equipment that had bacterial isolates determined from the sample in this study. Of the bacteria found, the most frequent isolates were *E. Coli*, *Klebsiella pneumonia*, *staph aureus* and *strep pyogen* 2 (4.2%) equally. A case study by Ameh *et al.*, investigating the prevalence of bacterial isolates on hospital surfaces in Nigeria (Ameh *et al.*, 2009) found the following isolates with prevalence: *E. coli* (34.4%), *Klebsiella* species. (21.9%), *Pseudomonas* species. (15.6%), *Staphylococcus* species. (12.5%), *Proteus* species (9.4%).

Most of the world literature indicates that *staph aureus* is one of the main pathogens accountable for a number of infections in hospital settings, with considerable morbidity and mortality (Ho *et al.*, 2009).

*S. aureus* represents one of the most severe gram-positive bacterial infections in nosocomial and community settings. The severity of *S. aureus* infections is related to the different potential infected tissues (ranging from skin to bloodstream). *S. aureus* strains were recovered from various materials and analyzed with traditional microbiological tests to specify their antimicrobial susceptibility. Hospitalized patients show a high frequency of *S. aureus* infections due to their weak immune system and frequent injections and catheterizations. Moreover, in these types of patients, *S. aureus* can lead to life-threatening infections such as endocarditis and osteomyelitis. Our study indicates a high rate of resistance of isolates to Septrin than Doxycycline. Analogous to our results, in the study of Sadeghi *et al.*, 2021.

## 5. Conclusion

In conclusion, the problem of antibiotic resistance is very serious in Libya and appears to be on the rise. In this study, the prevalence of antibiotic resistance was high in most common pathogenic strains. Therefore, the results of this study demonstrated that both antibiotics with a high resistance level must be less used for the treatment of infections. Moreover, to prevent the

spread of resistance among various strains and improve the effectiveness of antibiotics, it is suggested to establish a precise schedule for antibiotic use in this region based on their antibiotic resistance pattern. The health authorities in particular and society in general should address the problem of antimicrobial resistance urgently. Establishing monitoring systems based on routine testing of antimicrobial sensitivity and education of healthcare workers, pharmacists, and the community on the health risks associated with the problem and benefits of prudent use of antimicrobials are some of the steps that can be taken to tackle the problem in the future.

## **6. Ethical Approval**

The study was approved by the Ethics Committee of Al Saleem Medical Laboratory.

## **7. Consent**

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

## **8. Limitation of the study**

The study did not determine the resistance detected was due to hospital-acquired or community acquired infection, also the study did not show the trends of antibiotics resistance from year to year.

## **9. Author Contributions**

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

## **10. Conflict of interest and funding**

The authors declare no conflict of interest and not funding or benefits from any source.

## **11. Funding**

Not applicable.

## **12. Acknowledgements**

The authors are appreciative of the support of colleagues in the Bacteriology Department at Al Saleem laboratory.

## **12. References**

Alawkally, Noor Alhooda & Nag, Miftah & Ali, Maree & Al-Awkally, Nesrine & Al-Awkally, Abeer & Al-Awkally, Alreda. (2022). Antibiotic sensitivity of *Escherichia coli* isolated from different clinical specimens of patient human in Al-jala hospital-Benghazi Libya. International Journal of Current Research in Science Engineering & Technology. 8. Pages: 509-517.

- Al-awkally, N. A. M., DokallyAli, M., Ali, F. M., & Abouserwel, A. (2019). Antibiotic Susceptibility and Resistant Pattern of Isolates of *Pseudomonas aeruginosa* recovered from Infected Swabs, Abscess, Burn, Medical Tips and Blood from Patients at 4 Geographical Locations in Libya (Al-Bayda, Shahat, Derna and Benghazi). *Int. J. Curr. Microbiol. App. Sci*, 8(10), 143-149.
- Ameh EA, Mshelbwala PM, Nasir AA, Lukong CS, Jabo BA, Anumah MA, Nmadu PT. Surgical site infection in children: prospective analysis of the burden and risk factors in a sub-Saharan African setting. *Surg Infect*. 2009; 10(2):105–9.
- Aung, M.S.; Zi, H.; Nwe, K.M.; Maw, W.W.; Aung, M.T.; Min, W.; Nyein, N.; Kawaguchiya, M.; Urushibara, N.; Sumi, A. Drug resistance and genetic characteristics of clinical isolates of staphylococci in Myanmar: High prevalence of PVL among methicillin-susceptible *Staphylococcus aureus* belonging to various sequence types. *New Microbes New Infect*. 2016, 10, 58–65. [CrossRef] [PubMed]
- CDC. Types of healthcare-associated infections. Healthcare associated infections (HAIs). 2016 [Online] Available from: <https://www.cdc.gov/HAI/infectionTypes.html> [Accessed on 10th August, 2016]
- Clinical and Laboratory Standards Institute (CLSI), Performance Standards for Antimicrobial Disk Susceptibility Tests, Clinical and Laboratory Standards Institute, Wayne, PA, USA, 13 edition, 2018.
- Clinical and Laboratory Standards Institute. Methods for antimicrobial dilution and disk susceptibility testing of in frequently isolated or fastidious bacteria M45-A2, vol. 30. 2nd ed. Wayne: Clinical and Laboratory Standards Institute; 2010
- Daneman, A. H. Rishu, A. H. Pinto et al., “7 versus 14 days of antibiotic treatment for critically ill patients with blood- stream infection: a pilot randomized clinical trial,” *Trials*, vol. 19, no. 1, p. 111, 2018
- Edwards R, Harding K. Bacteria and wound healing. *Curr Opin Infect Dis*. 2004;17:91–6. *Emerging Infectious Diseases* • [www.cdc.gov/eid](http://www.cdc.gov/eid) • Vol. 24, No. 1, Januar 2018
- Fakron, A., Alawkally, N., El-warred, S., Aldouakali Ali, M., El-amari, M., Awkally, A., Soutiyah, M., Suliman, N. and Ibrahim, H. 2022. Risk Factors for Ciprofloxacin and Gentamycin Resistance among Gram Positive and Gram Negative Bacteria Isolated from Community-Acquired Urinary Tract Infections in Benghazi city. *Scientific Journal for the Faculty of Science-Sirte University*. 2, 1 (Apr. 2022), 76-87. DOI:<https://doi.org/10.37375/sjfssu.v2i1.204>
- H. A. Khan, A. Ahmed, and R. Mehboob, “Nosocomial infections and their control strategies,” *Asian Pacific Journal of Tropical Biomedicine*, vol. 5, no. 7, pp. 505–509, 2015.
- HO K.M., ROBINSON J.O. (2009). Risk factors and outcomes of methicillin-resistant *Staphylococcus aureus* bacteraemia in critically ill patients: a case control study. *Anaesth. Intensive Care*. 37, 457-463.
- K. Inweregbu, J. Dave, and A. Pittard, “Nosocomial infections,” *Continuing Education in Anaesthesia, Critical Care & Pain*, vol. 5, no. 1, pp. 14–17, 2005.

- Kassam, N.A.; Damian, D.J.; Kajeguka, D.; Nyombi, B.; Kibiki, G.S. Spectrum and antibiogram of bacteria isolated from patients presenting with infected wounds in a Tertiary Hospital, northern Tanzania. *BMC Res. Notes* 2017, 10, 1–6. [CrossRef] [PubMed]
- Kaur, A. and Singh, V. (2014) Bacterial isolates and their antibiotic sensitivity pattern in clinically suspected cases of fever of unknown origin. *JK Science* 16: 105–109.
- Kokku PK, Priyadarshini T, Sravanthi CH. Bacteriological profile and antimicrobial sensitivity pattern of blood culture isolates. *J Evid Based Med Healthc* 2020; 7(38), 2092-2095. DOI: 10.18410/jebmh/2020/434
- Lapeña, Jose Florencio. "Mirrors and reflections: the evolution of indirect laryngoscopy." *Annals of Saudi Medicine* 33.2 (2013): 177-181
- M. J. Bizzarro, V. Shabanova, R. S. Baltimore, L.-M. Dembry, R. A. Ehrenkranz, and P. G. Gallagher, "Neonatal sepsis 2004- 2013: the rise and fall of coagulase-negative staphylococci,"(e *Journal of Pediatrics*, vol. 166, no. 5, pp. 1193–1199, 2015.
- Noor-alhoda Milood Al-awkally, Maree Dokally Ali, Reeda Miloud Al-awkally, Abeer Miloud AL-awkally, Fowziya M. Ali and Ahmed Abouserwel. 2019. Antibiotic Susceptibility and Resistant Pattern of Isolates of *Pseudomonas aeruginosa* recovered from Infected Swabs, Abscess, Burn, Medical Tips and Blood from Patients at 4 Geographical Locations in Libya (Al- Bayda, Shahat, Derna and Benghazi). *Int.J.Curr.Microbiol.App.Sci.* 8(10): 143-149.  
doi: <https://doi.org/10.20546/ijcmas.2019.810.015>
- Hussein, A., Sayed, A. and Mohamed, A. (2005) Seroepidemiological study on human brucellosis in Assiut Governorate. *Egypt J Immunol* 12: 49–56.
- Sadeghi et al. *BMC Res Notes* (2021) 14:88 <https://doi.org/10.1186/s13104-021-05503-0>
- Siddiqui A, Bernstein J. Chronic wound infection: Facts and controversies. *Clin Dermatol.* 2010; 28:516–26.
- Hess C. Checklist for factors affecting wound healing. *Adv Skin Wound Care.* 2011;24(4):192.
- Sweileh, Waleed M. "A prospective comparative study of gentamicin-and amikacin-induced nephrotoxicity in patients with normal baseline renal function." *Fundamental & clinical pharmacology* 23.4 (2009): 515-520.
- Tian, Y., Zheng, B., Wang, B., Lin, Y., and Li, M. (2016). Rapid identification and multiple susceptibility testing of pathogens from positive-culture sterile body fluids by a combined MALDI-TOF mass spectrometry and vitek susceptibility system. *Front. Microbiol.*7:523. doi: 10.3389/fmicb.2016.00523.
- WHO. The burden of health care-associated infection worldwide. 2016 [Online] Available from: [http://www.who.int/gpsc/country\\_work/burden\\_hcai/en/](http://www.who.int/gpsc/country_work/burden_hcai/en/) [Accessed on 10th August, 2016].