

Original Research Article

KNOWLEDGE OF MALARIA PREVENTIVE MEASURES IN A HYPERENDEMIC RURAL AREA OF WEST AFRICA

ABSTRACT

The Abuja declaration on Roll Back Malaria in Africa on 25th April 2000 by 44 African heads of state and government adopted the use of insecticide treated bed nets (ITNs) as preventive measure against the adverse effects of malaria among pregnant women in Africa. Similar optimism is expressed for the coverage of intermittent presumptive treatment (IPT) of Malaria infection among pregnant women across the continent. We conducted a community-based survey of recently pregnant women randomly selected from eight communities of Ikwo Local Government Area of Ebonyi State, Nigeria in June 2016, to study the level of knowledge of rural dwellers on the use of insecticide-treated bed nets by pregnant women. Among 300 households surveyed, 76.5% stated their high knowledge of and use of insecticide treated bed nets during pregnancy. Greater population of women of these communities (88.5%) knew the dangers of malaria in pregnancy and to the unborn child. The coverage of nationally recommended strategy to prevent malaria during pregnancy was high across all the communities studied. More sensitization and expansion of access to this service are required to meet the fullest meaning of the name "ROLL BACK malaria to where it came from". The sealing up of malaria prevention program through antenatal care (ANC) services should be encouraged with 78.2% of women still indulging in self-treatment of malaria during pregnancy. Issues of commodity supply and service-free costs to clients should be encouraged and sustained.

Keywords: Malaria, Measure, Hyperendemic, Pregnancy, Prevalence.

Introduction

Susceptibility to malaria infection during pregnancy and incidence of anemia in children are determined by the level of immunity prior to pregnancy, which depends on intensity and stability of malaria transmission (Ndyomugenyiet al, 2003). Populations in Africa live under a wide range of exposure to *Plasmodium falciparum* but how this affects the clinical and mortality of malaria remains unclear. As transmission falls from high to lower levels, immunity to severe disease is acquired more slowly and consequently the average age of severe malaria tend to increase. Increase in the age of susceptibility to severe malaria have been associated with increased

Comment [PDU1]: Declaration

Comment [PDU2]: insecticide-treated

Comment [PDU3]: a preventive

Comment [PDU4]: insecticide-treated

Comment [PDU5]: A greater

Comment [PDU6]: a nationally

Comment [PDU7]: in

Comment [PDU8]: is

Comment [PDU9]: the intensity

Comment [PDU10]: affects

Comment [PDU11]: tends

Comment [PDU12]: An increase

Comment [PDU13]: has

rate, increased risk of cerebral malaria and increased fatality due to severe malaria. These observations have led to concerns that reductions from high to lower levels of transmission intensity of *P.falciparum* may not necessarily reduce the burden of the disease in the long term and could even result in paradoxical increase. A definitive answer to the question of how the burden of malaria will change following reductions in transmission may only be possible by maintaining a controlled trial of an effective intervention such as insecticide-treated nets for several years (Hugh *et al*, 2003). The value of insecticide-treated nets in preventing malaria is now established beyond doubt but their level of use is still low in malaria endemic countries. Also there is evidence from several sources that the community-wide mass effect of widespread use of treated nets is as important, than the personal protection effect to the individual user (Curtis *et al*, 2003). This is relevant to the current debate about the relative efficiency of free provision versus marketing of treated nets. In Africa, malaria control is reliant upon antimalarial chemotherapy and insecticide based vector control and in each case successful intervention is threatened by the emergence and spread of resistance to the compounds used. Insecticide-treated nets have been shown to be effective in increasing child survival through preventing malaria in various settings, at least over one or two year period (Schellenberg, 2003). There is an optimism expressed by international roll back malaria (RBM) for the coverage of insecticide-treated nets (ITNs) among pregnant women across the continent by at least 60% by 2005 (<http://www.rbm.who.int>). However, poor knowledge of rural-dwellers on malaria vector control and the absence of public partnerships on sustainable availability of insecticide-treated nets pose a challenge to Roll Back Malaria initiative. Therefore, there is a need to know what communities think about different approaches to malaria control measures. New preventive health interventions such as ITNs can only be considered successful if they are also socially and culturally acceptable. Planning and monitoring of effective vector-orientated approaches to malaria control requires information about the level of knowledge of the disease across the populations of an area. This work, therefore, studied the need to characterize knowledge of malaria disease control measure over a small area in order to define the important and complex relationship between persistence and prevalence of malaria.

Comment [PDU14]: a paradoxical

Comment [PDU15]: malaria-endemic

Comment [PDU16]: insecticide-based

Comment [PDU17]: Insecticide-treated nets effectively increase child survival by preventing malaria in various settings

Comment [PDU18]: years

Comment [PDU19]: citation writing must be consistent

Comment [PDU20]: require

Comment [PDU21]: measures

Comment [PDU22]: to

Methodology

The study was conducted between April and June 2017, the peak of malaria transmission and high infection rate in the country. It was carried out in eight communities across Ikwo local district of Ebonyi State, Nigeria as part of research work funded by National TETFund under the auspice of Ebonyi State University. Maps of enumeration areas (EA), created during 2005 National Population Census of the country and representing approximately 100 people or 10 homesteads. Random sampling of EAs was carried out in each mapped community up to 250 homesteads as representatives of the community.

Comment [PDU23]: A random

Comment [PDU24]: community representatives.

Survey procedures

We designed a structured interview questionnaire adapted from our earlier study performed in another state of the country Anambra state, Nigeria, Onwujekwe *et al*, (2007). The interview team members were indigenes of the communities who had high education training. They were fluent in both English and the local dialect of the communities. We selected 4 women and 4 men as interviewers, and they were given a week intensive training on translation and administration of questionnaire to household heads or the representatives. The questionnaire consisted of information on household demographics, education background of the caregiver or head of the household and the knowledge of the caregiver or the representative on insecticide-treated nets and other malaria preventive measures. We also trained 4 Community Health Extension Workers, who monitored the interviewers and served as quality control measures to the administration of the questionnaires. Our Cartographer used streams, roads and other natural landmarks to subdivide the EAs to make it easier for the team of interviewers to count the households into 10-50 households each.

Comment [PDU25]: week's

Comment [PDU26]: the educational

Comment [PDU27]: administer

We defined a household as all persons who share earned income and farm produce and eat from the same cooking pot. Most often household members shared the same house, although at times teenage boys would have their own house while remaining a member of the economic family unit.

We interviewed all women who had completed a pregnancy within the last 2 years or who are still giving birth, in each household where there were children within 5 years of age. The team members revisited each selected household the same day or the following day if one or the representative of the person from who information was needed was not at home at time of the

Comment [PDU28]: the information

Comment [PDU29]: the time

first visit. Households where no one was at home and households where the occupants refused to participate were not replaced.

Results

Our teams visited 350 households. Five percent (5%) of the study participants (women) were absent but the primary heads of the households answered on their behalf. Eighty-nine percent (88.5%) knew the dangers of malaria in pregnancy and to the unborn child. They agreed that malaria prevention strategies during pregnancy are laudable policies and should be sustained. (Table I): Seventy-eight percent (78.2%) depended on self-medication for the treatment of malaria during pregnancy while eighty-nine percent (88.7%) knew the repercussions of self-medication. (Table II): Seventy-seven percent (76.5%) stated that they used insecticide-treated nets during pregnancy. (Table III): Most of the nets were obtained from government health facilities. Eighty-two percent (82.3%) had knowledge of malaria prevention strategies and that the policies are government recommendations. (Table IV): The use of insecticide treated nets was similar in all the communities.

Comment [PDU30]: to treat

Comment [PDU31]: knew

Comment [PDU32]: insecticide-treated

Table I: Opinions on malaria prevention and treatment

	Frequency (%)
Malaria prevention options are adequate	
Yes	284 (88.5)
No	37 (11.5)
People over self-prescribe/self-medicate	
Yes	251 (78.2)
No	70 (21.8)
Repercussion to the practice of self-medicating	
Yes	282(88.7)
No	36(11.3)

Table II: Malaria prevention services to pregnant women

	Frequency (%)
Drugs	
Yes	162 (85.7)
No	27 (14.3)

Insecticide treated nets	
Yes	179 (76.5)
No	55 (23.5)
Other	
Yes	12 (15.0)
No	68 (85.0)
Are the services based on govt. recommendations	
Yes	242(82.3)
No	52 (17.7)

Table III: Information of Government Policies on ITN

	Frequency (%)
Knowledge of government policies on ITN use	
Yes	246 (82.3)
No	53 (17.7)
Does policy provide guidelines for prevention of malaria in pregnancy	
Yes	96 (35.7)
No	173 (4.3)

Table IV: Information about ITN and first source of information

	Frequency (%)
Heard about ITN	
Yes	242 (90.0)
No	27 (10.0)
First Source of information	
Internet	61 (20.3)
Educational journal	2 (0.7)
Friends	50 (16.6)
Co-workers	31 (10.3)
Radio	63 (20.9)
Television	81 (26.9)

Others	13 (4.3)
--------	----------

Discussion and conclusion

This study, across eight communities of similar malaria endemicity in Ebonyi State, Nigeria, indicates that the knowledge and use of insecticide-treated nets by pregnant women met the recommended malaria preventive measures (knowledge and use of ITNs, 82% and 77% >60% of the targets set by RBM for 2005). Greater number of the study population slept under an insecticide-treated bed nets. Moreso, 82% of the women studied stated their knowledge of insecticide-treated bed nets as a malaria preventive measure during pregnancy. The source of the nets to the communities is mostly from government facilities. Surprisingly, none of the women stated that the net was obtained from commercial sector. This underscores the importance of government involvement in distribution of ITNs especially in rural settings. This raises concerns regarding the recent RBM strategic document for expanding bed net coverage that recommends the commercial sector as the main distributor of nets for the general population (WHO, 2002). The data from this study suggest that this may act only to exaggerate the disparities between those accessing health care in government health facilities and those patronizing traditional birth attendants with the later increasing in number. It will also disrupt their orientation that ITNs is a government policy and could only be accessed in government own facilities. However, the WHO document does allude to the use of ANC clinics for targeting pregnant women, it is less specific in modalities and position of this delivery system than it is for other social marketing or private sector approaches. It is also contrary to the objective of making all pregnant women access their health needs in government health facility as a means of reducing maternal-child morbidity and mortality rate. This study corroborates the relevance of relative efficiency of free provision versus marketing of treated nets. The level of knowledge and use of ITNs (82% and 77%) shows the irrelevance of social marketing which is an appealing tool for getting insecticide-treated nets to poor rural-dwellers of African communities and often involves targeted or general price subsidies. A study in Malawi showed that bed net ownership following a social marketing campaign was higher in urban than rural areas (Guyatt *et al*, 2004), and major reason given by most households throughout Africa for not owning a net is financial (Guyatt *et al*, 2002b, Holtz *et al*, 2002). The UNICEF free net distribution is a good motivating factor to achieving RBM

Comment [PDU33]: A greater

Comment [PDU34]: the commercial

Comment [PDU35]: only

Comment [PDU36]: increase

Comment [PDU37]: are

Comment [PDU38]: their government

Comment [PDU39]: However, the WHO document alludes to the use of ANC clinics to target pregnant women; it is less specific in modalities and position of this delivery system than it is for other social marketing or private sector approaches.

Comment [PDU40]: a government

Comment [PDU41]: the relative

Comment [PDU42]: net bed

Comment [PDU43]: the major

initiative. Although, in Nigeria, there is a USAID funded project (Net Mark) whose main purpose of study is to monitor changes in the markets, such as variations in sales volumes; prices at both retail and wholesale levels, in both urban and rural areas; supply systems brand diversity and competitive structure. The high knowledge of dangers of malaria in pregnancy with the unborn child along with high use of insecticide-treated nets during pregnancy highlights the need for improved supply, distribution and sustainability.

Comment [PDU44]: The high knowledge of the dangers of malaria in pregnancy with the unborn child and the high use of insecticide-treated nets during pregnancy highlights the need for improved supply, distribution, and sustainability.

REFERENCES

- Curtis, C., Maxwell, C., Lamnge, M. *etal* (2003). Sealing up coverage with insecticide-treated nets against malaria in Africa: who should pay? *Lancet Infectious Diseases*, 3, 304 – 307.
- Gruyalt, H.L., Ochola, S.A. & Snow, R.W. (2002b). Too poor to pay: charging for insecticide-treated bednets in highland Kenya. *Tropical Medicine and International Health* 10, 846-850.
- Gruyalt, H.L. & Ochola, S.A. (2003). The use of bednets given free to pregnant women attending antenatal care in Kenya. *The Lancet* 362, 1549-1550.
- Holtz, T.H., Marum, L.H., Mkandala, C. *etal* (2002). Insecticide-treated bednet use, anaemia and malaria parasitaemia in Blantyre District, Malawi. *Tropical Medicine and International Health*, 7, 220 -230.
- Hugh, R., Chris, D., Ilona, C. *etal* (2003). The epidemiology of severe malaria due to *Plasmodium falciparum* at different transmission intensities in NE Tanzania. *LSHTM Malaria Centre Report 2002 -2003*.
- Joanna Schellenberg (2003). Long-term effect of treated mosquito nets on child's survival in an area of high *Plasmodium falciparum* transmission: a cohort study. *LSHTM Malaria Centre Report 2002 – 2003*.
- Clarke, S., Ndyomugenyi, R., Neema, S., Magnussen, P. (2003). Malaria in the highlands of Mgada: Strategy for control. *LSHTM Malaria Centre Report 2002 -2003*
- Onwujekwe, O., Uzochukwu, B., Ezumah, N. & Shu, E. (2005). Increasing the coverage of insecticide-treated nets in rural Nigeria: Implications of consumer knowledge, preferences and expenditures for malaria prevention. *Malaria Journal*, 18, 4 (1);29.

WHO(2002). Sealing-up Insecticide-treated Netting Programmes in
Africa.WHO/CDS/RBM/2002.43.

UNDER PEER REVIEW