

Evaluation of Anti-diabetic and Antioxidant Properties of Ethanol Leaf Extract of *Jathropa Tanjorensis*

Abstract

The aim of this study was to evaluate the anti-diabetic and antioxidant properties of ethanol leaf extract of *Jathropa tanjorensis*. Freshly harvested leaves of *Jathropa tanjorensis* were thoroughly processed into extract. Twenty five (25) adult male wistar rats were divided into five groups of five rats each. **Group I** was the normal control allowed unrestricted access to food and water only. **Group II** was the negative control and was induced with diabetes without treatment, **Groups III and IV** were diabetic rats administered with ethanol leaf extract of *Jathropa tanjorensis*, while **Group V** was diabetic administered with the standard drug. After treatment had been concluded, animals were sacrificed and blood and tissue obtained were analyzed using standard procedures. A significantly ($P < 0.05$) high blood sugar level was reported for diabetic rats which however was significantly ($P < 0.05$) reversed with oral administration of ethanol leaf extract of *Jathropa tanjorensis* in a dose dependent manner. The activity of the liver antioxidant enzymes was significantly ($P < 0.05$) reduced in diabetic rats which however was significantly ($P < 0.05$) increased following oral administration of *Jathropa tanjorensis* leaf extract in a dose dependent manner across treatment periods. In conclusion, the study unveils the antidiabetic potential of *J. tanjorensis* leaf.

Keywords: *Jathropa tanjorensis*, antioxidant enzymes, Anti-diabetic, Diabetes, Liver

Introduction

Diabetes mellitus (DM) is a disease condition which is characterized by an abnormally level of sugar in the blood. It abounds globally and has been identified as the one of most lethal diseases of mankind. An estimated 4.4% of the world population is projected to suffer from diabetes in 2030 [1].

Diabetic neuropathy, nephropathy and retinopathy among others have been implicated in prolonged diabetes mellitus [2] while complications arising from the condition are hypertention, atherosclerosis and micro circulatory disorders [3].

In diabetic patients, mononuclear cells are effectively engaged in the generation of reactive oxygen species (ROS) and consequently, macromolecules such as lipid, protein and DNA are damaged [4].

J. tanjorensis is a member of the *Euphorbiaceae* family. It is cultivated widely in the Southern part of Nigeria. Interest among researchers in the aforementioned plant has grown over the years owing to its immense health significance, availability and affordability [5]. Analysis on the leaf of *J. tanjorensis* unveiled the presence of alkaloids, flavonoids, tannins, cardiac glycoside, anthroquinones and saponins [6]. Reports abound on the antihypertensive, antimicrobial, antimalarial and hypolipidemic activity of *Jathropha tanjorensis* [7] hence, the need to probe the leaf further in an effort to unveil more health benefits of the said plant.

Materials and Methods

Collection of plant material

Fresh leaf of *J. tanjorensis* obtained a local market was identified at the herbarium unit of the Department of Forestry, Michael Okpara University of Agriculture, Umudike, Abia State.

Sample Preparation

Leaves of *J. tanjorensis* were air dried for three days after which they were ground to fine powder with the aid of an electric blender. The powdered sample was stored in a moisture free, air-tight container until further use. 500 g of powdered plant sample was soaked in 96%

ethanol for 2 hr. The extract obtained was filtered and concentrated with a rotary evaporator. The brownish residue obtained was dried in desiccator

Animals

Adult male wistar rats weighing between 150-250 g were obtained from the animal house of Abia State University Uturu. The rats were housed and maintained in well ventilated plastic cages under standard laboratory conditions and were allowed unrestricted access to food and water. Acclimatization of the animals occurred within three weeks before the experiment.

Median Lethal Dose 50% (LD50%)

The LD 50% was determined using three groups of three wistar rats and was each subsequently administered with 10, 100 and 1000 mg/kg of extract orally. Animals were studied for 24 h to observe signs of toxicity. Upon confirmation of absence of mortality in any of the groups, another three groups of one rat each was each administered with 1600, 2900 and 5000 mg/kg of extract separately and animals were observed for 48 h for signs of toxicity according to Lorke [8].

Animal Grouping

A total of 25 rats divided into 5 groups of 5 rats each were used in this study.

Group 1: Non diabetic without treatment. (Normal control)

Group 2: Diabetic rats without treatment (Negative control)

Group 3: Diabetic rats treated with 200 mg/kg .bw *J. tanjorensis* leaf extract.

Group 4: Diabetic rats treated with 400 mg/kg .bw *J. tanjorensis* leaf extract.

Group 5: Diabetic rats treated with 2.5 mg/kg Metformin daily.

Induction of Diabetes

Induction of diabetes mellitus was by a single intraperitoneal injection of 120 mg/kg body weight of alloxan [9] and blood sugar level evaluated with the aid of a glucometer (Acc-cheek Advantage Roche diagnostics GmbH, Germany) after three days, and the rats with fasting blood glucose level in excess of 126 mg/dl (11.1mmol/L) were deemed eligible for the study.

Preparation of Liver Homogenate

In 1:5 of 0.9% sodium chloride (ice cold), the liver tissue was homogenized and subsequently centrifuged at 3500 rpm for 20 minutes to obtain the supernatant which was used to assay for the activity of superoxide dismutase (SOD) and catalase (CAT).

Biochemical analysis

Determination of Superoxide Dismutase Activity

The method described by Martin et al. [10] was employed to determine the superoxide dismutase activity. Exactly 920 μL of phosphate buffer (0.05 M, pH 7.8) was added to 40 μL of the sample. A reagent test was prepared by replacing the sample with 40 μL of sample dilution buffer (0.85% NaCl). The mixtures were incubated for 2 min at 25⁰C before 40 μL of hematoxylin was added. Following the addition of 40 μL of hematoxylin, absorbance of the sample test and reagent test was read at 560 nm immediately and after 5 minutes against the sample blank which is distilled water.

SOD concentration in the sample was calculated thus;

Absorbance Reagent test (AR) = Absorbance Reagent test 2 – Absorbance Reagent test 1

Absorbance sample test (As) = Absorbance sample test 2 – Absorbance sample test 1

% inhibition = $[1 - As / AR] \times 100$

SOD (μ /ml) = $[1 - As / AR] \times 100 \times 1.2$

Determination of Superoxide Dismutase Activity

To 1 ml of reaction mixture containing 1.96 mL phosphate buffer (0.01 M, pH 7.0), 1.0 mL hydrogen peroxide (0.2 M) and 0.04 mL of homogenate in a final volume of 3.0 mL. 2 ml of dichromate acetic acid reagent was introduced and subsequently heated for 10 minutes before being cooled and absorbance read at 570 nm [10].

Statistical Analysis

Data were analysed using one way Analysis of Variance (ANOVA). Differences in mean were compared using Turkey Test. *p-values* less than 0.05 was considered statistically significant.

Table 1: Blood Glucose Levels of Diabetic rats treated with Ethanol Leaf extract of *J. tanjorensis*

Groups	Treatment	Blood Glucose Levels (mg/dl)		
		Week 1	Week 7	Week 14
Group I	Normal control	98.12±2.05 ^a	97.89±3.70 ^a	98.02±3.05 ^a
Group II	Diabetic-treatment	230.20±3.70 ^a	229.10±3.20 ^a	231.05±0.05 ^a
Group III	Diabetic+200 mg/kg extract	184.20±3.60 ^c	136.21±3.02 ^b	120.81±3.41 ^a
Group IV	Diabetic+400 mg/kg extract	110.00±4.61 ^c	106.21±2.02 ^b	100.00±3.02 ^a
Group IV	Diabetic+Std mg/kg extract	102.10±2.22 ^c	100.31±2.13 ^b	98.87±4.11 ^a

Results are expressed as mean ± Standard deviation of three determinations. Values with different superscript in a row are significantly ($P < 0.05$) different.

Table 2: Activity of Liver Antioxidant Enzymes in Diabetic rats treated with Ethanol Leaf extract of *J. tanjorensis*

GROUPS	TREATMENT	Enzyme activity	
		SOD (U/ml)	CAT(U/ml)
Group I	Normal control	2.60±2.02 ^c	49.33±2.40 ^d
Group II	Diabetic-treatment	0.90±3.20 ^a	32.00±0.58 ^a
Group III	Diabetic+200 mg/kg extract	1.60±1.20 ^b	41.00±1.15 ^b
Group IV	Diabetic+400 mg/kg extract	1.76±1.23 ^{bc}	41.33±2.60 ^b
Group IV	Diabetic+Std mg/kg extract	2.00±1.30 ^c	45.66±2.73 ^c

Results are expressed as mean \pm Standard deviation of three determinations. Values with different superscript in a column are significantly ($P < 0.05$) different.

RESULTS AND DISCUSSION

Diabetes mellitus is a multifactorial disease characterized by increased blood sugar level beyond the normal range [11]. It is characterized by a deficiency in antioxidant enzymes [12]. Table 1 shows the blood glucose levels of diabetic rats treated with ethanol leaf extract of *J. tanjorensis*. An increased blood sugar level which progressed with time was reported in diabetic rats. However, oral administration of ethanol leaf extract of *J. tanjorensis* significantly ($P < 0.05$) caused a reduction in the blood sugar level of diabetic rats in a dose dependent manner. The antidiabetic potential of the aforementioned plant could be attributed to the presence of polyphenolic compounds which had been reportedly present in leaf of the aforementioned plant [13]. This result is consistent with the finding of Asuk et al. [14] which established that the ethanol-methanol extracts of the leaf, stem, bark and root of *Jatropha curcas* demonstrated appreciable anti-diabetic property. There is a correlation between diabetes and oxidative tissue damage arising from free radical generation [15]. Diabetes causes increased free radical generation which results in the alterations of the liver tissue superoxide dismutase (SOD) and catalase [15]. Table 2 shows the activity of liver antioxidant enzymes in diabetic rats treated with leaf extract of *J. tanjorensis* showing a significantly ($P < 0.05$) decreased activity of liver antioxidant enzymes in diabetic rats which however was significantly ($P < 0.05$) increased following oral administration of *J. tanjorensis* leaf extract in a dose dependent manner across treatment periods. This could be attributed to the antioxidant property of the plant extract which creates a balance between free radical generated and eliminated. This result is in tandem with the

finding of Asuk et al [14] which reported the antioxidant activity of the leaf, root and stem of *Jatropha curcas* a member of the *Euphorbiaceae* family to which *J. tanjorensis* belongs.

Conclusion

It is evident that the leaf of *Jatropha tanjorensis* yields impressive antidiabetic and antioxidant properties and thus should be analysed further to unveil the active components characteristic of the said function for drug development purposes.

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

REFERENCES

- [1] Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care*. 2004; 27(05):1047-1053.
- [2] Long L, Wang J, Lu X, Xiaofang L, Yuxia X, Shuhui Z, Canqiao L, and Yubin L. Protective effects of scutellarin on type II diabetes mellitus-induced testicular damages related to reactive oxygen species/Bcl-2/Bax and reactive oxygen species/microcirculation/ staving pathway in diabetic rat. *J Diabetes Res*. 2015; 2015:252530.
- [3] Saidu A N, Mann A, Onuegbu C D. Phytochemical screening and hypoglycemic effect of aqueous Blighiasapida root bark extract on normoglycemic albino rats. *Br. J. Pharm. Res.*2012; 156, 357-361.
- [4] Giugliano D, Marfella R, Coppola L, Verrazzo G, Acampora R, Giunta R, Nappo F, Lucarelli C. and D'Onofrio F. Vascular effects of acute hyperglycemia in humans are reversed by LArgenine. *Circ*. 1997; 95,1783-1790.
- [5] Omobuwajo OR, Alade GO, Akanmu MA, Obutor EM, Osasan SA. Microscopic and toxicity studies on the leaves of *J. tanjorensis*, *Afr J Pharm Pharmacol*. 2011; 5(1):12-17.
- [6] Omoregie ES, Osagie AU. Phytochemical screening and antianaemic effect of *Jatropha tanjorensis* leaf in protein malnourished rats. *Plants Arch*. 2007; 7:509-516.
- [7] Omoregie ES, Sisodia BS. Invitro antiplasmodial activity and cytotoxicity of leaf extracts from *J. tanjorensis*. *Pharmacology Online*. 2011; 2:656-673.
- [8] Lorke D. A new approach to practical acute toxicity testing. *Arch Toxicol*.1983; 54(4):275-287.
- [9] Burcelin R, Eddouks M, Maury I, Kande J, Assan R. and Girard J. Excessive glucose production, rather than Insulin resistance, account for hyperglycemia in recent onset streptozotocin-diabetic rats.1995; *Diabetologia*. 35:283-290.
- [10] Sinha AK. Colorimetric assay of catalase. *Anal Biochem*.1972; 47:389-394.
- [11]Ugochukwu NH, Babady NE, Cobourne M, and Gasset SR. The effect of *Gangronematifolium* extracts on serum lipid profile and oxidative stress in hepatocytes of diabetic rats. *J. Biosci*. 2003; 28(1), 1-5.
- Unwin N, Sobngwi E and Alberti KGMM. Type2 diabetes: the challenge of preventing a global epidemic, *Diabetes Int*.2001; 4-8.
- [13] Boussageon R, Bejan-Angoulvant T, Saadatian-Elahi M, Lafont S, Bergeonneau C, Kassai B, et al. Effect of intensive glucose-lowering treatment on all cause mortality,

cardiovascular death, and microvascular events in type 2 diabetes mellitus: a meta-analysis of randomized controlled trials. *Brit Med J.* 2011; 343:d4169.

[14] Asuk AA, Dasofunjo K, Okafor A I and Mbina FA. Antidiabetic and antioxidative effects of *Jatropha curcas* extracts in streptozotocin-induced diabetic rats. *BJMMR.* 2015; 5(3): 341-349.

[15] Matough FA, Budin SB, Hamid ZA, Alwahaibi N, Mohamed J. The role of oxidative stress and antioxidants in diabetic complications. *Squ Med J.* 2012; 12(1):5-18.

UNDER PEER REVIEW