

## Original Research Article

ANTI-DIABETIC EVALUATION OF METHANOL EXTRACT OF FICUS EXASPERATA VAHL. (MORACEAE)

### **ABSTRACT**

#### Background

Diabetes mellitus is a metabolic disorder characterized by chronic hyperglycaemia with disturbances of carbohydrate, fat and protein metabolism resulting from impairments in insulin secretion, insulin action or both. The side effect of orthodox is alarming and expensive. This prompted a search for a cheaper and reliable alternative drug medicinal plant for the management of diabetes.

#### Aim

This study aimed at carrying out anti-diabetics evaluation of methanol extract of *Ficus exasperata* leaves.

#### Methology

The cold maceration method was used to obtain the extract using methanol as solvent. Lorkes method was used to carry out the acute toxicity study. Alloxan induced model was used to evaluate the antidiabetic potentials and blood glucose level was measured using glucometer and statistically analyzed.

#### Result

Acute toxicity study indicated no mortality or any behavioral changes even at 5000mg/kg Body weight. The extract at 200mg/kg and 400mg/kg exhibited anti-diabetics activity in dose dependent manner. 400mg/kg body weight reduces blood glucose level significantly at p value less than 0.05 when compared to the standard drug.

#### Conclusion

The methanol extract of *Ficus exasperata* leaves have demonstrated a reasonable antidiabetics activity. The research has given an insight in the use of the plant leaves in controlling blood sugar level and substantiate its folkloric uses. Further studies on isolation of bioactive compound, establishment of the mechanism for the observed anti-diabetic activity and hepato-renal and hematological effects.

Keywords: Diabetes, Phytochemicals, Medicinal Plants, and blood glucose level

### **Introduction**

Diabetes mellitus is a disease caused by chronic hyperglycaemia due to defects in insulin secretion, insulin action or both [1]. Diabetes disorder is characterized by frequent urine, presence of glucose in urine and hyperglycemia [2]. There are two major clinical manifestation of diabetes, the type 1 diabetes or insulin dependent diabetes mellitus (IDDM) and type 2 diabetes or non-insulin dependent diabetes mellitus (NIDDM). Based on pathophysiology, type 2 diabetes mellitus is the predominant form of the disease mostly as a

result of consumption of diets that promote obesity coupled with physical inactivity. Globally, type 2 diabetes mellitus is considered one of the most rapidly growing non-communicable diseases with high mortality rate [3].

According to the International Diabetes Federation (IDF) 2021 updates, approximately 536 million adults between age 20-79 are living with diabetes while undiagnosed diabetes cases are estimated to be 183 million. The prevalence is projected to rise to 642 million adults by 2030 and 783 million by 2045. In Nigeria, there are estimated 3.6 million diabetic cases, with another 1.9 million undiagnosed cases between ages 20-79 years. It is projected to increase to 4.9 million by 2030 and 7.9 million by 2045 [4].

Currently, diabetes therapy is based on the use of hypoglycemic drugs (sulfonamides, biguanides, and insulin), on Diet measures, exercise, and requires a lifelong treatment. Considering the level of poverty in underdeveloped and developing countries, the need for a better and cheaper medication cannot be over emphasized. Medicinal plants have always provided a cheaper and time trusted alternative for the treatment and management of various diseases over time.

Medicinal plants have continued to attract attention in global search for effective methods of using plants parts: seeds, leaves, stem and root barks for many diseases affecting humans [5]. They are playing a major role in the treatment of various diseases and ailment in humans and animals since days immemorial with proven efficacy [6]. However, their acceptability by some cultures of the world has been limited by lack of specific dose regimen and thorough scientific information on toxicity profile to validate their safety [7].

*F. exasperate* Vahl. Belong to the family of moraceae, a terrestrial tropical shrub or small tree with scabrous, with ovate leaves that grows up to about 20m tall and prefers evergreen and secondary forest habitats. It is widespread in tropical Africa [8]. Traditionally different parts of this plant (fruit, leaf, sap, bark, and root) are considered medicinally important. They are

used as analgesic, diuretic, vermifuges, febrifuge, abortifacient, wound healing, leprosy, ophthalmic and oral infections, nasopharyngeal afflictions, arthritis, rheumatism, gout, edema, kidney disorders, diarrhea, dysentery, hemorrhoids, venereal diseases and animal fodder [9]. Previous researchers have reported that *F. exasperata* leaves possesses anti-inflammatory, antioxidant, antiulcer, anticonvulsant, antiarthritic, antimicrobial and hypotensive properties [11-14].

However, this study was carried out to evaluate methanol extract of *F. exasperata* for in vivo anti-diabetic potentials on alloxan induced diabetic rats

## **Material and methods**

### **Plant Sample collection**

The leaves of *F. exasperata* was collected from Agbani in Nkanu West Local Government Area of Enugu State, Nigeria on November, 2021. It was identified by a Taxonomist, Mr Ozioko Alfred of the International Centre for Ethnomedicine and Drug Development (InterCEDD), 110 Aku road Nsukka Enugu state and herbarium sample was deposited in the research Centre.

### **Animals**

Twenty-eight (28) apparently healthy male and female rats of body weights ranging between 57-144g and 12 Wistar albino mice, weighing 17-27g were used for the experiment. The animals were sourced from animal house of the Department of Pharmacology, Enugu State University of Science and Technology, Agbani Enugu under standard conditions. They were housed in aluminium cages in soft wood shavings as beddings in a room, at room temperature with free access to drinking water and rat chow food ad libitum. The rats were acclimatized to normal laboratory conditions prior to study.

### **Preparation of plant sample**

The fresh plant sample was collected from the bush, and the good leaves were separated from the unwanted parts of the plants. The fresh leaves samples were then washed under a running tap water to remove dust particles and other contaminants. The leaves were air dried under a shade at room temperature for 14 days. Thereafter the leaves were pulverized to a fine course powder using an electrical blender.

### **Extraction**

A 400g of the air dried, finely pulverized leaves sample was soaked in 2.5litres of methanol and left for 72 hours at room temperature of  $25^{\circ} \pm 2^{\circ}\text{C}$  with constant agitation. The extract was filtered first, using a fine grade cloth and second filtration was done using funnel clogged with cotton wool. The filtrate was concentrated using rotary evaporator under reduced temperature to obtain methanol extract. The percentage yield of the extract was determined and then transferred into an air-tight container and stored at  $4^{\circ} \pm 2^{\circ}\text{C}$  in a refrigerator until when needed.

### **Acute Toxicity Study**

This was performed according to method described previously [18]. 12 mice were used after acclimatization. An initial investigation involving administering (10, 100 and 1000 mg/kg) of the plant extract to three different groups of three mice each. After 24 hours and there was no death recorded. In this second stage, three dose levels were used (1600, 2900 & 5000 mg/kg) and physical observations were made up to 24 hours for mortality and behavioral changes

### **Alloxan-induced diabetes model**

The baseline blood glucose levels of the rats and weight were determined before they were induced with diabetes by intra peritoneal (IP) injection of 150mg/kg body weight of alloxan monohydrate solution. After 48 hours, the glucose levels of the rats were checked, the rats had increased glucose level using a glucometer. The diabetic rats were regrouped into 5

animals per cage in 4 different cages. The treatments were given to the rats in each group with a known anti-diabetic drug (positive control), the plant extract (200mg and 400mg) and normal saline (negative control). The blood glucose level of the diabetic rats was checked at 0 days (baseline), 4 days, 8 days and 12 days during the treatment period. The treatment lasted for 12 days after which the rats were sacrificed. The animals were grouped into 4 as follow: Group I- Negative Control rats received normal saline, Group II - Positive control treated with 5mg/kg body weight glibenclamide, Group III- Treatment with 200mg/ kg. body weight methanol extract, and Group IV- Treatment with 400mg/ kg. body weight methanol extract

### **Statistical analysis**

The data obtained was analyzed using SPSS version 23 and the significance difference between the control and treated groups determined using one-way analysis of variance (ANOVA) followed by turkey t-test. P values less than 0.05 were considered to be statistically significant.

## **RESULTS**

### **Percentage Yield of the Extract**

The extract gave a percentage yield of 4.5%.

Weight of crude sample = 400g

Total yield of the extract = 18.26g

Percentage yield = total yield / weight of the crude sample  $\times$  100

= 18.26g / 400g  $\times$  100

= 4.5%

### Acute Toxicity Study

The acute toxicity study result obtained showed no mortality in both phase 1&2 experiment as presented in Table 1 and 2.

**Table 1: phase 1 acute toxicity test**

Samples	Dose	Dose	Dose
	10mg/kg	100mg/kg	1000mg/kg
Methanol extract	0/3	0/3	0/3

**Table 2: phase 2 acute toxicity test**

Dose level	1600 mg/kg	2900 mg/kg	5000 mg/kg
Surviving rat	3/3	3/3	3/3

### Effect of Extract on the Blood Glucose Level of Alloxan Induced Diabetic Rats

The extract at 200mg/kg and 400mg/kg exhibited antidiabetics activity in dose dependent manner. 400mg/kg body weight reduces blood glucose lowering effect significantly at  $p < 0.05$  when compare to standard drug at 5mg/kg body weight as presented in table 3.

**Table 3.** Effects of *F. exasperata* leaves extract on blood glucose level of alloxan induced diabetic rats

Groups	Mean Anti-diabetic response per Day					
	IGLBE	Weight	0 day (GLAI)	Day 4	Day 8	Day 12
Negative control N. Saline	108.6±0.5	111.8±0.24	345.6± 0.37	215.2± 0.19	172.8± 0.12	114.8± 0.48

Positive control Glibenclamide 5mg/kg	112.8± 0.6	81.2± 0.67	500.4± 0.27	327.4± 0.3	211± 0.7	146± 0.23
200mg/kg extract	114.2±0.12	95.8± 0.84	369.2± 0.4	237.6± 0.23	137.6± 0.73	106.6± 0.42
400mg/kg extract	115± 0.11	122 ± 0.63	438.8± 0.47	267± 0.18	149.2± 0.68	89.4± 0.28*

Data are presented as Mean ± SEM; superscripts \* represents the level of significance at  $p < 0.05$  when compared to positive control. Repeated Measures ANOVA followed by turkey, test n=5

IGLBE = Initial Glucose Level before Induction, GLAI = Glucose Level after Induction, Positive control (Glibenclamide), Negative control N. saline = Normal saline




Fig: 1 Effect of extract and standard drug on experimental animals

## Discussion

This present research has shown that the methanol extract of *F. exasperata* leaves possess antidiabetic potentials due to its effect in lowering the blood glucose level of the alloxan induced diabetic rats. The effect is attributed to the phytochemical constituent such as phenols, flavonoids, alkaloids, tannins, saponins, steroids and triterpenes present in the plant extract as previously reported [11 - 18]. The result of acute toxicity study of extract orally administered showed no mortality at doses less or equal to 5000mg/kg which is in line with previous study [19]. This suggests that the methanol extract is safe for ethnopharmacological research but hepato-renal and hematological parameters must be established before administering it to humans.

The methanol leaves extract of *Ficus exasperata* significantly decreased the blood glucose level as observed in this present study. At 400mg/kg body weight there is a significant difference at  $p < 0.05$  when compared to standard drug whereas no significant difference occurred between 200mg/kg dose and standard drug. The result corroborates earlier findings that established the blood sugar lowering effect of aqueous leaves extract of *Ficus exasperata* [20-22]. From our findings, as the treatment progresses a significant decrease is observed due to increase in the concentration of extract administered. Therefore, the antidiabetic activity of the methanol extract follows a dose dependent manner. It is worthy of note that the use of the plant should be monitored at intervals as there is a tendency of causing hypoglycemic condition when consumed in large amounts. We recommend 200mg of the methanol extract as optimum dose to be administered. The significant decrease in blood glucose levels suggested that the extract contains therapeutic phytoconstituents with antidiabetic potentials. It has been established that the possible mechanism by which plant extract brings about its blood glucose lowering action is by potentiating the insulin effect and by increasing the pancreatic secretion of insulin from  $\beta$  cells. This lowering blood sugar effect observed might be the synergistic action of the phytochemical constituents present in *F. exasperata* leaves. The results obtained

from this study suggest that the methanol leaves extract of *Ficus exasperate* contained bioactive constituents that may have relevant antidiabetic properties used in the management of diabetes mellitus.

### **Conclusion**

The methanol extract of *F. exasperata* leaves has significant anti-diabetic activity. This is as a result of present of phytochemical constituent with proven therapeutic potentials. The study also established safety of the leaves extract with respect to doses administered. These observations justified the folkloric use of the plant in management of diabetic diseases. Further studies are needed to establish the hepato-renal and hematological effects of the methanol extract and also to isolate the active ingredient responsible for the observed anti-diabetic activity of the leaves extract.

### **REFERENCES**

1. Oyinlola OO, Lyndy JM, Ilse Janse van Rensburg, Jacobus NE and Vinny N. Antidiabetic activity of the ethyl acetate fraction of *Ficus lutea* (moraceae) leaf extract: comparison of an in vitro assay with an in vivo obese mouse model. BMC complementary and alternative medicine. 2016; 16:110. Doi.org/10.1186/s12906-016-1087-z
2. N'guessan K, Amoikon, KE and Soro D. Effect of aqueous extract of *Persea americana* seeds on the glycemia of diabetic rabbits. European journal of scientific research. 2014; 26, 376-385.
3. Lloyd A, Sawyer W, and Hopkinson P. Impact of long-term complications on quality of life in patients with type 2 diabetes not using insulin. Value in Health. 2001; 4(5), 392-400. Doi.org/10.1046/j.1524-4733.2001.45029.x

4. International diabetes federation (IDF). The diabetes atlas 10th edition update 2021  
[www.idf.org/diabetesatlas](http://www.idf.org/diabetesatlas). accessed 18/04/ 2022
5. Sofowora A, Ogunbodede E and Onayade A. The role and place of medicinal plants in the strategies for disease prevention. African journal of traditional, complementary, and alternative medicines. 2013; 10(5), 210–229. Doi:10.4314/ajtcam.v10i5.2
6. Mee-young L, Chang-Seob S, In-shik S, Young-bum K, Jung-hoon K and Hyeun-kyoo S. Evaluation of oral subchronic toxicity of soshiho-tang water extract. The traditional herbal formula in rats. Evidence-based complement. altern. Med. 2013; 2:1–9. Doi/org/10.1155/2013/590181
7. Firenzuoli F and Gori L. Herbal medicine today: clinical and research issues. Evidence based complement. altern. med. 2007; 4(S1):37–40. Doi.org/10.1093/ecam/nem096
8. Burkill HM. The useful plants of tropical West Africa. 3rd ed. kew: royal botanic gardens; 1997; 166–17
9. Tropical plants database, ken fern. tropical.theferns.accessed 18/04/22
10. Muibat OB, Misbaudeen A and Precious O. Nutrient and anti-nutrient phytochemicals in *Ficus exasperata* vahl leaves. International journal of scientific and engineering research 2014;5(1),2177-2181
11. Woode E, Poku RA, Ainooson GK, Boakye-gyasi E, Abotsi WK and Mensah TL. An evaluation of anti-inflammatory, antipyretic and antinociceptive effects of *Ficus exasperata* (vahl) leaf extract. J. pharmacol toxicol.2009; 4(4),138-51.
12. Adebayo EA, Ishola OR, Taiwo OS, Majolagbe ON and Adekeye BT. Evaluations of the methanol extract of *Ficus exasperata* stem bark, leaf and root for phytochemical

- analysis and antimicrobial activities. African journal of plant science. 2009; vol. 3 (12), pp. 283-2
13. Adebayo SO, Adenowo T, Naicker T, Ojewole JA. hypoglycaemic and hypotensive effects of *Ficus exasperata* vahl. (moraceae) leaf aqueous extract in rats. Afr j tradit complement altern med, 2011; 8:275-283.
  14. Abotsi WK, Woode E, Ainooson GK, Amo-Barimah AK, Boakye-Gyasi E. Antiarthritic and antioxidant effects of the leaf extract of *Ficus exasperata* p. beauv. (moraceae). pharmacognosy res. 2010; 2:89-97
  15. Banu KS and Cathrine L. General techniques involved in phytochemical analysis. *International Journal of Advanced Research in Chemical Science*, 2015 2 (4), 25-32
  16. Gusthinnadura OD, Achala TA and Malalige MW. Extraction methods, qualitative and quantitative techniques for screening of phytochemicals from plants. Nat Prod, 2017 5(2):29-32
  17. Lorke D. A new approach to practical acute toxicity testing. Archives of toxicology. 1983; 54: 275-287
  18. Malviya N, Jain S and Malviya S. Antidiabetic potential of medicinal plants. Acta Poland Pharmaceuticals. 2010; 67(2): 113-118.
  19. Victoria NU, Emmanuel EI, Daniel LA, Earnest OE, Goodies EM and Peter AA. Wound healing activity of the aqueous leaf extract and fractions of *Ficus exasperata* (moraceae) and its safety evaluation on albino rats. Journal of Traditional and Complementary Medicine. 2014; vol. 4, no. 4, pp. 246-252. Doi.org/10.4103/2225-4110.139105
  20. Adeyi AO, Idowu AB, Mafiana CF, Oluwalana SA and Ajayi OL. Effects of aqueous leave extract of *Ficus exasperata* on pathophysiology and histopathology of alloxan-induced diabetic albino rats. Journal of medicinal plants research 2012; vol. 6(46), pp. 5730-5736. Doi.org/10.5897/JMPR10.163

21. Toyin MY, Ahmad K. Salau OBO, and Musbau AA. Effect of aqueous leaf extract of *Ficus exasperata* in alloxan-induced diabetic wistar rats. Cameroon journal of experimental biology. 2014; vol.10 (1) 35-43. Doi.org/10.4314/cjeb.v10i1.5
22. Kazeem MI, Oyedapo BF, Raimi OG and Adu OB. Evaluation of *Ficus exasperata* leaf extracts in the management of diabetes mellitus in vitro. j. med. Sci. 2013; 13(4);269-275. Doi.org/10.3923/jms.2013.269.275

UNDER PEER REVIEW