

# Effect of Plant Growth Promoting Bacteria on Growth and Nutrient Content of Rice

---

## ABSTRACT

**Aims:** Application of large number of chemical fertilizers for crop production alters sustainability of the environment and creates pollution. This problem can be minimized by the application of Plant Growth Promoting Bacteria (PGPB). Plant Growth Promoting Bacteria (PGPB) are beneficial bacteria that promote plant growth and development through a variety of mechanisms. Therefore, a pot experiment was conducted at the department of Soil Science, BSMRAU to investigate the effect of different PGPB isolates on growth, nutrients content and uptake by rice.

**Materials and Methods:** The experiment was carried out following a completely randomized design (CRD) with four replicates. Twenty bacterial isolates were used as treatments with a control treatment. Thirty-five days old seedlings of rice were inoculated with different bacterial isolates for thirty minutes and planted in plastic pot. Broth culture was applied at one month interval and crops were harvested at 65 days after planting.

**Results:** Experimental results disclose that application of PGPB isolates resulted in a significant increase in plant height, SPAD value, root length, root volume, straw fresh weight, straw dry weight, root fresh weight, root dry weight, nutrient content and uptake by the rice plant compare to control. The highest values of most of the parameters were recorded from the plant inoculated with the isolate BU Ls 28.

**Conclusion:** The present study suggests that the use of PGPB isolate BU Ls 28 might be used as suitable inoculant for rice production due to the ability of this isolate to promote plant growth.

*Keywords: PGPB, isolate, growth, nutrient content, rice.*

## 1. INTRODUCTION

Rice (*Oryza sativa*) is an important cereal crop being a staple food for a large part of the world's human population including Bangladesh. In terms of production, Bangladesh is the fourth largest rice producer in the world [1]. In the fiscal year of 2019-2020 total production of rice in Bangladesh was estimated at about 36,604 metric tons from 28213 acres [2]. In Bangladesh, about 74.85 % of the total cultivable land is used for rice cultivation [3]. The contribution of rice in the daily dietary energy is almost 69 per cent and in the daily protein intake contribution of rice is 55 percent [1]. Rice production depends on large amount of

chemical fertilizers which leads to health hazard and environmental pollution for the rice growing areas. Therefore, for maintaining environmental sustainability, there is a need to think about eco-friendly approaches in order to minimize the use of chemical fertilizers by application of beneficial microorganisms [4]. Soil is abundant with many types of microorganisms and microorganisms especially in the rhizosphere play an important role in plant growth and development through various mechanisms [5]. Plant growth promoting bacteria (PGPB) that are beneficial to the plant, are capable of associating with many plants and are mainly classified into two types: free-living rhizosphere bacteria and endophytic bacteria [6, 7]. Plant growth promoting bacteria can directly cause seed emergence, plant growth, or improvement in crop yields by generating plant growth regulators such as auxins, gibberellins (GAs), and cytokinins, supplying biologically fixed nitrogen, and increasing the phosphorous uptake by solubilization of inorganic phosphates [8] while indirect mechanisms that involve suppression of bacterial, fungal, viral, and nematode pathogens [9]. Microbial population is closely related with the improvement of soil fertility [10] which ultimately favors the growth of the plant. PGPB have been applied to various crops to enhance growth, seed emergence and crop yield. In the findings of [11] who indicated that Plant growth-promoting rhizobacteria (PGPR) strains provided a significant increase in shoot and root length, and shoot and root biomass of rice. A significant increase in shoot N content (up to 76%) and root N content (up to 32%) was observed over the un-inoculated control. [12] reported that root length, shoot length, fresh and dry biomass of rice were significantly increased over control due to inoculation with different *Rhizobium* strains. It has been demonstrated that most of the parameters i.e. number of tillers/plant (46%), yield (43%), plant biomass (18%), straw dry weight (45%) and 1000-grain weight (25%) were improved in rice due to inoculation with *R. leguminosarum* [13]. Very little work has so far been done in Bangladesh regarding assess the effect of PGPB on growth of rice in Bangladesh. Therefore, a pot experiments was conducted to assess the effectiveness of PGPB on growth enhancement, nutrient content in rice plant as well as nutrient uptake by the rice plant.

## 2. MATERIAL AND METHODS

### 2.1 Experimental details

A pot experiment was carried out in the net house of the Department of Soil Science, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur-1706, Bangladesh during boro rice growing season of 2017 (9 February to 11 May 2017). Geographically the experimental site is located at 24.09° North Latitude and 90.25° East Longitudes with an elevation of 8.2 m from mean sea level [14].

Soil used for the experiment was collected from the experimental field of the Department of Soil Science, BSMRAU. The soil belongs to Shallow Red Brown Terrace Soil (Salna series) of the Madhupur Tract (AEZ 28) and is classified as Inceptisols [15]. This soil is characterized by clay within 50 cm of the surface and is slightly acidic in nature. The pH value of the soil sample collected from the study site was 5.70. This soil contained 0.75% organic carbon (OC), 1.30% organic matter (OM), 0.07% nitrogen (N), 2.57 ppm phosphorus (P) and 0.212 meq 100 g<sup>-1</sup> soil exchangeable potassium (K). Collected soil sample was sterilized by autoclaving. An amount of 1 kg soil was placed into a 12.9 cm x 12.7 cm plastic pot. The experiment was conducted in CRD design with four replications.

Twenty bacterial isolates viz. BU Ca 14, BU Ca 15, BU Ca 16, BU Ca 17, BU Ps 6, BU Ps 7, BU Ps 8, BU Ps 9, BU Ls 26, BU Ls 27, BU Ls 28, BU Ls 29, BU Ls 30, BU Ls 31, BU Le 16, BU Le 17, BU Le 18, BU Le 19, BU Le 20 and BU Le 21 isolated from the different leguminous crop viz. chickpea (*Cicer arietinum*), pea (*Pisum sativum*), grasspea (*Lathyrus sativus*) and lentil (*Lens esculentus*) were collected from Soil Microbiology Laboratory of

BSMRAU. A loopful of the respective PGPB isolate was transferred to the liquid medium of 100 mL conical flask and incubated for 2 days on a rotary shaker. Two days old broth culture of *Rhizobium* isolates were centrifuged at 8000 rpm, supernatant was discarded then the cell pellet was re-suspended in sterile distilled water. Bacterial cells per milli liter was adjusted ( $10^8$ cfu /ml) by taking absorbance reading using spectrophotometer. An un-inoculated control was also maintained. BRR1 dhan 28, a high yielding rice variety was used as the test crop in the boro season. Thirty five days old rice seedlings were washed with sterile water seven times and then treated with bacterial isolates for 30 min. Inoculated seedlings with each strain were transplanted into the pots. Un-treated seedlings were designated as control. Broth culture of *Rhizobium* isolates were also applied in root zone after transplantation of rice up to tillering stage @ 1 ml / plant at one month interval. Hoagland solution was applied to soil for supplying plant nutrients. All intercultural operations like weeding, irrigation etc. were done as and when needed.

## 2.2 Harvesting of the plants and analysis

Rice plants were harvested after 65 days of planting through separating of plants from soil. The plants were washed through dipping into a vessel. Plant height (cm plant<sup>-1</sup>), leaf greenness (SPAD Value), root length (cm), root volume (cm<sup>3</sup>), fresh weight and dry weight of shoot and root (g) of each plant was recorded.

Dry weight of straw and root were recorded after drying in an oven for 48 hours at 70°C. Total nitrogen content of plant samples was estimated by Micro-kjeldahl method [16]. Phosphorus in digested samples was determined by Vanadomolybdophosphoric yellow color method [17] by using spectrophotometer at 440 nm wavelength. Potassium in the aliquot was estimated with the help of flame photometer after appropriate dilution [16]. Sulfur content in the digest was determined by adding acid solution and then precipitation with BaCl<sub>2</sub> and measuring the turbidity calorimetrically at 420 nm wavelength [18].

The nutrient uptake by rice straw was measured by using the following formula:

$$\text{Nutrient Uptake} = \frac{\% \text{ Nutrient} \times Y \text{ (g plant}^{-1}\text{)}}{100}$$

Here,

Y (g plant<sup>-1</sup>) = Total dry matter production of plant


Data was analyzed statistically by using the computer package Statistix 10 program. Mean separation was done by LSD (Least significant Difference Test) at 5% level of probability. Computation and preparation of graphs were made by the use of Microsoft Excel 2010 program.

## 3. RESULTS AND DISCUSSION

### 1. 3.1 Influence of PGPB on growth parameters of rice

#### 3.1.1 Plant height


Results presented in Figure 1 show that all the treatments produced significantly higher plant height over the control treatment where no bacterial isolate was applied. The highest plant height (80.48 cm) was obtained from treatment receiving isolate BU Ls 28. The effect of BU Ls 28 was statistically similar to the isolates BU Ca 14, BU Ca 16, BU Ps 6, BU Ps 7, BU Ps 9, BU Ls 26, BU Ls 29, BU Ls 30 and BU Ls 31. The lowest plant height (54.95 cm) was obtained from the control treatment. Higher plant height in treatments receiving PGPB isolates might be due to production of growth promoting substances by the bacterial isolates. PGPR can affect plant growth through production and release of different types of secondary metabolites (plant growth regulators/ phytohormones/ biologically active substances), preventing deleterious effects of phytopathogenic organisms in the rhizosphere and/or facilitating the availability and uptake of certain nutrients from the root environment [19]. This result corroborates with the finding of [12] who found higher plant height with *Rhizobium* inoculation in rice.



**Fig. 1. Influence of plant growth promoting bacteria on plant height of rice (Error bar indicates the standard deviation of mean)**

### **3.1.2 Leaf greenness (SPAD value)**

All the treatments produced significantly higher SPAD value over control (Figure 2). The highest leaf greenness (46.1) was obtained from the treatment receiving isolate BU Ls 28 while the lowest (36.5) was found from control (without inoculation). The effect of BU Ls 28 was statistically identical to the isolates BU Ca 14, BU Ls 26, BU Ls 29 and BU Ls 31. The SPAD value is correlated with leaf chlorophyll content which represents the greenness of leaf. It has been demonstrated that leaf nitrogen content significantly influences the SPAD value [20]. The nitrogen fixing ability from the atmosphere and nutrient uptaking capability from the root zone of the PGPB might have enhanced chlorophyll content in rice plant as compared to the un-inoculated control treatment. This result corroborates with the findings of [21] who reported significant increase in SPAD values in rice with *Rhizobium* inoculation.



**Fig. 2. Influence of plant growth promoting bacteria on leaf greenness (SPAD Value) of rice (Error bar indicates standard deviation of mean)**

### **3.1.3 Root length**

Inoculation of plant with different bacterial isolates resulted in higher root length compared to uninoculated control plants (Table 1). The maximum root length was found in treatment receiving isolates BU Ls Ca 14 (26.75 cm) followed by the treatment receiving BU Ls 28 (24.18 cm) and BU Ls 26 (23.25 cm). The lowest root length (14.6 cm) was obtained from the uninoculated control treatment. The increased root length in *Rhizobium* inoculated treatments might be due to production of phytohormones by the bacterial isolates which encourage root elongation in rice. This result corroborates with the findings of [22] who reported that *Rhizobium* inoculation significantly increased the root length of rice. These results are also in harmony with that of [12] who found higher root length with *Rhizobium* inoculation in rice. [23] also reported significantly higher root length of chickpea with bacterial inoculation as compared to the control treatment.

### **3.1.4 Fresh biomass yield of shoot and root**

Fresh biomass yield of shoot and root of rice plant was increased significantly over control due to different plant growth promoting bacterial inoculation (Table 1). The highest fresh biomass yield of rice shoot (52.35 g) and root (26.54g) were obtained from treatment receiving isolate BU Ls 28. In case of shoot, the lowest (16.15 g) biomass yield was obtained from control treatment. The lowest root fresh biomass (4.26 g) was also obtained from the control treatment. This result correlated with the finding of [12] who reported that inoculation of rhizobia to rice produced significantly higher shoot fresh biomass yield.

### 3.1.5 Dry biomass yield of shoot and root

A significant increase in dry biomass yield of rice shoot and root was observed in bacterial inoculated plants in comparison to the untreated control plants (Table 1). The highest dry biomass yield both for shoot (12.56 g) and root (6.37 g) were obtained from treatment receiving BU Ls 28 followed by BU Ca 14 and BU Ls 26. The lowest shoot biomass (3.88 g) and root biomass (1.02 g) were obtained from control (Table 1) treatment where no PGPB isolate was applied. This result was at par with the findings of [24] who reported that PGPR inoculants increase shoot and root dry matter yield in rice which also corroborates with the findings of [13]. The increase in the shoot and root dry mass directly correlates with the increased plant growth promotion due to the bacterial inoculation. In our investigation, bacterial treatments increased plant height, leaf greenness, root length and root volume thereby suggesting that the amount of photosynthates produced in rice plant may have been higher in the bacterial inoculated plants in comparison to the uninoculated control plants, thus, resulting in an increase in the shoot and root biomass of the rice plants. Moreover, PGPB changes the physico-chemical properties of the soils surrounding the host plants root which facilitate better nutrient acquisition by the host plant at the cost of minimum input [24] and therefore obtained higher dry biomass yield of rice plants. [25] also reported higher dry matter yield of bottle gourd due to inoculation with *Trichoderma*.

**Table 1. Influence of plant growth promoting bacteria on root length, fresh and dry biomass yield of rice**

PGPB isolates	Root length (cm)	Shoot Fresh Weight (g)	Root fresh weight (g)	Shoot Dry Weight (g)	Root Dry Weight (g)
Control	14.6 j	16.15 n	4.26 m	3.88 l	1.02 m
BU Ca 14	26.75a	41.88 b	19.98 c	10.05 b	4.59 c
BU Ca 15	20.5fg	31fgh	11.32 g	7.44ghi	2.60 h
BU Ca 16	21.25 def	36 d	14.28 f	8.64 de	3.28 f
BU Ca 17	17.75i	24.61 kl	7.95ijk	5.91 j	1.83jkl
BU Ps 6	20.63ef	32.3efg	11.79g	7.75fgh	2.83gh
BU Ps 7	20.93ef	32.75ef	12.13g	7.86fgh	2.91 g
BU Ps 8	17.98i	23.9 kl	7.46jkl	5.92 j	1.79 kl
BU Ps 9	20.8ef	32.55ef	12.04g	8.06efg	2.89 g
BU Ls 26	23.25bc	40.875bc	21.82 b	9.81bc	5.24 b
BU Ls 27	17.65i	21.75lm	6.54 l	5.22jk	1.57 l
BU Ls 28	24.18b	52.35a	26.54a	12.56 a	6.37a
BU Ls 29	22.88bcd	38 cd	17.93 d	9.12 cd	4.30 d
BU Ls 30	21.23 def	35 de	13.27 f	8.4 def	3.18 f
BU Ls 31	22.25cde	36.63 d	16.44 e	8.79 de	3.95 e
BU Le 16	20.13fgh	30.48fghi	9.63 h	7.31ghi	2.31i
BU Le 17	18.5hi	28.88hij	8.99 hi	7.22 hi	2.07ij
BU Le 18	18.88ghi	29.28ghij	9.06 hi	7.32ghi	2.08ij
BU Le 19	17.35i	19mn	6.85 kl	4.75 k	1.58 l
BU Le 20	18.1i	27.75ij	8.67hi	6.94i	1.99jk
BU Le 21	18.05i	26.75jk	8.08ij	6.69i	1.86jk
SE ( $\pm$ )	0.85	1.57	0.56	0.38	0.13
CV (%)	5.95	7.07	6.50	7.08	6.50

SE and CV mean Standard Error and co-efficient of variation respectively. Means followed by common letter(s) are not significantly different at 5% level of probability by LSD

## **3.2 Influence of PGPB on nutrient content and nutrient uptake by rice plant**

### **3.2.1 Nitrogen content and uptake by rice shoot**

Plant growth promoting bacterial inoculation had positive effect on nitrogen content and uptake by rice shoot (Tables 2-3). The highest nitrogen content in shoot (1.80 %) was obtained from treatment receiving isolate BU Ls 28. The effect of isolate BU Ls 28 was statistically similar to the rest of the treatments except BU Le 20, BU Le 21 and control. The lowest nitrogen content in rice shoot (1.00%) was obtained from control treatment. This result was in harmony with that of [13] who reported that nitrogen content of rice grain was increased significantly with inoculation of *Rhizobium* isolates as compared to uninoculated control. Our findings are also supported by the findings of [26], [27] who found higher nitrogen content in *Rhizobium* inoculated chickpea shoot as compared to the un-inoculated control plants.

The highest nitrogen uptake by rice shoot (225.78 mg/plant) was obtained from treatment BU Ls 28 followed by BU Ca 14 and BU Ls 26. The lowest nitrogen uptake by rice shoot (38.94 mg /plant) was obtained from control treatment where no bacterial isolates were applied. Higher nitrogen content in rhizobacterial inoculated treatments might be due to fixation of atmospheric nitrogen by rhizobacteria and which eventually might lead to higher nitrogen content in rice plant and also increased N uptake by increased availability of this element by different metabolites produced by rhizobacteria. [11] found higher nitrogen content in shoot of rice plant due to *Rhizobium* inoculation which also corroborates with the findings of [28] as well. These results are also in harmony with that of [29] who reported that N uptake was increased by rice plants inoculated with rhizobia.

### **3.2.2 Phosphorus content and uptake by rice straw**

Results presented in Table 2 and 3 reveal that phosphorus content and uptake by rice shoot were increased significantly over control due to inoculation of plant growth promoting bacteria. Treatment receiving isolate BU Ls 28 showed higher phosphorus content in rice shoot (0.489%) whose effect was statistically similar to BU Ca 14, BU Ca 16, BU Ls 26, BU Ls 29 and BU Ls 31. The lowest phosphorus content in rice shoot (0.286%) was obtained from control. The highest phosphorus uptake by rice shoot (61.35 mg/plant) was also obtained from BU Ls 28 followed by BU Ca 14 and BU Ls 26. The lowest phosphorus uptake by rice shoot (11.09 mg/ plant) was obtained from control treatment. This result is in agreement with that of Solaiman et al. (2010) [30] who found higher phosphorus content and uptake by shoot of chickpea due to *Rhizobium* inoculation. Higher phosphorus content and uptake by shoot as reported might be due to increased availability of this element by different mechanisms mediated by rhizobacteria. Soil of the present study was slightly acidic in reaction. In acidic soil, phosphorus fixation is a common phenomenon. It has been reported that phosphorus solubilizing ability of the PGPB [9] which might improve P availability in the studied soil and thereby obtained higher P content as well as higher P uptake by the rice plants inoculated with PGPB isolates as compared to the un-inoculated control plants. The result corroborates with the findings of [13] who reported that phosphorus content of rice grain increased significantly with inoculation of *Rhizobium* strains as compared to un-inoculated control.

**Table 2. Influence of plant growth promoting bacteria on nitrogen, phosphorus, potassium and sulfur contents in rice shoot**

<b>PGPB isolates</b>	<b>%N</b>	<b>%P</b>	<b>% K</b>	<b>%S</b>
Control	1.00 c	0.286i	1.41 g	0.139 n
BU Ca 14	1.79 a	0.486a	2.6a	0.362 ab
BU Ca 15	1.61ab	0.372g	2.02 def	0.244 kl
BU Ca 16	1.71ab	0.473ab	2.42 ab	0.331cdef
BU Ca 17	1.61ab	0.373g	2 def	0.247 kl
BU Ps 6	1.67ab	0.442cd	2.34abc	0.302fgh
BU Ps 7	1.70ab	0.456bc	2.38 ab	0.316 def
BU Ps 8	1.72ab	0.457bc	2.39 ab	0.329cdef
BU Ps 9	1.67ab	0.447c	2.35abc	0.311efg
BU Ls 26	1.75ab	0.488 a	2.44 ab	0.352abc
BU Ls 27	1.67ab	0.425 de	2.28bcd	0.286ghi
BU Ls 28	1.80a	0.489 a	2.62 a	0.382 a
BU Ls 29	1.73ab	0.484 a	2.42 ab	0.341bcd
BU Ls 30	1.58ab	0.366 g	1.97ef	0.237 kl
BU Ls 31	1.73ab	0.475 ab	2.42 ab	0.334bcde
BU Le 16	1.65ab	0.408ef	2.28bcd	0.279hij
BU Le 17	1.62ab	0.394 f	2.08cdef	0.261 ijk
BU Le 18	1.62ab	0.402 f	2.18bcde	0.266ijk
BU Le 19	1.62ab	0.400 f	2.04 def	0.250jk
BU Le 20	1.57 b	0.358 g	1.89 f	0.218 l
BU Le 21	1.56 b	0.320 h	1.91ef	0.182 m
SE ( $\pm$ )	0.09	0.01	0.14	0.015
CV (%)	8.32	3.45	9.22	7.49

SE and CV mean Standard Error and co-efficient of variation respectively. Means followed by common letter(s) are not significantly different at 5% level of probability by LSD

### **3.2.3 Potassium content and uptake by rice shoot**

Potassium content and uptake by rice shoot were increased significantly over control due bacterial inoculation (Tables 2-3). The highest potassium content in rice shoot (2.62%) was obtained from treatment receiving isolate BU Ls 28. The effect of BU Ls 28 was statistically similar to BU Ca 14, BU Ca 16, BU Ps 6, BU Ps 7, BU Ps 8, BU Ps 9, BU Ls 26, BU Ls 29 and BU Ls 31. The lowest potassium content in rice shoot (1.41 %) was obtained from control. The highest potassium uptake by rice shoot (328.40 mg/plant) was obtained from treatment receiving isolate BU Ls 28 followed by BU Ca 14 and BU Ls 26. The lowest potassium uptake (54.48 mg/ plant) by rice shoot was obtained from uninoculated control treatment whose effect was statistically inferior to all others treatments receiving different PGPB isolates. Higher potassium content and uptake as found might be due to increased availability of this element by the activity of rhizobacteria. This result is in harmony with that of [31] who found higher potassium content in rice shoot due to *Rhizobium* inoculation. These results corroborate with the findings of [32] who reported that application of *Rhizobium* enhanced the K-uptake by pea.

**Table 3. Influence of plant growth promoting bacteria on nitrogen, phosphorus, potassium and sulfur uptake by rice shoot**

<b>PGPB isolates</b>	<b>N (mg/ plant)</b>	<b>P (mg/ plant)</b>	<b>K (mg/ plant)</b>	<b>S (mg/ plant)</b>
Control	38.94 m	11.09 j	54.48 l	5.40 k
BU Ca 14	179.76 b	48.81 b	261.49 b	36.32 b
BU Ca 15	119.89 ghi	27.63efg	150.31 hi	18.15 hi
BU Ca 16	147.64 def	40.82 c	208.56 de	28.57 de
BU Ca 17	94.99jkl	22.02 hi	118.12jk	14.55 j
BU Ps 6	129.21 fgh	34.22 d	180.76 fg	23.41fg
BU Ps 7	133.32efg	35.80 d	187.27ef	24.82 f
BU Ps 8	101.45ijk	27.01fg	141.03hij	19.51 h
BU Ps 9	135.22efg	35.95 d	188.93ef	25.02ef
BU Ls 26	171.89bc	47.88 b	240.19bc	34.57 bc
BU Ls 27	87.13 kl	22.19 hi	119.02 jk	14.93 ij
BU Ls 28	225.78 a	61.35 a	328.40 a	47.81 a
BU Ls 29	157.70 cd	44.10 c	220.30 cd	31.13 cd
BU Ls 30	132.68efg	30.73 e	165.48fgh	19.91 gh
BU Ls 31	152.38cde	41.74 c	212.02 de	29.40 d
BU Le 16	121.19 ghi	29.86ef	166.64fgh	20.39gh
BU Le 17	116.66ghi	28.41ef	150.29 hi	18.83 h
BU Le 18	118.46 ghi	29.42ef	159.29 gh	19.50 h
BU Le 19	76.75 l	19.02 i	96.85 k	11.87 j
BU Le 20	109.21hij	24.81 gh	131.3ij	15.12 ij
BU Le 21	104.74ijk	21.37 hi	127.73 ij	12.17 j
SE ( $\pm$ )	10.29	1.72	13.61	1.79
CV (%)	11.52	7.48	11.21	11.29

SE and CV mean Standard Error and co-efficient of variation respectively. Means followed by common letter(s) are not significantly different at 5% level of probability by LSD

### **3.2.4 Sulfur content and uptake by rice shoot**

Sulfur content and uptake by rice shoot were influenced significantly due to inoculation of plant growth promoting bacteria (Tables 2-3). Treatment receiving isolate BU Ls 28 showed higher sulfur content in shoot (0.382 %) of rice. This result is in harmony with [28] who reported higher sulfur content in shoot of rice due to *Rhizobium* inoculation. The effect of isolate BU Ls 28 was statistically similar to the treatments BU Ca 14 and BU Ls 26. The lowest sulfur content (.139%) was obtained from control. The highest sulfur uptake by shoot (47.81 mg/plant) of rice was obtained from *Rhizobium* isolate BU Ls 28. The lowest sulfur uptake (5.40 mg/plant) was found in control treatment. [28] reported higher sulfur uptake by shoot of rice due to *Rhizobium* inoculation. We observed an increase in shoot sulfur content as well as higher sulfur uptake by the rice shoot due to PGPB inoculation over un-inoculated control. The enhanced sulfur content and uptake might be due to the enhanced root volume due to bacterial inoculation thereby leading to an increase in the translocation of sulfur from root to shoot, consequently shoot sulfur content was also found to be increased. Additionally, higher dry biomass yield of rice plant due to bacterial inoculation might be responsible for higher nutrient uptake by the rice plant.

#### 4. CONCLUSION

Results of the present study reveal that inoculation of plant growth promoting rhizobacteria resulted in higher vegetative growth and also facilitated increased dry matter accumulation of the rice plants. In addition of improving growth of the plants, inoculation with PGPR also resulted in improving the nutrient content in rice plant as well as ensure higher nutrient uptake by the plant. Among the twenty bacterial isolates, BU Ls 28 performed better than all other isolates in improving the different growth parameters, nutrient content and uptake by the rice plant which may imply that this isolate could be used as PGPB for promoting rice growth followed by BU Ca 14 and BU Ls 26. Therefore, the study provides an insight of utilizing PGPB isolates as sustainable and eco-friendly approach for rice cultivation with reduced inorganic nitrogenous fertilizer. However, further studies in field level are needed to confirm their potential.

#### REFERENCES

1. Mottaleb KA. Why hybrid rice is not gaining popularity in Bangladesh. *The Financial Express*. Dhaka. 2020.
2. BBS. *Statistical Year Book*. (2019). Bangladesh Bureau of Statistics, Dhaka, Bangladesh. 2020.
3. BBS. *Statistical Year Book*. (2020). Bangladesh Bureau of Statistics, Dhaka, Bangladesh. 2021.
4. Rahman MM, Alam MS, Islam MM, Kamal MZU, Rahman GKMM, Haque MM, Miah MG, Biswas JC. Potential of legume-based cropping systems for climate change adaptation and mitigation. In *Advances in Legumes for Sustainable Intensification*. Academic Press. 2022;381-402.
5. Rahman MM, Alam MS, Kamal MZU, Rahman GKMM. Organic sources and tillage practices for soil management. In *Resources Use Efficiency in Agriculture*. Springer, Singapore. 2020;283-328.
6. Kloeppel JW, Rodriguez-Kabana R, Zehnder AW, Murphy JF, Sikora E, Fernandez C. Plant root-bacterial interactions in biological control of soil borne diseases and potential extension to systemic and foliar diseases. *Australas Plant Pathol*. 1999;28(1):21-26.
7. Gray EJ, Smith DL. Intracellular and extracellular PGPR: commonalities and distinctions in the plant-bacterium signaling processes. *Soil Boil. Biochem*. 2005; 37(3):395-412.
8. Podile AR, Kishore GK. Plant growth-promoting rhizobacteria. In *Plant-associated bacteria*. Springer, Dordrecht. 2007;195-230.
9. Kirankumar R, Jagadeesh KS, Krishnaraj PU, Patil MS. Enhanced growth promotion of tomato and nutrient uptake by plant growth promoting rhizobacterial isolates in presence of tobacco mosaic virus pathogen. *Karnataka J. Agric. Sci*. 2010;21(2).

10. Rahman GKMM, Rahman MM, Alam MS, Kamal MZ, Mashuk HA, Datta R, Meena RS. Biochar and organic amendments for sustainable soil carbon and soil health. In Carbon and nitrogen cycling in soil. Springer, Singapore. 2020; 45-85.
11. Majeed A, Abbasi MK, Hameed S, Imran A, Rahim N. Isolation and characterization of plant growth-promoting rhizobacteria from wheat rhizosphere and their effect on plant growth promotion. *Front Microbiol.* 2015;6:198.
12. Solaiman ARM, Hossain GMA, Mia MAB. Effect of *Rhizobium* on growth and biomass production of rice. *Bangladesh J. Microbiol.* 2011; 28(2):64-68.
13. Hussain MB, Mehboob I, Zahir ZA, Naveed M, Asghar HN. Potential of *Rhizobium* spp. for improving growth and yield of rice (*Oryza sativa* L.). *Soil. Environ.* 2009; 28(1):49-55.
14. Anonymous. Annual weather report. IPSA Metrological station, Salna, Gazipur. 1989;6-17.
15. Brammer H. The geography of the soils of Bangladesh. The University Press Limited, Dhaka, 1996, pp 132–133.
16. Page AL, Miller RH, Keney DR. Methods of Soil Analysis. Part II, 2nd edn. American Society of Agronomy. Inc. Madison. Wisconsin, USA; 1989.
17. Jackson MZ. Soil Chemical Analysis, Practice Hall of India Private Limited. New Delhi, India; 1973.
18. Black CA. Methods of soil analysis, Part-I and II. Amer. Soc. Agronomy. Inc. Pub. Madison, Wisconsin, USA 1965;30-48.
19. Zahir ZA, Arshad M, Frankenberger WT. Plant growth promoting rhizobacteria: applications and perspectives in agriculture. *Adv. Agron.* 2003;81:97-168.
20. Xiong D, Chen J, Yu T, Gao W, Ling X, Li Y, Peng S, Huang J. SPAD-based leaf nitrogen estimation is impacted by environmental factors and crop leaf characteristics. *Sci. Rep.* 2015;5(1), 1-12.
21. Biswas JC, Ladha JK, Dazzo FB, Yanni YG, Rolfe BG. Rhizobial inoculation influences seedling vigor and yield of rice. *J. Agron.* 2000a;92(5):880-886.
22. Hari K, Srinivasan TR. Response of sugarcane varieties to application of nitrogen fixing bacteria under different nitrogen levels. *Sugar Tech* 2005;7(2):28-31.
23. Solaiman ARM, Rabbani MG, Hossain D, Hossain GMA, Alam MS. Influence of phosphorus and inoculation with *Rhizobium* and AM fungi on growth and dry matter yield of chickpea. *Bangladesh J. Sci. Res.* 2012;25(1):23-32.
24. Sharma A, Shankhdhar D, Shankhdhar SC. Growth promotion of the rice genotypes by PGPRs isolated from rice rhizosphere. *J. Soil Sci. Plant Nutr.* 2014;14(2):505-517.
25. Barua S, Molla AH, Haque MM, Alam MS. Performance of Trichoderma-enriched bio-organic fertilizer in N supplementation and bottle gourd production in field condition. *Hort. Internat. J.* 2018;2:106-114.
26. Alam MS, Rahman MM, Rahman GKMM, Islam MM. Evaluation of *Rhizobium* isolates in terms of nodulation, growth and yield of chickpea. *Bangladesh J. Soil Sci.* 2015;37(1):35-45.
27. Alam, MS, Solaiman ARM, Islam MM. Screening of *Rhizobium* for high nitrogen fixing potential in chickpea. *The Agriculturists.* 2010;8(1):75-84.
28. Biswas JC, Ladha JK, Dazzo FB. Rhizobia inoculation improves nutrient uptake and growth of lowland rice. *Soil Sci. Soc. Am. J.* 2000b;64(5):1644-1650.
29. Yanni YG, Rizk RY, Corich V, Squartini A, Ninke K, Philip-Hollingsworth S, et al. Natural endophytic association between *Rhizobium leguminosarum* bv. *trifolii* and

- rice roots and assessment of its potential to promote rice growth. *Plant Soil*. 1997;194:99-114.
30. Solaiman ARM, Talukder MS, Rabbani MG. Influence of some *Rhizobium* strains on chickpea: nodulation, dry matter yield and nitrogen uptake. *Bangladesh J. Microbiol*. 2010;27(2):61-64.
  31. Naher UA, Othman R, Shamsuddin ZH, Saud HM, Ismail MR. Growth enhancement and root colonization of rice seedlings by *Rhizobium* and *Corynebacterium* spp. *Int. J. Agric. Biol*. 2009;11:586-590.
  32. Solaiman ARM, Rabbani MG. Effects of *Rhizobium* inoculant, compost and nitrogen on edible podded pea. *Bangladesh J. Microbiol*. 2005;22(1):5-9.