

Bioactives and Pharmacology of *Acacia Nilotica*- A Review

ABSTRACT:

This review article aims to give information about the main phytochemicals present in *Acacia* and its many pharmacological activities. This review is helpful for researchers in the development of new drug products.

1. INTRODUCTION

Acacia is a genus of shrubs and trees belonging to the subfamily Mimosoideae of the family Fabaceae (Baravker et al., 2008).

They are pod-bearing, with sap and leaves typically bearing large amounts of tannins and condensed tannins that historically in many species found used as pharmaceuticals and preservatives. This name derives from the Greek word for its characteristic thorns, (*as*, thorn). The generic name derives from ἀκακία (*akakia*), the name given by early Greek botanist-physician Pedanius Dioscorides (ca. 40-90) to the medicinal tree *A. nilotica* in his book *Materia Medica* (Anonymous, 2003). The genus *Acacia* previously contained roughly 1300 species, about 960 of them native to Australia, with the remainder spread around the tropical to warm temperate regions of both hemispheres, including Europe, Africa, southern Asia, and America. The extract of pods or legumes is known as *Aqaqia* in the Unani system of medicine (Antaki AD et al, 1998) when it is obtained from unripe fruit then called *Qurz*. It is native to Egypt, and seen throughout the greater part of India, Ceylon, Baluchistan, Waziristan, Arabia, Egypt and tropical Africa (Brenan J.P, 1983).

Acacia nilotica known as babul is the most important tree in the dried parts of India. Almost all its parts are used in different aspects including root, bark, leaves, flower, gum, pods, etc. (Fosberg FR. Et al, 1980), The bark of the babul tree contains tannin made up of strong and astringent acid (gallic acid) and used in tanning and dyeing, inks and pharmaceuticals (Dymock W. et al, 2005). It is an evergreen tree, growing up to 10 meters in height. It is a small tree with dark brown or black longitudinally fissured bark; branchlets are slender,

terete, and pubescent when young. The tree generally attains a height of 15 m and girth of 1.2 m, though trees up to a height of 30 m with a girth of 3 m have also been recorded.¹ The leaves are 2-pinnate, 5-10 cm long; main rachis downy, often furnished with glands; petioles are 2.5-5 cm long; stipular spines very variable, 0.6-5cm long, and smooth.(Rashed I. Kitabul Kulliyat,1987) The leaflets are opposite in 10 to 20 pairs of crowded, sessile, linear-oblong, obtuse or acute, rigid, greyish green about 1/6 inch long.(Khan R .2009). The flowers are yellow, golden–yellow, fragrant, crowded in long-stalked, globed heads, 1.5 cm in diameter, forming axillary clusters of 2-5 heads,¹ and pubescent; bracteoles 2, calyx campanulate, 1.25 mm long; teeth very short. Corolla 3 mm long; lobes short, triangular.¹⁰ Pod shortly stalked, 3 or 4 inches long by about ¾ wide, more or less constricted between the 2-6 seeds, flat except over the seeds, smooth, pale membranous, with a strong fibrous marginal rib and fainter transverse reticulating veins. Seeds with a long funicle slightly dilated at the hilum, roundish in outline¹⁵, and persistently grey downy. It is very bitter to taste The gum exudes from the cuts in the bark in form of ovoid tears. The tears are glossy and marked with minute fissures and are brittle. The color of the gum varies from pale yellow to black. It is soluble in water (Gupta, R.K, 1970)




Figure 1: Picture of *Acacia nilotica*


2. BIOACTIVES

Acacia species contain secondary metabolites including amines and alkaloids, cyanogenic glycosides, cyclitols, fatty acids, seed oils, fluoroacetate, gums, nonprotein amino acids,


terpenes (including essential oils, diterpenes, phytosterol, and triterpene genins and saponins, hydrolyzable tannins, flavonoids, and condensed tannins. The plant is a richer source of cysteine, methionine, threonine, lysine, tryptophan, potassium, phosphorus, magnesium, iron, and manganese (Dymock W, 2005). The plant chemical compounds like diester, pentacosane dioic acid dihexadecyl ester and alcohol, heptacosane 1, 2, 3-triol. Seeds: It contains a high percentage of phenolic constituents consisting of m-digallic acid, gallic acid, protocatechuic and ellagic acids, leucocyanidin, mdigallic dimer 3,4,5,7-tetrahydroxy flavan-3-ol, oligomer 3,4,7- trihydroxy flavan 3,4-diol and 3,4,5,7tetrahydroxy flavan-3-ol and (-) epicatechol. The mature seed also contains crude protein, crude fiber, crude fat, carbohydrates, potassium, phosphorus, magnesium, iron, and manganese occurring in high concentrations and it is a richer source of cysteine, methionine, threonine, lysine, and tryptophan (Eline M.B, et al 2005). Fruit also contains mucilage and saponins. Pods: It contains gallic acid & its Me-este-n-digallic acid and condensed tannins. Leaf: It contains apigenin, 6-8-bisD-glucoside, rutin, and 8% digestive protein (12.4% crude protein). Relative levels of tannin in different parts of the plant are deseeded pods (50%), pods (5.4%), leaves (7.6%), bark (13.5%), and twigs (15.8%).Bark: It contains tannin (12-20%), terpenoids, saponins and glycosides, Phlobetannin, gallic acid, protocatechuic acid pyrocatechol, (+) – catechin, (-) epigallocatechin digallate (Bachayaa H.A, Et al 2009).Its extract contains total phenolic content ranging from 9.2 to 16.5 g/100 g. Root: It contain s octaconsanol, betulin, B-amyryn and Bsitosterol. Gum: It is composed of galactoaraban which gives on hydrolysis L-arabinose,, D-galactose, Lrhamnose, D-glucuronic acid, and 4-O-methyl- Dglucuronic acid (Ali A.J, Et al 2010).




Apigenin




Threonine




Kaempferol




Pyrocatechol




Tryptophan




Cystine




^{OH}Cyanidin



Catechin



Gallic acid



Lysine

Figure 2: Bioactives from *Acacia nilotica* Linn

3. PHARMACOLOGY

3.1. Antioxidant activity

Acacia species are a rich source of polyphenolic compounds, known to have strong antioxidant properties that help in the prevention and therapy of various oxidative stress-related diseases including cardiovascular, neurodegenerative, and cancer. Methanolic extract of the plant contains Kaempferol which is responsible for the antioxidant activity of the plant.

(Singh B.N, et al 2009).

3.2. Antiplatelet Aggregatory & Anti-quorum sensing

Activity: The extract of the plant block platelet aggregation mediated by platelet agonists, arachidonic acid, ADP, platelet-activating factor, and collagen. It blocks the aggregation of platelet induced by the Ca^{+2} -channel (Dymock W, 2005)

3.3. Antibacterial activity

The plant extract showed potent antibiotic activity against four bacterial species: gram-positive; *Bacillus subtilis*, *Staphylococcus albus*, *Streptococcus faecalis*; gram-negative, *Escherichia coli* and two fungal species: *Candida albicans* and *Aspergillus flavus* (Chaubal R, Tambe A, 2006)

3.4. Antifungal activity and Antiviral activity

Acacia nilotica species can be regarded as promising resources for antibacterial drugs due to their highly active nature. Antifungal activity of methanolic extracts and aqueous extract of *A. nilotica* with percentage inhibition ranging from 34.27 ± 1.45 to 93.35 ± 1.99 . Dried fruits of *Acacia nilotica* are active against *C. albicans* and are used to treat oral (Jansen, P.C.M, 2005). Methanolic extract of the plant is active against two animal viruses: Newcastle Disease and Fowlpox Viruses.

3.5. Antimicrobial Activity

Antimicrobial activity of the crude ethanolic leaf extract of *Acacia nilotica* Linn against *Campylobacter coli* isolated from goats. The highest zone of inhibition was observed with the 70 mg/ml concentration has studied the antimicrobial activity of ethanolic extracts of the stem bark against *Streptococcus viridans*, *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, and *Shigella sonnei* using the agar diffusion method and found the minimum inhibitory concentration of the stem bark extract of the plant ranged between 35 and 50 mg/ml while the minimum bactericidal concentration ranged between 35 and 60 mg/ml. has explored the antimicrobial activities of the crude ethanolic extracts of five plants against multidrug-resistant (MDR) strains of *Escherichia coli*, *Klebsiella pneumonia*, and *Candida albicans* and ATCC strains of *Streptococcus mutants*, and different strains of microorganism (Saini M.L., 2008 and Mahesh B, 2008)

3.6. Antimalarial activity

The root extracts of *A. nilotica* were active against *Plasmodium berghei* and *Plasmodium falciparum* in mice. *In vitro* Antimalarial activity against CQ sensitive (3D7) and CQ-resistant (Dd2 and INDO) strains of *P. falciparum* in culture using the fluorescence-based SYBR (Ali A.J., 2010). *A. nilotica* was reported with significant activity and IC₅₀ was found

at 13µg/mL *Acacia nilotica* has the highest Antiplasmodial activity *in vitro* against *Plasmodium falciparum* 3D (chloroquine-sensitive) and Dd2.(Chloroquine-resistant and pyrimethamine sensitive) organisms (Khan Rosina,2009)

3.7. Anti-Diarrhoeal Activity

Acacia nilotica has been reported to be very useful in treating diarrhea and cough in humans. The powdered bark of the plant with a little salt is used for treating acute diarrhea. Methanol and chloroform extract exhibit activity against *Escherichia coli*. (Chaubal R. et al 2006). It reduces the number of unformed feces and decreased the intestinal transit of charcoal.

3.8. Antispasmodial activity

The aqueous extract of seeds of *Acacia nilotica* shows spasmogenic activity on the isolated guinea-pig ileum. The mechanism behind it may be an increase in calcium influx that results in muscle spasms (Tahir A.E, et al, 1999)

3.9. Antihypertensive activity & Vasoconstriction activity

Methanol extract of *Acacia nilotica* pods causes a decrease in arterial blood pressure at a dose (3–30 mg/kg). It produces an inhibitory effect on force and rate of contraction in guinea-pig paired atria. Aqueous extract of *Acacia nilotica* possesses vasoconstriction by increasing calcium ion influx which is responsible for vasoconstriction action in guinea-pig ileum. The extract shows dose-dependent sustained contractile activity Eline (M.B. Ouedraogo, 2004)

Conclusion:

The various extracts of *Acacia nilotica* have nine different pharmacological activities, proved in most animal studies. Further in-vivo evaluation can help to develop the new drug dosage forms.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

References

1. Brenan J.P. Manual on taxonomy of *Acacia* species: present taxonomy of four species of *Acacia* (*A. albida*, *A. senegal*, *A. nilotica*, *A. tortilis*). *FAO, Rome*, 1983, 20-24.

2. Khan Rosina; Barira Islam; Mohd Akram; Shazi Shakil. Antimicrobial Activity of Five Herbal Extracts Against Multi Drug Resistant (MDR) Strains of Bacteria and Fungus of Clinical Origin. *Molecules*. 2009, 14(2), 586-597.
3. Chaubal R; Tambe A. Isolation of new straight chain compound from *Acacia nilotica*. *Ind. J. Chem.* 2006, 45(B), 1231-1233.
4. Bushra Farooq A; Roman Antioxidant activity of phenolic components present in barks of *Azadirachta indica*, *Terminalia arjuna*, *Acacia nilotica*, and *Eugenia jambolana* Lam. Trees, *Food Chemistry*, 2007, 104, 3, 1106-1114.
5. Saini M.L. Comparative Pharmacognostical and antimicrobial studies of *Acacia* species (Mimosaceae). *Journal of Medicinal Plants Research*, 2008, 2(12):378-386.
6. Khan R. Antimicrobial Activity of Five Herbal Extracts Against Multi Drug Resistant (MDR) Strains of Bacteria and Fungus of Clinical Origin. *Molecules*, 2009, 14(2): 586-597.
7. Mashram N Antimicrobial activity of methanol extracts of medicinal plants against bacterial species. *Int. Res. J*, 2009, I (3 &4), 147150.
8. Mahesh B; Satish S. Antimicrobial Activity of Some Important Medicinal Plant against Plant and Human Pathogens. *World Journal of Agricultural Sciences*, 2008, 4 (S), 839843.
9. Ali A.J; Akanya H.O.; Dauda B.E.N. Polygalloyltannin isolated from the roots of *Acacia nilotica* Del. (Leguminosae) is effective against *Plasmodium berghei* in mice. *J. Med. Plants. Res*, 2010, 4(12), 1169-1175.
10. Tahir A.E; Satti G.M.H; Khalid S.A . Antiplasmodial Activity of Selected Sudanese Medicinal Plants with Emphasis on *Acacia nilotica*. *Phytother. Res*, 199, 13, 474-478.
11. Singh B.N; Singh B.R.; Singh R.L. Antioxidant and anti-quorum sensing activities of green pod of *Acacia nilotica* L. *Food. Chem. Toxicol*, 2009, 47(4), 778-786.
12. Bachayaa H.A; Zafar I; Nisar K.M. Anthelmintic activity of *Ziziphus nummularia* (bark) and *Acacia nilotica* (fruit) against *Trichostrongylid* nematodes of sheep, *J. Ethnopharmacol*, 2009, 123, 325-329.
13. Eline M.B; Ouedraogo Z.L; Heide D Effect of aqueous extract of *Acacia nilotica* ssp *adansonii* on milk production and prolactin release in the rat, *J. Endocrin*, 2004, 182, 257-266.

14. Anonymous. The Wealth of India, Raw materials. Vol 9. New Delhi CSIR, 2003; 37-41.
15. Antaki AD. Tazkiradu Ulil Albab. Ed 1, Beirut: *Dar Alkutub Al Ilmiyah*, 1998; 1-3.
16. Dymock W; Warden CJH; Hooper DA History of the principal drugs of vegetable origin. Vol 1. *Pharmacographica Indica. New Delhi Shrishti book distributors*, 2005, 556.
17. Fosberg FR; Dasanayakke MD. Flora of Ceylon. *New Delhi: Oxford and IBH publishing company*, 1983, 532.
18. Ghani MN. Khazainul Advia. New Delhi Idarae Kitabul Shifa YNM, 1980, Vol 1,488,254.
19. Gupta; R.K. Resource survey of gummiferous acacias in western Rajasthan. *Tropical Ecology*, 1970, 11,148-161.
20. Hakim A; Bustanul Mufredat. New Delhi. *Idarae Kitabul Shifa*, 2002,Part I, 64: 65.
21. Jansen, P.C.M; Cardon D. Plant resources of tropical Africa.'Dyes and Tannins'. *Proto Foundation / Backhuys Publishers / CTA Wageningen, Netherlands*, 2005, 3.
22. Orwa, C; Mutua, A; Kindt R;Jamnadass; R.;Anthony; S.Agro. forestry Database: a tree reference and select ion guide version 4.0 (<http://www.worldagroforestry.org/sites/treed/treedatabases.asp>),2009.
23. Rastogi RP; Mehrotra BN. Compendium of Indian Medicinal Plants. *New Delhi NISC*, 2001, Vol 3, 4-7.
24. Rushd I; Kitabul Kulliyat. *New Delhi: CRUM*, 1987, Vol 2, 56, 114-117, 210, 222, 259.
25. Said H.M. Hamdard Pharmacopeia of Eastern Medicine. New Delhi,S ri Satguru publications, *Journal of Pharmacognosy and Phytochemistry*, 1997, Ed 2,353.
26. Shetty, K.A.B., Social forestry in Tamil Nadu. *Indian Farming*, 1977, 26, 82.
27. Sina I; Al Qanoon Fil Tib. (Eng translation). Book II. New Delhi: *Jamia Hamdard*, 1993, 70, 71, 433.
28. Basu, S.; Haldar, N.; Bhattacharya, S.; Biswas, S.; Biswas, M. Hepatoprotective activity of *Litchi chinensis* leaves against paracetamol-induced liver damage in rats. *Amer. Eurasian J. Scientific Res.* 2012, 7, 77-81.
29. Besra, S.E.; Sharma, R.M.; Gomes A. Anti-inflammatory effect of petroleum ether extract of leaves of *Litchi chinensis* Gaertn. (Sapindaceae). *J. Ethnopharmacol.* 1996, 54, 1-6.

30. Bhat, R.S.; Al-daihan, S. Antimicrobial activity of *Litchi chinensis* and *Nephelium lappaceum* aqueous seed extracts against some pathogenic bacterial strains. *J. King Saud Univ.-Sci.* 2014, 26, 79-82.
31. Bhoopat, L., Srichairatanakool, S., Kanjanapothi, D., Taesotikul, T., Thananchai, H., Bhoopat, T. Hepatoprotective effects of lychee (*Litchi chinensis* Sonn.): A combination of antioxidant and anti-apoptotic activities. *J. Ethnopharmacol.* 2011, 136, 55-66
32. Castellain, R.C.L.; Gesser, M.; Tonini, F.; Schulte, R, V.; Demessiano, K, Z.; Wolff, F.R.; Delle- Monache, F.; Netz, D.J.A.; Cechinel-Filho, V.; de Freitas, R.A.; de Souza, M.M.; Meyre- Silva, C. Chemical composition, antioxidant and antinociceptive properties of *Litchi chinensis* leaves. *J. Pharm. Pharmacol.* 2014, 66, 1796-1807.
33. Cooray HC, Janvilisri T, van Veen HW, Hladky SB, Barrand MA. Interaction of the breast cancer resistance protein with plant polyphenols. *Biochem. Biophys. Res. Commun.* 2004. 317:269–75.
34. Guo, C.J.; Yang, J.J.; Wei, J.Y.; Li, Y.F.; Xu, J.; Jiang, Y.G. Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutri. Res.* 2003, 23(12), 1719-1726
35. Guo, J.W.; Liao, H.F.; Pan, J.Q.; Ye, B.B.; Jian, X.B.; Wei, D.L.; Dai, L.Y. Effects of saponin of Litchi on decreasing blood glucose and controlling blood lipid in hyperlipidemia-fatty liver rats fed by high-sugar-fat. *Zhonguo Linchuang Yaolixueyu Zhiliao* 2004, 9, 1403-1407.
36. Huang, F.; Guo, Y.; Zhang, R.; Yi, Y.; Deng, Y.; Su, D.; Zhang, M. Effects of drying methods on physicochemical and immunomodulatory properties of polysaccharide-protein complexes from litchi pulp. *Molecules* 2014, 19, 12760-12776.
37. Jiang, W.F.; Chen, J.Z.; Zhang, J.; Peng, J. Inhibitory effects of total saponins extracted from the semen lychee against Hepatitis B virus in-vitro. *J. Four Military Medi. Univ.* 2008, 29(2), 100-103.
38. Jing, Y.; Huang, L.; Lv, W., Tong, H.; Song, L.; Hu, X.; Yu, R. Structural characterization of a novel polysaccharide from pulp tissues of *Litchi chinensis* and its immunomodulatory activity. *J. Agri. Food Chem.* 2014, 62, 902-911.
39. Kanlayavattanakul, M., Ospondpant, D., Ruktanonchai, U., Lourith, N., 2012. Biological activity assessment and phenolic compounds characterization from the

- fruit pericarp of *Litchi chinensis* for cosmetic applications. *Pharmaceutical Biology* 50, 1384-1390.
40. Lin, Y.; Chang, J.; Cheng, S.; Wang, C.; Jhan, Y.; Lo, I.; Hsu, Y.; Liaw, C.; Hwang, C.; Chou, C. New bioactive chromanes from *Litchi chinensis*. *J. Agri. Food Chem* 2015, 63, 2472-2478.
 41. Liu, L.; Xie, B.; Cao, S.; Yang, E.; Xu, X.; Guo, S. A-type procyanidins from *Litchi chinensis* pericarp with antioxidant activity. *Food Chemistry* 2007, 105, 1446-1451.
 42. Luximon-Ramma, A., Bahorun, T. and Crozier, A. 2003. Antioxidant actions and phenolic and vitamin C contents of common Mauritian exotic fruits. *Journal of the Science of Food and Agriculture* 83:496-502. Marchand LL. Cancer preventive effects of flavonoids — a review. *Biomed. Pharmacother.* 2002, 56, 296–301.
 43. Menzel, C.M. *The Lychee Crop in Asia and the Pacific*. Food and agriculture organization of the United Nations regional office for Asia and the Pacific Bangkok, Thailand, 2002, pp. 8-13.
 44. Menzel, C.M.; Waite, G.K. *Litchi and longan: Botany, Production, and Uses*. CAB e BOOK, 2005, 26.
 45. Nacif, S.R.; Paoli, A.A.S.; Salomão L.C.C. Morphological and anatomical development of the litchi fruit (*Litchi chinensis* Sonn. cv. Brewster). *Fruits* 2001, 56, 225-233.
 46. Nimmanpipug, P.; Lee, V.S.; Wolschann, P.; Hannongbua, S. *Litchi chinensis*-derived terpenoid as anti-HIV-1 protease agent: structural design from molecular dynamics simulations. *Molecular Simulation* 2009, 35, 673-680.
 47. Pan, J.Q., Liu, H.C., Liu, G.N., Hu, Y.L., Chen, L.X. and Qiu, Z.Q. 1999. A study in blood sugar reducing, blood lipid controlling and antioxidant activities in litchi seed. *Guangdong Pharmaceut. J.* 9(1):47-50.
 48. Pan, J.Q.; Guo, J.W.; Han, C.; Liu, H.C. Survey of pharmacological experimental studies on Litchi seeds. *Zhonguo Xinyao Zazhi* 2000, 9, 14-16.
 49. Qi, S.; Huang, H.; Huang, J.; Wang, Q.; Wei, Q. Lychee (*Litchi chinensis* Sonn.) seed water extract as potential antioxidant and anti-obese natural additive in meat products. *Food Control* 2015, 50, 195-201.
 50. Queiroz, E.d.; de Abreu, C.M.P.; Rocha, D.A.; Simao, A.A.; Bastos, V.A.A.; Botelho, L.N.S.; Braga, M.A. Anti-nutritional compounds in fresh and dried lychee fraction (*Litchi chinensis* Sonn.). *African J. Agric. Res.* 2015, 10, 499-504

51. Ren, S.; Xu, D.; Pan, Z.; Gao, Y.; Jiang, Z.; Gao, Q. Two flavanone compounds from litchi (*Litchi chinensis* Sonn.) seeds, one previously unreported, and appraisal of their α -glucosidase inhibitory activities. *Food Chem.* 2011, 127, 1760-1763.
52. Rodgers, E.H.; Grant, M.H. The effect of the flavonoids, quercetin, myricetin and epicatechin on the growth and enzyme activities of MCF7 human breast cells. *Chem.-Biol. Interact.* 1998, 116, 213–218.
53. Shen, R.; Duo-Duo, X.; Yang, G.; Yu-Ting, M.; Qi-Pin, G. Flavonoids from Litchi (*Litchi chinensis* Sonn.) seeds and their inhibitory activities on α -glucosidase. *Chemical Res. in Chinese Universities* 2013, 29, 682-685.
54. Su, D.; Ti, H.; Zhang, R.; Zhang, M.; Wei, Z.; Deng, Y.; Guo, J. Structural elucidation and cellular antioxidant activity evaluation of major antioxidant phenolics in lychee pulp. *Food Chem.* 2014, 158, 385-391.
55. Sung, Y.Y.; Yang, W.K.; Kim, H.K. Antiplatelet, anticoagulant and fibrinolytic effects of *Litchi chinensis* Sonn. extract. *Mol. Med. Rep.* 2012, 5, 721-724
56. Lchinose T, Musyoka TM, Watanabe K, Kobayashi N. Evaluation of antiviral activity of Oligonol, an extract of *Litchi chinensis*, against betanodavirus. *Drug discoveries & therapeutics.* 2013, 7(6), 254-260.
57. Tu, P.; Luo, Q.; Zheng, J. Studies of chemical constituents in seed of *Litchi chinensis*. *Zhongcaoyao* 2002, 33, 300-303.
58. Wang, L.; Lou, G.; Ma, Z.; Liu, X. Chemical constituents with antioxidant activities from litchi (*Litchi chinensis* Sonn.) seeds. *Food Chem.* 2011, 126, 1081-1087.
59. Wang, X., Yuan, S., Wang, J., Lin, P., Liu, G., Lu, Y., Zhang, J., Wang, W., Wei, Y., 2006. Anticancer activity of litchi fruit pericarp extract against human breast cancer in vitro and in vivo. *Toxicol. Appl. Pharmacol.* 215, 168-178.
60. Wen, L.; Wu, D.; Jiang, Y.; Prasad, K.N.; Lin, S.; Jiang, G.; He, J.; Zhao, M.; Luo, W.; Yang, B. Identification of flavonoids in litchi (*Litchi chinensis* Sonn.) leaf and evaluation of anticancer activities. *J. Functional Foods* 2014, 6, 555-563.
61. Winter, C.A., Risley, E.A. and Nuss, G.W. (1962) Carrageenan-induced oedema in hind paw of the rat as an assay for anti inflammatory drugs. *Proc. Soc. Experim. Biol. Med.* 111, 544 - 547.
62. Wu, Y.S.; Chiu, C.; Yang, D.; Lin, Y.; Jung-Kai, T.; Chen, Y. Inhibitory effects of litchi (*Litchi chinensis* Sonn.) flower-water extracts on lipase activity and diet-induced obesity. *J. Functional Foods* 2013, 5, 923-929.

63. Xu, X.; Xie, H.; Wang, Y.; Wei, X. A-Type Proanthocyanidins from Lychee seeds and their antioxidant and antiviral activities. *J. Agric. Food Chem.* 2010, 58, 11667-11672.
 64. Yamanishi, R.; Yoshigai, E.; Okuyama, T.; Mori, M.; Murase, H.; Machida, T.; Okumura, T.; Nishizawa M. The anti-inflammatory effects of flavanol-rich lychee fruit extract in rat hepatocytes. *PLoS ONE* 2014, 9, e93818.
 65. Yang, D.J.; Chang, Y.Y.; Hsu, C.L.; Liu, C.W.; Wang, Y.; Chen, Y.C. Protective effect of a litchi (*Litchi chinensis* Sonn.)-flower-water-extract on cardiovascular health in a high- fat/cholesterol-dietary hamsters. *Food Chem.* 2010, 119, 1457-1464.
 66. Zhang, J.; Zhang, C. Research progress on the antineoplastic pharmacological effects and mechanisms of Litchi seeds. *Chinese Medicine* 2015, 6, 20-26.
 67. Zhao, M.; Yang, B.; Wang, J.; Liu, Y.; Yu, L.; Jiang, Y. Immunomodulatory and anticancer activities of flavonoids extracted from litchi (*Litchi chinensis* Sonn.) pericarp. *Intern. Immunopharmacol.* 2007, 7, 162-166.
 68. Zheng, M.S.; Zheng, Y.F. Experimental studies on the inhibition effects of 1000 Chinese medicinal herbs on the surface antigen of Hepatitis B virus. *J. Trad. Chinese Med.* 1992, 12(3):193.
-