

Original Research Article

Estimates of genetic variability for seed yield and its component characters in green gram (*Vigna radiata*, Fabaceae)

Abstract

Sixteen green gram (*Vigna radiata*, Fabaceae) genotypes were evaluated under **randomized block design (RBD)** with three replications for genetic variability for eight characters. The analysis of variance revealed significant differences among genotypes for all the characters. Analysis of variance indicated that experimental material possesses a good amount of variability for all the characters under study. A high magnitude of genotypic coefficient of variance was observed for seed yield per plot, seed yield per hectare, plant height, days to flowering, and test weight while moderate GCV was recorded for the number of seeds per pod, pod length, and days to maturity suggesting the possibility of their improvement by selection. High heritability coupled with high genetic gain as a percent of mean was observed for test weight, seed yield per hectare, seed yield per plot, plant height, and days to flowering while high heritability and moderate genetic gain were recorded for days to maturity, pod length and the number of seeds per pod suggesting that these characters were controlled by additive gene action hence, selection may be effective. High heritability coupled with high genetic gain and high GCV has been exhibited by the characters viz., seed yield per hectare, seed yield per plot, plant height, test weight, and days to flowering. There is substantial scope for improvement of these characters. Hence emphasis should be given to improving these characteristics in a future breeding program to enhance the seed yield of green gram.

Keywords: Green gram, Genetic variability, GCV, heritability, seed yield

Introduction

“Green gram (*Vigna radiata*, Fabaceae) is one of the important pulse crops in the arid region because of its short growth duration, adaptation to low water requirements, and low soil fertility. It is favored for consumption due to its easy digestibility and low production of flatulence” (Shil and Bandopadhyaya, 2007; Dadepeer *et al.*, 2009). “Pulses are extensively grown in tropical regions of the world

as a major protein-rich crop bringing considerable improvement to the human diet. The average protein content in the seeds is around 24 percent. The protein is comparatively rich in the amino acid lysine but predominantly deficient in cereal grains” (Baskaran *et al.*, 2009; Dhanajay *et al.*,2009). Presently, “the yield of green gram is well below the optimum level compared to other pulses. The average yield of green gram is very low not only in India (425 kg/ha) but in entire tropical and sub-tropical Asia. India is the largest producer of green gram in the world and accounts for 65 percent of the acreage and 54 percent of production” (Kumar *et al.*,2005 and Pratap *et al.*, 2012).

To develop high-yielding varieties a systematic breeding approach has to be adopted. Assessment of genetic variability is the first step in any breeding program. The greater the diversity in the material better are the chances of improvement, provided the heritability is high and genetic gain is more. Further, the selection is more effective when it is practiced simultaneously for the characters who have desired nature of association with the characters of ultimate interest.

Materials and Methods

Sixteen genotypes of green gram were raised in RBD with three replications during *Kharif*, 2019 at the research farm of Agricultural Research Station, Fatehpur-Shekhawati, Sikar (Rajasthan) under rainfed conduction. These genotypes of green gram were obtained from the All India Coordinated Research Project on MULLaRP, RARI, Durgapur (Jaipur). Each genotype was given in a four-row plot of 4 m in length with a spacing of 30 cm between rows and 10 cm between plants. Ten plants were selected at random from each plot and data were recorded on eight characters viz., plant height, pod length, number of seeds per pod, test weight of seeds, seed yield per plot, and seed yield per ha., whereas for days to 50% flowering and days to maturity data were recorded on a whole plot basis. The data so obtained were subjected to analysis of variance and estimation of different variability parameters.

Statistical methods

- (i) Analysis of variance (Panse and Sukhatme, 1985).
- (ii) Variability parameters.
 - a) Phenotypic Coefficient of variability (PCV) (Burton, 1952).
 - b) Genotypic Coefficient of variability (GCV) (Burton, 1952).
 - c) Heritability (Burton and dvane, 1953).

- d) Genetic gain (Johnson *et al.*, 1955).

Results and Discussion

The analysis of variance revealed highly significant differences among the treatments for all the characters suggesting the presence of a high degree of genetic variability in the experimental material (Table-1). In general, the estimates of PCV for all the characters were higher than the GCV indicating the environmental influence over the expression (Table-2). This finding was in accordance with Anand *et al.* (2015), Hemavathy *et al.* (2015), Mehandi *et al.* (2013), and Raturi *et al.* (2015) in green gram. Characters viz., seed yield per plot (23.21%) had the highest genotypic coefficient of variation (GCV) followed by seed yield per hectare (23.04%), plant height (8.61%), days to flowering (5.84%), and test weight of seeds (5.01%). Similar results were reported by Gadakh *et al.* (2013), Raturi *et al.* (2015), Garg *et al.* (2017), and Muthuswamy *et al.* (2019) in green gram. Moderate estimates of GCV were observed for no. of seeds per pod, pod length, and days to maturity indicating the limited scope of selection for these characters. Similar results were reported by Mehandi *et al.* (2013) and Makeen *et al.* (2007) in green grams.

The selection of characters based on heritability and genetic gain as a percent of the mean is of great importance (Mehandi *et al.*, 2013; Narasimhulu *et al.*, 2013a). The highest heritability estimate was recorded for test weight (90.90%) followed by seed yield per hectare (89.80%), seed yield per plot (89.70%), plant height (87.90%), days to flowering (87.20%), and days to maturity (60.10%) suggesting these characters are governed by the additive genetic effect to great extent and improvement of these characters would be effective through phenotypic selection. Similar findings were also observed by Makeen *et al.* (2007), Mehandi *et al.* (2013), Sowmini and Jayamani (2013), Prasanna *et al.* (2013), Rashid and Biswas (2015), Hemavathy *et al.* (2015) and Garg *et al.* (2017). The high heritability with high genetic gain for test weight, seed yield per hectare, seed yield per plot, plant height, and days to flowering was due to additive gene effects. The characters, pod length, and no. of seeds per pod exhibited moderate heritability indicating non-additive gene action and environmental influence on the expression of characters, where selection is ineffective and could be better exploited through recombination breeding. Similar results were reported by Mehandi *et al.* (2013) for pod length and Makeen *et al.* (2007) for the number of seeds per pod in green gram (Figure-1).

Genetic gain percent of mean was observed highest for seed yield per plot (45.29%) followed by seed yield per hectare (45.00%), plant height (16.64%), days to flowering (11.23%), test weight (9.85%) and no. of seeds per plant (5.38%). Comparatively moderate values of genetic gain were recorded for days to maturity and pod length displayed moderate genetic gain as a percent of mean suggesting that all these characters were governed by additive genes and selection will be rewarding for improvement of such characters. Similar results were reported by Mishra and Yadav (1992), Dashora and Nagda (2002), Rahim *et al.* (2010), and Srivastava and Singh (2012).

The characters having a higher quantum of genetic variability along with high heritability and greater genetic gain can be easily improved through selection (Johnson *et al.*, 1955). Panse (1957) also suggested that the characters which have high heritability with higher genetic gain are expected to have a greater proportion of genetic effects. High heritability coupled with high genetic gain and high GCV has been exhibited by the characters *viz.*, seed yield per hectare, seed yield per plot, plant height, test weight, and days to flowering indicating control of additive gene action for these characters. Similar, results were reported by Jain and Ramgiry (2000), Singh *et al.* (2000), and Degefa *et al.* (2014).

Conclusion

High GCV coupled with high heritability and genetic gain of a character provides a good selection advantage. In the present study, the characters, seed yield per hectare, seed yield per plot, plant height, test weight, and days to flowering indicate that the variation in the characters was most likely due to additive gene effects, hence, simple directional selection may be effective to improve these characters in green gram.

References

- Anand, G., Anandhi, K. and Paulpandi, V.K. 2015. Genetic variability, correlation and path analysis for yield and yield components in F₆ families of Green gram (*Vigna radiata* (L.) Wilczek) under rainfed condition. *Electronic J. of Plant Breeding*, **7**(2): 434-437.

- Baskaran, L., Sundararmoorthy, P., Chidambaram, A.L.A. and Ganesh, K.S. 2009. Growth and physiological activity of green gram (*Vigna radiata* (L.) Wilczek) under effluent stress. *Bot. Res. Int.*, 2: 107-114.
- Burton, G.W. 1952. Quantitative inheritance in grasses. *Proc. Sixth Int. Grassld Congr.* 1: 277-283.
- Burton, G.W. and Devane, E.M., 1953. Estimation of heritability in tall fescue (*Festuca arundinaceae*) from replicated clonal material. *Agron. J.* 45: 478-481.
- Dashora, A. and Nagda, A.K. 2002. Genetic variability and character association in Spanish bunch groundnut. *Research on Crops* 3: 416-420.
- Dadepeer, Peerajade, Ravi Kumar, R. L. and Salimath, P.M. 2009. Genetic variability and character association in local green gram genotypes. *Environment and Ecology.* 27(1): 165-169.
- Degefa, I., Petros, Y. and Andargie, M. 2014. Genetic variability, heritability and genetic advance in Mungbean (*Vigna radiata* L. Wilczek) accessions. *Plant Science Today*, 1(2): 94-98
- Dhananjay, Ramakant, Singh, B. N. and Singh, G. 2009. Studies on genetic variability, correlations and path coefficients analysis in mung bean. *Crop Res. Hisar.* 38(1/3): 176-178.
- Gadakh, S.S., Dethé, A. M., and Kathale, M. N. 2013. Genetic variability, correlations and path analysis studies on yield and its components in mungbean (*Vigna radiata* (L.) Wilczek) *Bioinfolet*, 10 (2a), 441-447.
- Garg, G. K., Verma, P. K. and Kesh, H. 2017. Genetic Variability, Correlation and Path Analysis in Greengram [*Vigna radiata* (L.) Wilczek]. *Int. J. Curr. Microbiol. App. Sci.*, 6(11): 2166-2173.
- Hemavathy, A. T., Shunmugavalli, N. and Anand, G. 2015. Genetic variability, correlation and path co-efficient studies on yield and its components in mungbean [*Vigna radiata* (L.) Wilczek]. *Legume Res.*, 38(4): 442-446.
- Jain, P.K. and Ramgiry, S.R. 2000. Genetic variability of metric traits in Indian germplasm of green gram. *Adv. in Plant Sciences.* 13: 127-131.
- Johnson, H. W., Robinson, H. F. and Comstock, R. E. 1955. Estimation of genotypic and environmental variability in soybean. *Agron. J.*, 47: 314-318.
- Kumar, U., S.P. Singh and Vikas 2005. Variability and character association in mungbean. (*Vigna radiata* (L.) Wilczek). *New Agriculturist*, 16 (1, 2): 23-28.

- Makeen, K., Abraham, G., Jan, A. and Singh, A. K. 2007. Genetic variability and correlations studies on yield and its components in Greengram [*Vigna radiata* (L.) Wilczek]. *Journal of Agronomy*, **6**(1): 216-218.
- Mehandi, S., Singh, C. M. and Kushwaha, V. K. 2013. Estimates of genetic variability and heritability for yield and yield component traits in Greengram [*Vigna radiata* (L.) Wilczek]. *The Bioscan*, **8**(4): 1481-1484.
- Mishra, L.K. and Yadav, R.K. 1992. Genetic variability and correlation studies in summer groundnut. *Advances in Plant Science* **5** : 106-110.
- Muthuswamy, A., Jamunarani, M. and Ramakrishnan, P. 2019. Genetic Variability, Character Association and Path Analysis Studies in Green Gram (*Vigna radiata* (L.) Wilczek). *Int.J.Curr.Microbiol.App.Sci.*, **8**(4): 1136-1146.
- Narasimhulu, R., Naidu, N.V. Priya, M.S. Rajarajeswari, V. and Reddy, K.H.P. 2013a. Genetic variability and association studies for yield attributes in mungbean (*Vigna radiata* (L.) Wilczek). *Indian J. Plant Sci.*, **2**(3): 82-86.
- Panse, V.G. 1957. Genetics of qualitative characters in relation to plant breeding. *Indian Journal of Genetics* **17**: 318-328.
- Panse, V.G. and Sukhatme, P.V. 1985. Statistical methods for research workers, ICAR, New Delhi.
- Prasanna, L. B., Rao, P.J.M., Murthy, K.G.K. and Prakash, K. 2013. Genetic variability, correlation and path coefficient analysis in mungbean. *Enviro. and Ecolo.*, **31**(4): 1782- 1788.
- Pratap, A., Gupta., D.S. and Rajan., N. 2012. Breeding Indian Field Crops. *Agro bios Publishers*, New Delhi, India. p 208-227.
- Rahim, M. A., Mia, A. A., Mahmud, F., Zeba, N. and Afrin, K. S. 2010. Genetic variability, character association and genetic divergence in mungbean (*Vigna radiata* (L.) Wilczek). *Plant Omics*. **3**(1): 1-6.
- Rashid, K. M. and Biswas, M. 2015. Genetic Variability, Correlation and Path Analysis in Mungbean (*Vigna radiata* L). *J. of Environmental Science and Natural Resources*, **7**(1): 131-138.
- Raturi, A., Singh, S. K., Sharma, V. and Pathak, R. 2015. Genetic variability, heritability, genetic advance and path analysis in Greengram [*Vigna radiata* (L.) Wilczek]. *Legume Res.*, **38** (2): 157-163.
- Shil, S. and Bandopadhyaya, P.K. 2007. Retaining seed vigor and viability of mungbean by dry dressing treatments. *J. Food Legumes.*, **20**: 173-175.

- Singh, J., Parmar, R.P., Yadav, H.S. and Singh., J. 2000. Assessment of genetic variability and selection parameters in early generation of green gram. *Adv. in Plant Sciences*. 13: 227-232.
- Sowmini, K. and Jayamani, P. 2013. Genetic variability studies for yield and its component traits in RIL population of blackgram (*Vigna mungo* (L.) Hepper). *Electronic J. of Plant Breeding*, 4 (1): 1050-1055.
- Srivastava, R.L and Singh, G. 2012. Genetic variability, correlation and path analysis in mungbean (*Vigna radiata* (L.) Wilczek). *Indian J. Life Sci.*, 2(1): 61-65.

UNDER PEER REVIEW

Table-1: Analysis of variance showing values of mean squares for different characters in green gram (*Vigna radiata*, Fabaceae).

Characters	Source o variance		
	Replications [2]	Treatments [15]	Error [30]
Days to flowering	5.395**	17.016*	0.795
Days to maturity	6.812*	7.420*	1.345
Plant height	4.036	41.335**	1.808
Pod length	3.203**	0.216*	0.079
No. of seeds per pod	2.270*	1.516*	0.670
Seed yield per plot	39.062	11758.020**	432.395
Test weight	0.071	8.099**	0.261
Seed yield per ha.	1181.73	89205.532**	3239.501

[] = df

*,** significant at 5% level and 1% level, respectively.

Table-2: Variability parameters for different characters in green gram(*Vigna radiata*, Fabaceae).

Characters	Coefficient of variation		Heritability (broad sense) (%)	Genetic gain (%)	Genetic gain % of mean
	GCV (%)	PCV (%)			
Days to flowering	5.84	6.25	87.20	4.47	11.23
Days to maturity	2.32	3.00	60.10	2.27	3.71
Plant height	8.61	9.18	87.90	7.01	16.64
Pod length	2.77	4.56	36.80	0.26	3.46
No. of seeds per pod	4.80	8.84	29.60	0.59	5.38
Seed yield per plot	23.21	24.50	89.70	119.89	45.29
Test weight	5.01	5.26	90.90	3.17	9.85
Seed yield per ha.	23.04	24.31	89.80	330.53	45.00

Figure-1: Genetic parameters [PCV, GAM, GCV and h^2 (broad sense)] for different characters in green gram(*Vigna radiata*, Fabaceae).

