

Original Research Article

ASSESSMENT OF GENETIC DIVERSITY FOR COTTON LEAF CURL DISEASE (CLCuD) AND QUALITATIVE TRAITS AMONG ELITE COTTON CULTIVARS

ABSTRACT

Cotton is a major fibrous cash crop and is cultivated in more than 80 countries because global textile industry depends on it. In Pakistan, the major factor reducing the cotton production is cotton leaf curl disease which is caused by a begomovirus, cotton leaf curl virus vectored through a whitefly, *Bemisia tabaci*. This disease shows a wide variety of symptoms including vein thickening, stunted plant growth, and cup-shaped outgrowth known as Enation. This study was conducted to find out the genetic diversity among 50 cotton varieties by using 10 SSR primers. DNA extraction was done by using CTAB method which was followed by DNA quantification through agarose gel to confirm its quantity. PCR and PAGE were performed to check the quality of DNA. Statistical analysis was performed by using Power marker v. 3.25. Primer NAU 2083 showed the maximum genetic diversity and PIC value of 0.8736 and 0.8621 respectively. The unweighted pair-group method analysis (UPGMA) generated the phylogenetic tree to divide the varieties into two clusters which represented the greater genetic diversity in three cotton varieties. These varieties can be utilized in future breeding programmes for cotton improvement.

Keywords: Cotton, CLCuD, Elite, Cultivars, Diversity

INTRODUCTION

Cotton (*Gossypium hirsutum* L), a major fibrous cash crop and performs a crucial function in oil and forage industries [1]. It is also named as "White Gold" as its consumptions as lint and its end-products and foreign exchange earnings [2]. It is grown widely across the globe because of its economic status. In many cotton producing countries, cotton diligence is of great worth as it contributes to their GDPs [3]. The upland cotton is accountable for 90% of total world's cotton production [4]. It has yearly production of over 20 million tons and is produced in >80 countries because worldwide textile diligence depends on both natural and artificial sources of fiber production [2]. In agronomic countries like Pakistan, cotton is a significant crop and it subsidizes 0.8% in Gross Domestic Product (GDP) and 4.5% in the agriculture division in Pakistan [4]. Pakistan is the 5th prevalent producer and the 3rd prevalent exporter of cotton throughout the entire globe [5]. Economy of Pakistan chiefly

Comment [IG1]: I think it should be reformulated with something like: This study was conducted to find out the genetic diversity for cotton leaf curl disease among 50 cotton varieties by using 10 SSR primers.

depends on cotton fabrication because it is not only the basis of fiber but also offers vegetable oil and low-quality oil for soup diligence. Cotton crop is regarded as the king of fiber and cotton plant delivers 70-80% of natural fiber [2]. Cotton fibers are used for the production of nano-fibrillate cellulose materials via nano-technological appliances. Woven fabrics, foodstuffs, cosmetic and soap are also manufactured from cotton fiber [3]. Edible oil and seed cake are also acquired from cotton seeds [4]. Due to several biotic and abiotic causes, cotton production is reduced [6]. Cotton crop, the cultivated shrub, is susceptible to several pathogens and pests out of which the most destructive aspect is cotton leaf curl disease (CLCuD) [7]. CLCuD is caused by multiple begomoviruses which are vectored through the whitefly, *Bemisia tabaci* (Zubair et al. 2021). Across Pakistan and northwestern India, cotton plants are destroyed by CLCuD causing drastic reduction in production [8]. The objective of study is to analyze the genetic diversity for Cotton Leaf Curl Disease among elite cotton cultivars.

MATERIALS AND METHODS

Leaf samples of fifty cotton (*G. hirsutum*) varieties were gathered from Cotton Research Station and Central Cotton Research Institute Multan in the months of September and October (Table 1). Susceptible and resistant plants were selected according to the infection scale [9]. For assessment of genetic diversity for CLCuD and phenotypic data for selected traits was also measured. Fresh and young leaves were selected for genomic DNA extraction via Cetyltrimethyl ammonium bromide (CTAB) method [1]. Extracted DNA was assessed to confirm its quantity through agarose gel electrophoresis. For the evaluation of genetic diversity, SSR primers were used to amplify 50 samples of cotton by using PCR. PAGE was executed to assess the PCR products.

Comment [IG2]: You should insert 1-2 more sentences related to genetic diversity and analyzes in this sense because this is the purpose of your study. And it is necessary to bring in attention other studies in the field with which you can compare yourself later in the discussions. Add other sources to be found in the final list of References. In fact, what is missing or what has not been sufficiently studied so that it is necessary to do additional research. Then come up with the purpose of your study.

Comment [IG3]: Separate the part describing the material collected and used from the part where you describe the genetic analyses.

Provide details about the material. In what conditions were the plants from which you collected leaves maintained. Did all 50 varieties have the same conditions or were they different?

Comment [IG4]: The work methodology would be much improved if you brought an image of the work. For example 1-2 pictures of the collection or leaves from the relevant varieties. Optional.

Comment [IG5]: What does PAGE mean? Explain the abbreviation mentioned for the first time in the manuscripts. Then you can only use it as an abbreviation. So that any reader can understand what you are referring to.

Table 1: Names of 50 Cotton varieties

Sr. No.	Variety Name	Sr. No.	Variety Name
1	AA 802	26	IUB 222
2	Barberton	27	LaOkr 541
3	BH 160	28	LB 391
4	Bt. CIM 602	29	MNH 329
5	CA 325 IRABLT	30	MNH 886
6	CEMB 33	31	NIAB 112
7	Chilala 76/2	32	NIAB 2009
8	CIM 1100	33	NIAB 2010
9	CIM 443	34	RH 112
10	CIM 446	35	S 14
11	CIM 448	36	S 32
12	CIM 473	37	Samaru 72
13	CIM 482	38	SI Okra1 23
14	CIM 499	39	Sitar 008
15	Cris 613	40	SLH 06
16	DP Acala 90	41	SLH 119
17	DPL NEW COTTON 33	42	SLH 12
18	FBS 30	43	SLH 13
19	FBS 37	44	SLH 317
20	FH 113	45	SLS 90/2
21	FH 87	46	SLS B7/175
22	FVH 53	47	TARZAN 1
23	GM 90	48	TARZAN 2
24	Gomal 105	49	VH 305
25	IR 3701	50	VH 363

Comment [IG6]: What is Sr.?

Comment [IG7]:

RESULTS

Comment [IG8]:

Analysis of Allele number

For the estimation of genetic diversity among 50 cotton accessions, 10 SSR primers were used. The total 57 loci have an average of 5.7 loci per primer. Number of alleles on every primer ranges from 1-8. The total number of alleles amplified by these SSR primers was 57. Primers NAU 980 and BNL 827 exhibited maximum number of polymorphic bands while the minimum numbers of polymorphic bands were exhibited by the primers NAU 2273 and NAU 2437. Primers and their allele number are denoted in the Table 2.

Calculating the Allele frequency, Genetic diversity and Polymorphism

Genetic diversity was evaluated by calculating the major allele frequency, polymorphism and allele number using power marker v 3.25. The mean value of genetic diversity was 0.4566 and in these SSR markers, the maximum genetic diversity was demonstrated by the primers NAU 2083 (0.8736) and NAU 2273 (0.6616) whereas the minimum genetic diversity was exhibited by the primers NAU 3414 (0.0000) and NAU 4042 (0.1512) represented by the Table 3.

Polymorphism of 10 SSR primers was finding out by calculating the PIC values. Mean PIC value of these primers was 0.4209. Maximum polymorphism was exhibited by the primers NAU 2083 and NAU 2273 having PIC values of 0.8621 and 0.5874 respectively whereas minimum polymorphism was exhibited by the primers NAU 3414 and NAU 4042 having PIC values of 0.0000 and 0.1471 respectively.

Comment [1G9]: Detail the abbreviation

Table 2 Primer, chromosome and allele number

Primers	Chromosome	Number of alleles
NAU 2083	17	7
NAU 2954	2	5
NAU 4042	4	7
NAU 3414	1	5
NAU 5046	5	6
NAU 2838	6	6
NAU 980	5	8

BNL 827	7	8
NAU 2273	3	2
NAU 2437	3	3
Total		57

Table 3 Major Allele Frequency, Gene diversity and Polymorphism

Marker	Major.Alele.Frqency	AlleleNo	Gene Diversity	PIC
NAU 2083	0.2400	17.0000	0.8736	0.8621
NAU 2954	0.8400	2.0000	0.2688	0.2327
NAU 4042	0.9200	4.0000	0.1512	0.1471
NAU 3414	1.0000	1.0000	0.0000	0.0000
NAU 5046	0.5000	5.0000	0.6344	0.5738
NAU 2838	0.6000	6.0000	0.5856	0.5452
NAU 980	0.7600	5.0000	0.3944	0.3619
BNL 827	0.6400	7.0000	0.5552	0.5258
NAU 2273	0.3800	3.0000	0.6616	0.5874
NAU 2437	0.7000	3.0000	0.4408	0.3728
Mean	0.6580	5.3000	0.4566	0.4209

Phylogenetic tree

The unweighted pair group method (UPGM) was used for the assessment of genetic diversity among 50 cotton genotypes by using SSR primers. Nei and Li's 1973 bootstrap neighbor joining method generated the phylogenetic tree. This phylogenetic tree classified into

two main clusters A and B. Cluster A comprises of 8 cotton genotypes and classified into two sub-clusters L and M. Sub-cluster L contains two sub-sub clusters L1 and L2. Sub-sub cluster L1 denoted three cotton genotypes including LaOkr 541, IR 3701, and NIAB 2010. Sub-sub cluster L2 comprises two cotton genotypes SLH 119 and SLH 13. Sub-cluster M further differentiates into two sub-sub clusters M1 and M2. M1 denoted the one genotype CEMB 33 whereas M2 designated the two cotton genotypes TARZAN 1 and FH 113.

Cluster B comprises of 42 cotton genotypes and distinguished further into two sub-clusters D and E. Sub-cluster D distributed into two sub-sub clusters D1 and D2. Sub-cluster D1 indicated the two cotton genotypes NIAB 112 and AA 802 while D2 exhibited the only one genotype NIAB 2009. Sub-cluster E distinguished into two sub-sub clusters E1 and E2. E1 further classified into E1-a comprising three cotton genotypes FBS 30 and VH 305 and SLH 12 in a distinct group. E1-b contains three genotypes Samaru 72 in one group and the other two genotypes Sitar 008 and MNH 886 in a distinct group. E2-a comprises only one genotype LB 391 however E2-b further distributed into two groups E2-ba and E2-bb. E2-ba comprising 7 genotypes further distinguished into two groups. One group comprises four genotypes Gomal 105, DPL NEW COTTON 33 and SI Okra1 23 and TARZAN 2 in a distinct group. The second group comprises three genotypes BH 160, FBS 37 and Cris 613. E2-bb classified into two groups E2-bb1 and E2-bb2. E2-bb1 further classified into two groups. One group comprising 9 genotypes include S 32, CIM 446, DP Acala 90, CIM 482, Bt. CIM 602, FH 87 and an isolated group of SLH 317, RH 112, and MNH 329. The second group comprises two genotypes VH 363 and SLSB 7/175. E2-bb2 further distributed into two sub-groups. One sub-group comprising five genotypes including CIM 448, CIM 443, CIM 1100 and Barberton and the other S 14 in a distinct group. Second group of E2-bb2 again discriminated into two sub-sub groups. One sub-sub group comprises only 1 genotype CA 325 IRABLT and the other sub-sub group further distinguished into two groups one comprising three genotypes FVH 53, CIM 449 and SLS 90/2 and the other sub-sub group also contains three genotypes GM 90, CIM 473 and Chilala 76/2.




Figure 1: Phylogenetic tree of 50 cotton varieties

Comment [IG10]: Does this diagram/figure belong to you or did you take it from another source? If it is from another source, please mention it below the figure and it would be good to have the usage agreement

DISCUSSION

Cotton is a substantial fibrous crop which donates in the textile diligence and also the basis of cotton seed. But cotton fabrication is in danger due to various biotic and abiotic reasons. Apart from the abiotic factors, Cotton leaf curl disease caused by *Begomovirus* and vectored through *B. tabaci* declines the cotton yield on the basis of severity of infection. Broad genetic diversity of genotypes can control such diseases [10]. Genetic diversity is defined as the total number of alleles with variable frequencies within a species. Genetic diversity arises due to alterations in the alleles of genetic makeup and is involved in the marker-aided

selection. SSR markers are used for the estimation of genetic diversity in cotton genotypes. These markers are co-dominant in nature and reveal accurate results [11]. Genetic variability for favorable traits is very little among elite cotton accessions. For the development of desirable cotton germplasm, SSR markers are utilized in the marker-aided selection [10].

In our study, the genetic diversity of 50 cotton genotypes was assessed by using ten SSR primers and total 57 alleles were amplified which ranged from 8 to 2 alleles with an average of 5.7 alleles per primer. Maximum numbers of polymorphic bands were characterized by the primers NAU 980 and BNL 827 whereas minimum numbers of polymorphic bands were demonstrated by the primers NAU 2273 and NAU 2437. In current study, 10 SSR primers were used to observe major allele frequency, genetic diversity and polymorphic information content (PIC) among 50 cotton genotypes. Our results revealed that maximum genetic diversity of 0.8736 and maximum PIC values of 0.8621 were presented by the primer NAU 2083. Estimation of genetic diversity by using SSR markers have been discussed in earlier studies [12].

Comment [IG11]: Cut and use ...Also

Comment [IG12]: What studies are we talking about? You should continue with a description of them. That is, what others have found and how do you see your results in comparison with those from previous studies.

Phylogenetic tree was generated by Nei and Li's 1973 bootstrap neighbor joining method to evaluate genetic diversity among 50 cotton varieties. This phylogenetic tree classified these 50 varieties into clusters and various sub-clusters to reveal the similarity index. Our results showed that DPL NEW COTTON 33 and Gomal 105 are closely linked and reside in the same group. Varieties CIM 446 and S 32 are also closely related and present in the same group. [13] reported the same results. Phylogenetic tree was constructed by UPGMA revealed genetic diversity among clusters. SSR primers are beneficial used for the genetic evaluation of cotton such as genetic diversity and marker-aided selection to choose desirable cotton varieties.

CONCLUSION

This study discovered the genetic diversity by using SSR markers among selected cotton cultivars. From this study, it is found out that great genetic diversity is recognized in some genotypes of cotton. It is observed that greater genetic diversity is examined in genotypes Chilala 76/2, Cris 613 and LaOkr 541 as they are present on maximum distance in clusters. This assessment of genetic diversity in different cotton genotypes provides the opportunity

Comment [IG13]: I think it is much more appropriate to use...analyzed... The term to ...discover...is much too deep and it seems that you have discovered this kind of analysis for the first time

to the breeders to select parental lines for the production of new cotton genotypes with superior yield and quality.

REFERENCES

1. Ahmed N, Amin I, Zaidi SS-E-A, Rahman SU, Farooq M, Fauquet CM, Mansoor S (2021) Circular DNA enrichment sequencing reveals the viral/satellites genetic diversity associated with the third epidemic of cotton leaf curl disease. *Biology Methods and Protocols* 6:bpab005.
2. Sarwar G, Nazir A, Rizwan M, Shahzadi E, Mahmood A (2021) Assessment of genetic diversity of cotton cultivars by using correlation and principal component analysis for CLCuD tolerance, yield and quality traits. *J. Agric. Res* 59:1-5.
3. Ozyigit II, Dogan I, Kaya Y, Bajrovic K, Gozukirmizi N (2022) Cotton biotechnology: An efficient gene transfer protocol via *Agrobacterium tumefaciens* for a greater transgenic recovery. *Journal of Natural Fibers* 1-15.
4. Sahar A, Zafar MM, Razzaq A, Manan A, Haroon M, Sajid S, Rehman A, Huijuan M, Ashraf M, Maozhi R (2021) Genetic variability for yield and fiber related traits in genetically modified cotton. *Journal of Cotton Research* 4:1-10.
5. Azam S, Rao AQ, Shad M, Ahmed M, Gul A, Latif A, Ali MA, Husnain T, Shahid AA (2021) Development of broad-spectrum and sustainable resistance in cotton against major insects through the combination of Bt and plant lectin genes. *Plant Cell Reports* 40:707-721. doi:10.1007/s00299-021-02669-6
6. Khan MaU, Shahid AA, Rao AQ, Shahid N, Latif A, Ud Din S, Husnain T (2015) Defense strategies of cotton against whitefly transmitted CLCuV and Begomoviruses. *Advancements in Life Sciences* 2:58-66.
7. Farooq T, Umar M, She X, Tang Y, He Z (2021) Molecular phylogenetics and evolutionary analysis of a highly recombinant begomovirus, Cotton leaf curl Multan virus, and associated satellites. *Virus evolution* 7:veab054.
8. Zubair M, Zaidi SS-E-A, Shakir S, Farooq M, Amin I, Scheffler JA, Scheffler BE, Mansoor S (2017) Multiple begomoviruses found associated with cotton leaf curl disease in Pakistan in early 1990 are back in cultivated cotton. *Scientific reports* 7:1-11.
9. K, Haidar S, Khan M, Ahmad M, Sarwar N, Murtaza M, Aslam M (2010) Evaluation of *Gossypium* species for resistance to cotton leaf curl Burewala virus. *Annals of applied biology* 157:135-147.
10. Abbas A, Iqbal MA, Rahman M-U, Paterson AH (2015) Estimating genetic diversity among selected cotton genotypes and the identification of DNA markers associated with resistance to cotton leaf curl disease. *Turkish Journal of Botany* 39:1033-1041.
11. Çelik S (2022) Genetic Diversity Analysis of Some Upland Cotton (*Gossypium hirsutum* L.) Genotypes Using SSR Markers. *Türk Doğa ve Fen Dergisi* 11:80-89.
12. Ali I, Khan NU, Gul S, Khan SU, Bibi Z, Aslam K, Shabir G, Haq HA, Khan SA, Hussain I (2019) Genetic diversity and population structure analysis in Upland cotton Germplasm. *International Journal of Agriculture and Biology* 22:669-676.
13. Rehman A, Iqbal M (2022) Genome-wide Allelic Variations and Population Structure Analysis of Cotton Genotypes using SSR Markers. *Pakistan Journal of Biochemistry and Biotechnology* 3:

Comment [IG14]: Please add the references from the Introduction that I suggested. Be careful to keep a unified description; sometimes you mention the full name of the journal, sometimes only the abbreviation. Reference 13 seems unfinished, so please add the necessary information.

UNDER PEER REVIEW