

Morphological Characterization and Adaptive Trail of two prominent T. Aman rice (*Oryza sativa* L.) Lines under different locations of Bangladesh

ABSTRACT

Two advanced lines: BR8693-8-4-2-1 and BR8693-17-6-2-2, along with BRRI dhan49 (S. Ck) and BRRI dhan33 (R. Ck) as checks were tested at farmers' field in eight locations such as West Byde (BRRI Gazipur), Khulna (Dumuria), Habiganj (Sadar), Chapainawabganj (Gomostapur), Rangpur (Sadar), Feni (Dagonbhuyan), Chattagram (Hathazari) and Barishal (Sadar) during the season of **Transplanting Aman** 2018. The trial **was** replicated three times in each location. The unit plot size was 17.94 m² (4.6m x 3.9m). Seeding emergence for eight locations varied from 11-17 July 2018. Seedling age varied from 25-30 days among the locations. Seedlings were transplanted at 20 cm x 15 cm spacing. NPKS and Zn fertilizers were applied at the rate of 0, 15, 50, 12 and 3.6 kg ha⁻¹, respectively. All fertilizers except urea were applied as basal whereas, urea was applied in 3 equal splits at 10, 25 and 40 **days after transplanting (DAT)** Other standard management practices were followed as and when necessary. No insecticide was used because the two advanced lines were insect resistant and diseases were not controlled (to identify susceptibility and tolerance level of lines). Date of seeding, transplanting, flowering and maturity, lodging tolerance, pest and disease incidence, phenotypic acceptance at vegetative and ripening stage, yield and yield components were recorded. Among the two advanced lines, BR8693-8-4-2-1 **line was statistically significant from BR8693-17-6-2-2, BRRI Dhan49 and BRRI Dhan33 respectively and the disease reaction, farmers' opinion, and special character of insect resistant, BR8693-8-4-2-1 was found to be superior for further variety trial.** The main purpose of the experiment is to recommended insect resistant rice genotype for proposed variety trail. For yield estimation, 9 m² sample area from each plot was harvested at maturity and grain yields were adjusted to 14% moisture content.

Keywords: Genotype, insect resistant, disease resistant, growth, yield

INTRODUCTION

The rice (*Oryza sativa* L.) plant belongs to the tribe of Oryzaceae under the sub-family of Pooideae in the grass family Gramineae (Poaceae). Recently the genus *Oryza* divided into several sections and placed *Oryza sativa* under the series of *Sativa* in section *Sativae*. Rice is indigenous to Asia. For rice plant, root is a fasciculate system and seems at the herbaceous plant, chill and water deficiencies sensitive^[12, 11]. The role of the root is extracted and absorption of dissolved minerals and water from the flooded regions^[16 and 35]. Rice (*Oryza sativa* L.), one of the major food crops, which is capable of feeding over half of the global population^[33]. The people of Asia are dominating regarding rice production as well as its consumption, almost 90% of the global rice production are contributed by them^[4]. Rice is the staple food crop in Bangladesh and occupies about 80% of the country's total cropped area^[10]. The total area of the country is 14.86 million ha (147,570 sq. km)^[31] and the cultivable area is 8.57 million ha^[21]. Rice is cultivating nearly in 11.42 million ha of which Aus, Aman and Boro rice are covered 9.3, 48.9 and 41.8%, respectively^[23, 5]. Rice is extensively grown in Bangladesh in the three seasons namely, **aus, aman (broadcasted and transplanted) and boro**^[3]. Among these cropping seasons transplanted Aman is the most important and occupied about 46% of the rice cultivation land in 2019-2020. Area covered by Aman rice is almost 5.53 million hectares and production is about 13.19 million metric ton^[6]. To increase the cultivation area as well as production, research should be emphasized regarding morpho-physiological and genetic characteristics of Aman rice in which a little information has been reported earlier^[19, 31]. One of the main constraints to increasing rice production in Bangladesh is the infestation of insects. According to an estimate, annual yield loss due to insect pest alone is 16% for rice. Global crop losses in rice due to weeds, animal pests, and diseases at 10.2%, 15.1%, and 12.2% of the attainable yield, respectively^[28]. In Bangladesh, chemical control has been the primary method of insect control. Bangladesh needs to increase its rice production on a sustainable basis. Insects continue to cause serious damages to rice crop and the use of toxic pesticides is the main method of pest control and that such continued heavy reliance on chemicals would lead to serious environmental and human health problems, pest resurgence, new pest problems and development of resistance.

Insect pest attacks frequently occur with varying intensities and frequencies possibly induced by the changes in climate and cropping systems in modern rice cultivation. Intensive rice production with the primary goal of achieving high yield is often characterized by the excessive application of fertilizers and pesticides. It has led to many negative environmental effects, such as the reduction of biodiversity and natural biological control, high pesticide residues in rivers, drinking water and agricultural products, rapid and high insecticide resistance in pests, secondary pest outbreaks, environmental pollution and ecological imbalance. These severe negative effects will damage the ecosystem, lead to frequent pest outbreaks and in turn require an increased pesticide dosage, which form a vicious circle^[13]. Pesticides used in the paddy fields globally account for nearly 15% of the total pesticides used for crop production.

The breeding of insect resistant cultivar is an effective and economical way to address this problem and some of these cultivars have been released as a variety. However, the conventional breeding progress towards insect resistant rice cultivars has been relatively slow^[14, 34 & 7], whereas **modern induced breeding is need to be adopted** which is an effective way to enhance the genetic variability^[27, 26]. On the other hand, **there are many rice varieties but** still we need more suitable rice varieties to address different situation, which can fulfill farmers' demands like higher yield, shorter growth duration with attractive grain quality and also insect and disease resistant rice varieties. We need new insect resistant rice varieties for transplanted Aman, which can replace

old varieties to overcome harmful effect of insecticides and can help achieve the target of sustainable and environmentally friendly agricultural production. Bangladesh has achieved self-sufficiency in food, especially in rice ^[21]. According to the Sustainable Development Goals (SDG) set in 2015 the target is to end hunger and ensure access by all people, in particular the poor and people in vulnerable situations, including infants, to safe, nutritious and sufficient food all year round by 2030 ^[33].

In the present study, two advanced lines are selected and the lines are BR8693-8-4-2-1 and BR8693-17-6-2-2 to identify their insect resistant response. Morphological traits, agronomical traits and physical traits were used to evaluate the resistant tolerance advanced lines.

MATERIALS AND METHOD

Two advanced lines: BR8693-8-4-2-1 and BR8693-17-6-2-2, along with BRRRI dhan49 (S. Ck) and BRRRI dhan33 (R. Ck) as checks were tested at farmers' field in eight locations such as West byde (BRRRI Gazipur), Khulna (Dumuria), Habiganj (Sadar), Chapainawabganj (Gomostapur), Rangpur (Sadar), Feni (Dagonbhuyan), Chattagram (Hathazari) and Barishal (Sadar) during the season of transplanting aman, 2018. The trials were replicated thrice in each location. The unit plot size was 17.94 m² (4.6m x 3.9m). Seeding emergence time for eight locations varied from 11-17 July 2018. About 20-25 days old seedlings were uprooted from the different locations. Seedlings were then transplanted with 20 x 15 cm spacing. N, P, K, S and Zn fertilizers were applied at the rate of 90, 15, 50, 12 and 3.6 kg ha⁻¹, respectively. All fertilizers except urea were applied during land preparation as basal whereas, urea was applied in 3 equal splits at 10, 25 and 40 days to transplanting (DAT). Other standard management practices were followed as and when necessary. No insecticide was used because the two advanced lines were insect resistant and diseases were not controlled (to identify susceptibility and tolerance level of lines). Date of seeding emergence, seedling transplanting, flowering and maturity, lodging tolerance, pest and disease incidence, phenotypic acceptance at vegetative and ripening stage, yield and yield components were recorded. Feedback from farmers and department of agricultural extension (DAE) personnel were also recorded. For yield estimation, 9 m² sample area from each plot was harvested at maturity and grain yields were adjusted to 14% moisture content.

RESULTS AND DISCUSSION

The trial site of Barishal (Sadar) was damaged due to 4-time tidal submergence. On an average, in rest of the seven locations, the tested two entries BR8693-8-4-2-1 and BR8693-17-6-2-2 gave higher yield (4.84 and 4.89 t ha⁻¹) than the check varieties BRRRI dhan49 (S. Ck.) (4.48 t ha⁻¹) and BRRRI dhan33 (R. Ck.) (4.06 t ha⁻¹) (Table 1). There might be seen varietal differences of grain yield ^[8]. The variety/genotypes which produced higher number of effective tillers/hill and higher number of grain/panicle also showed higher grain yield in rice ^[15]. On the other hand, yield differences recorded by ^[20] due to variable grain yield among varieties. Among the two advanced entries, BR8693-17-6-2-2 (entry no.2) gave mean 4.89 t ha⁻¹ grain yield, ranged from 4.46 to 5.17 t ha⁻¹ which was a bit higher than the another advanced line BR8693-8-4-2-1 (entry no.1) which gave mean 4.84 t ha⁻¹ ranged from 4.42 to 5.20 t ha⁻¹. But, the grain yield of the two advanced lines was statistically similar. There may have variation in grain filling due to genetic, environmental or cultural management practices adopted where ^[15] reported that grain yield was affected by the filled grains/panicle and correlated significantly with grain yield ^[24]. The mean growth duration of the advanced line BR8693-8-4-2-1 (entry no.1) was 133 ranged from 131 to 136 days. Whereas, the mean growth duration of another advanced line BR8693-17-6-2-2 (entry no.2) was 135 days ranged from 134-138. Check variety BRRRI dhan33 (R. Ck.) was found to be matured within the earliest mean growth duration (120 days), followed by in another check variety BRRRI dhan49 (132

days). The highest yielder (entry no. 2) matured within 135 days which was longest among the advanced lines including check varieties. Days to maturity variation was recorded due to different varieties/genotypes^[17] where^[18] reported that genotypic variation in phenological events among different cultivars. Demonstrated significant differences in attaining phenological stages due not only to varieties but also to variable management practices reported by^[1]. Among the tested advanced lines and check varieties the highest mean plant height (125 cm) was found in entry no. 2 ranged from 120 cm to 132 cm followed by 121 cm in entry no. 1 ranged from 114 cm to 128 cm. However, mean plant height was much lower in both the check varieties (101 cm in BRRIdhan49 and 104 cm in BRRIdhan3e 1). The mean plant height of the check varieties ranged from 98 to 109 cm. Mean 1000-grain weight was found lowest (20.90 g) in the check variety BRRIdhan49, ranged from 20.11 to 21.91 g and it was 23.76g for the another check variety BRRIdhan33 (22.96 to 24.89 g). Whereas, the highest mean 1000-grain weight (26.03 g) was found in the BR8693-17-6-2-2 (entry no.2) varied from 24.89 to 27.27 g. Mean 1000-grain weight of BR8693-8-4-2-1 (entry no.1) was 23.12 g (ranged from 23.06 to 25.80 g) which was a bit lower than the entry no. 2 (Table 2).

Table 1. Grain yield (t/ha), Growth duration (days) and plant height of some advanced Lines.

Genotypes Locations	Gazip WB	Khulna	Habiganj	Chapainawa	Rangpur	Feni	Chattogra	Mean
	Yield (t ha ⁻¹)							
BR8693-8-4-2-1	5.20	4.70	5.13	4.42	5.02	4.90	4.52	4.84
BR8693-17-6-2-2	4.63	4.46	5.17	5.04	4.83	5.01	5.11	4.89
BRRIdhan49(S.Ck)	5.04	4.75	5.04	4.37	4.82	3.64	3.69	4.48
BRRIdhan33(R.Ck)	4.45	4.37	4.06	4.12	4.19	3.72	3.51	4.06
LSD (0.05)	0.56							
CV%	7.37							
Growth duration (days)								
BR8693-8-4-2-1	136	135	131	132	133	134	133	133
BR8693-17-6-2-2	138	134	134	134	135	134	135	135
BRRIdhan49(S.Ck)	133	132	134	133	129	133	133	132
BRRIdhan33(R.Ck)	117	117	120	120	122	120	121	120
LSD (0.05)	0.78							
CV%	0.37							
Plant height (cm)								
BR8693-8-4-2-1	124	120	128	120	119	114	119	121
BR8693-17-6-2-2	126	123	132	128	122	123	120	125
BRRIdhan49 (S.Ck)	107	105	101	98	99	101	99	101
BRRIdhan33 (R.Ck)	109	107	102	104	101	105	100	104
LSD (0.05)	2.52							
CV%	1.30							

1000 grain weight differed significantly among the aman cultivars^[25] and^[29] found found difference in thousand weight of grains due to morphological and varietal variation. Highest panicles m⁻² (231) was observed in

BR8693-8-4-2-1 (entry no. 1) followed by BRR1 dhan49 (226), BR8693-17-6-2-2 (215) and the lowest panicles m^{-2} was found in the R. Ck., BRR1 dhan33. Panicles m^{-1} varied from 205-282 for the advanced line BR8693-8-4-2-1 (entry no. 1) and it varied from 182 to 268 for the advanced line BR8693-17-6-2-2 (entry no. 2) (Table 2).

Table 2. Yield components of some advanced lines.

Genotypes Locations	Gazip WB	Khulna	Habiganj	Chapainawa	Rangpur	Feni	Chattogra	Mean
1000-grain weight (gm)								
BR8693-8-4-2-1	23.06	23.73	25.03	24.57	25.20	25.07	25.80	23.12
BR8693-17-6-2-2	25.28	24.89	25.51	27.27	27.09	26.35	25.83	26.03
BRR1 dhan49 (S.Ck)	20.79	20.11	21.91	21.30	21.69	20.27	20.25	20.90
BRR1 dhan33 (R.Ck)	23.98	23.43	23.74	23.22	22.96	24.13	24.89	23.76
LSD (0.05)	0.85							0.32
CV%	2.24							
Panicles/m^2								
BR8693-8-4-2-1	248	243	282	209	226	206	205	231
BR8693-17-6-2-2	234	235	268	193	182	198	193	215
BRR1 dhan49 (S.Ck)	270	261	245	230	197	191	186	226
BRR1 dhan33 (R.Ck)	237	242	245	215	202	131	129	200
LSD (0.05)	20.74							7.83
CV%	5.58							
Filled grain per panicle								
BR8693-8-4-2-1	107	101	95	107	113	110	111	106
BR8693-17-6-2-2	94	92	98	107	104	106	108	101
BRR1 dhan49 (S.Ck)	113	105	106	101	126	97	102	107
BRR1 dhan33 (R.Ck)	94	92	87	93	104	114	107	99
LSD (0.05)	8.68							3.28
CV%	5.20							
Sterility (%)								
BR8693-8-4-2-1	18	20	22	23	17	17	17	19
BR8693-17-6-2-2	21	25	27	26	28	17	13	23
BRR1 dhan49 (S.Ck)	15	17	26	31	24	15	13	20
BRR1 dhan33 (R.Ck)	18	21	30	29	26	12	14	22
LSD (0.05)	5.93							2.24
CV%	17.61							

Filled grains/panicle of the tested advanced lines and check varieties varied from 99-107 and it was the highest (107) in the check variety BRR1 dhan49 (Table 2). Highest yield (entry no. 2) produced a bit lower filled grains panicle⁻¹ (101) than the check variety BRR1 dhan49 but higher than BRR1 dhan33. Sterility (%) of the entries including checks ranged from 19-23 (Table 2). Sterility variation occurred might be due to supply of insufficient

food materials, moisture and light for plant in closer spacing and rice produces 15-20% sterile grains ^[9] whereas ^[32] stated that spikelet sterility difference varied significantly by variety and plant spacing. Difference in plant height of the cultivars/varieties were due to varietal variation ^[30]. ^[2] stated that 50% days to flowering was positively and significantly correlated with days to maturity.

INSECT INFESTATION

Different insects' infestation was found in the different experimental sites. Stem borer (05-15%), Rice bug (05-10%) and Leaf folder (05-15%) were found in some trial locations. However, No insecticide was used because the two advanced lines were insect resistant rice.

Table 3. Insect infestation of some advance rice lines

Genotypes	Disease infection (%)*				
	Rice bug	Stem borer	Leaf folder	Brown Plant Hopper	Green Leaf Hopper
V ₁ = BR8693-8-4-2-1	5-8 in 5 loc.	4-10 in 2 loc.	5-12 in 3 loc.	3-4 in 2 loc.	2-4 in 2 loc.
V ₂ = BR8693-17-6-2-2	5-10 in 4 loc.	5-12 in 2 loc.	5-15 in 2 loc.	3-5 in 2 loc.	3-4 in 2 loc.
V ₃ = BRRI dhan49 (S.Ck.)	4-9 in 5 loc.	5-15 in 1 loc.	4-14 in 2 loc.	4-7 in 3 loc.	2-5 in 3 loc.
V ₄ = BRRI dhan33 (R. Ck.)	8-10 in 3 loc.	10 in 1 loc.	4-12 in 2 loc.	5-7 in 2 loc.	3-5 in 2 loc.

DISEASE INFECTION

Different diseases infection was found in the different experimental sites. From Table 4, V₁ = BR8693-8-4-2-1 and V₂ = BR8693-17-6-2-2 both the lines are comparatively less infected by sheath blight, Bacterial leaf blight, leaf blast, false smut, Bacterial leaf streak, Tungro and Grain spot as per checked varieties of V₃ = BRRI dhan49 (S.Ck.) and V₄ = BRRI dhan33 (R. Ck.).

Table 4. Disease infestation of some advance lines

Genotypes	Disease infection (%)*						
	Sheath blight	Bacterial leaf blight	Leaf Blast	False smut	Bacterial leaf streak	Tungro	Grain spot
V ₁ = BR8693-8-4-2-1	1-4 in 5 loc.	3-4 in 2 loc.	4-6 in 3 loc.	1 in 2 loc.	0	10-15 in 1 loc.	0
V ₂ = BR8693-17-6-2-2	1-3 in 4 loc.	2-3 in 2 loc.	3-6 in 2 loc.	1 in 2 loc.	0	10-15 in 1 loc.	1-5 in 2 loc.
V ₃ = BRRI dhan49 (S.Ck.)	1-5 in 5 loc.	1-4 in 1 loc.	4-7 in 2 loc.	2-12, in 3 loc.	0	10-15 in 1 loc.	1-2 in 1 loc.
V ₄ = BRRI dhan33 (R. Ck.)	3 in 3 loc.	3 in 1 loc.	3-7 in 2 loc.	1-2 in 2 loc.	0	10-15 in 1 loc.	4 in 1 loc.

*Eye estimation of the number of hills showing the sign and symptom of disease infection. The percentage indicates the disease incidence level of the tested genotypes.

Lodging incidence: The two advanced lines were found lodging tolerant in all locations.

Table 5: PHENOTYPIC ACCEPTANCE

Entry no.	Characteristics						P. acp Score	
	Plant growth	Uniformity of flowering	Uniformity of maturity	Wrapping quality of culm	Grain type	Flag leaf	Veg.	Mat.
1	Fair	regular	regular	Well wrapped	Medium bold	Erect	5	5
2	Fair	irregular	irregular	Well wrapped	Medium bold	Erect	5	7
3	Good	regular	regular	Well wrapped	Medium fine	Erect	3	3
4	Good	regular	regular	Well wrapped	Medium bold	Erect	3	5

Phenotypic Acceptability: 1= Excellent, 3= Good, 5= Fair, 7= Poor, 9= Unacceptable

CONCLUSION

Most of the farmers showed their interest about the two advanced lines due to their higher grain yield than the check varieties i.e., BRRI dhan49 and BRRI dhan33. Farmers preferred entry no. 1 due to similar growth duration to BRRI dhan49. But most of the farmers dislike bold type grain size of the two advanced lines. In this study, correlative and comprehensive observations of physiological traits were performed including morphological development, agronomic traits. The results showed that both the advanced lines gave statistically higher yield than the check variety BRRI dhan49 and BRRI dhan33. But growth duration of BR8693-8-4-2-1 (entry no. 1) is lower than BR8693-17-6-2-2 (entry no. 2), but similar to BRRI dhan49. Though BR8693-8-4-2-1 (entry no. 1) is statistically significant from BR8693-17-6-2-2 (entry no. 2), BRRI dhan49 and BRRI dhan33 which would be recommended insect resistant rice genotype and considering the above results and following information regarding disease reaction, farmers' opinion, and special character of insect resistant, entry no. 1 (BR8693-8-4-2-1 line) was found to be suitable for Proposed Variety Trial (PVT).

ACKNOWLEDGEMENTS

Authors would like to express their sincere gratitude to the staffs of Bangladesh Rice Research Institute (BRRI), Gazipur, Bangladesh for their helpful cooperation during the entire research work.

REFERENCES

1. Ahmed, A. R., Dutta, B. K. and Ray, D. C. (2015). Response of some rice varieties to different crop management practices towards morphological and yield parameters. *Intl. J. Sci. Res. Pub.* 5(2), 1-6.
2. Alam, M. S., Islam, M. M., Hassan, L., Begum, S. N. and Gupta, R. (2014). Study of Correlation, Magnitude of Genetic Diversity and Selection Indices in Popular Rice (*Oryza sativa* L.) Landraces of Bangladesh. *International Journal of Innovation and Applied Studies*, 8 (3), 1329-1338.
3. AIS (Agricultural Information Service), Krishi Dairy. Agriculture Information Service, Khamarbari, Farmgate, Dhaka. p. 10-15, 2011.
4. Bandumula N., Rice production in Asia: Key to global food security. The Proceedings of the National Academy of Sciences, India, Section B: Biological Sciences DOI 10.1007/s40011- 017-0867-7, 2017.
5. BBS (Bangladesh Bureau of Statistics), Agriculture statistics of major crops. Statistics and Informatics Division (SID), Bangladesh Bureau of Statistics, Government of the People's Republic of Bangladesh. pp. 21-22, 2018.
6. BBS (Bangladesh Bureau of Statistics), Forty five years Agriculture Statistics of Major Crops (Aus, Amon, Boro, Jute, Potato & Wheat). Bangladesh Bureau of Statistics Division, Government of the People's Republic of Bangladesh. p. 22, 2018.
7. Bernier J, Atlin G N, Serraj R, Kumar A, Spaner D., Breeding upland rice for drought resistance. *Journal of the Science of Food and Agriculture*, **88**, 927–939, 2008.
8. Biswas, J. K., Hossain, M. A., Sarker, B. C., Hassan, M. and Haque, M. Z. (1998). Yield performance of several rice varieties seeded directly as late aman crops. *Bangladesh J. Life Sci.*, 10, 47–52.
9. BRKB (2016). Rice production problem, 5th edition. p.70. Retrived 13 January 2017 from <http://knowledgebank-brrri.org/Publications/Rice-production-problem/booklet-rice-production-problem.pdf>
10. BRRRI (Bangladesh Rice Research Institute), Adhunik Dhaner Chash (Cultivation of Modern Rice), Gazipur, Bangladesh. pp. 5-93, 2019.
11. Chun, L., Wang, G., Chen, P., Zhu, H., Wang, S., Ding, Y., Shoot-Root Communication Plays a Key Role in Physiological Alterations of Rice (*Oryza sativa*) Under Iron Deficiency, *Front Plant Sci.* 9:757, 2018.
12. Comas, L. H., Becker, S. R., Cruz, V. M. V., Byrne, P. F., Dierig, D. A., Root traits contributing to plant productivity under drought, *Front. Plant Sci.* 4:442, 2013.
13. Conway G R, Pretty J N., *Unwelcome Harvest: Agriculture and Pollution*. London: Earthscan Publications Ltd, 1991.
14. Cooper M, Fukai S, Wade L J., How can breeding contribute to more productive and sustainable rainfed lowland rice systems? *Field Crops Research*, **64**, 199–209, 1999.
15. Dutta, R. K., BasetMia, M. A. and Khanam, S. (2002). Plant architecture and growth characteristics of fine grain and aromatic rices and their relation with grain yield. *Intl. Rice Comm. Newslett.* 51, 51–56.
16. Fageria, N. K, Moreira, A., The Role of Mineral Nutrition on Root Growth of Crop Plants. *Advances in agronomy*, 110(1):251- 331, 2011.
17. Ghosh, U. K., Khan, A. R. and Karim, M. A. (2015). Growth performance of aus rice varieties under rainfed condition. *International Journal of Advanced Multidisciplinary Research* 2(11), 29–35.

18. Haque, K. M. S., Karim, M. A., Bari, M. N. and Islam, M. R. (2016). Genotypic variation in the effect of drought stress on phenology, morphology and yield of aus rice. *Intl. J. Biosci.* 8(6), 73-82. <https://doi.org/10.12692/ijb/8.6.73-82>
19. Hossain MS, Monshi FI, Kamal AMA, Miah MF, Grain yield and protein content of transplant aman rice as influenced by variety and rate of nitrogen. *Journal of Agroforestry and Environment* 3(2): 235-238, 2010.
20. Islam, M. S., Paul, S. K. and Sarkar, M. A. R. (2014). Varietal performance of modern transplant Aman rice subjected to level of nitrogen application. *J. Bangladesh Agril. Univ.* 12 (1), 55-60. <https://doi.org/10.3329/jbau.v12i1.21239>.
21. Kabir, M S, M U Salam, A Chowdhury, N M F Rahman, K M Iftekharuddaula, M S Rahman, M H Rashid, S S Dipti, A Islam, M A Latif, A K M S Islam, M M Hossain, B Nessa, T H Ansari, M A Ali and J K Biswas. (2015). Rice Vision for Bangladesh: 2050 and Beyond. *Bangladesh Rice J.* 19(2):1-18.
22. Karmakar, B, and M A R Sarkar., Optimizing seedling age of promising rice genotypes in rainfed environment. *J. Crop and Weed.* 11(Special Issue):149-160, 2015.
23. Karmakar, B and M A Ali, Production and Preservation of Quality Rice Seed. 1st Edition, Bangladesh Rice Research Institute, Gazipur. pp.136, 2019.
24. Kiani, G. and Nematzadeh, G. (2012). Correlation and path coefficient studies in F2 populations of rice. *Notulae Scientia Biologicae*, 4(2), 124-127.
25. Mondal, M. M. A., Islam, A. F. M. S. and Siddique, M. A. (2005). Performance of 17 modern transplant aman cultivar in the north region of Bangladesh. *Bangladesh J. Crop. Sci.* 16, 23-29.
26. Mondal S, Petwal V C, Badigannavar A M, Bhad P G, Verma V P, Goswami S G, Dwivedi J. (2017). Electron beam irradiation revealed genetic differences in radio-sensitivity and generated mutants in groundnut (*Arachis hypogaea* L.). *Applied Radiation and Isotopes*, **122**, 78–83.
27. Mostafa H A, Wang H P, Shen D, Yang Q, Li X., Sprout differentiation and mutation induction of garlic (*Allium sativum* L.) callus exposed to gamma radiation. *Plant Growth Regulation*, **75**, 465–471, 2015.
28. Oerke EC. 2006. Crop losses to pests. *J. Agric. Sci.* 144:31-43.
29. Roy, S. K., Ali, M. Y., Jahan, M. S., Saha, U. K., Ahmad-Hamdan, M. S, Hassan, M. M. and Alam, M. M. (2014). Evaluation of growth and yield attributing characteristics of indigenous boro rice varieties. *Life Sci. J.* 11, 122-126.
30. Sarkar, S. C. (2014). Performance of five selected hybrid rice varieties in aman season. M.S. Thesis, Dept of Agricultural Botany, Sher-e-Bangla Agril, Univ., Dhaka. pp. 25-26, 44-46. Retrived from http://archive.saulibrary.edu.bd:8080/bitstream/handle/123456789/342/Bot-SAU201401_12-08-02835.pdf?sequence=1
31. Shelley, I J, M T Nosaka, M K Nakata, M S Haque and Y Inukai., Rice Cultivation in Bangladesh: Present Scenario, Problems, and Prospects, *J. Intl. Cooper. Agric. Dev.* 14: 20-29, 2016.
32. Sohel, M. A. T., Siddique, M. A. B., Asaduzzaman, M., Alam, M. N. and Karim, M. M. (2009). Varietal performance of transplant aman rice under differnt hill densities. *Bangladesh J. Agril. Res.* 34(1), 33-39.
33. USDA, Rice Sector at a Glance. <https://www.ers.usda.gov/topics/crops/rice/rice-sector-at-a-glance/>, 2020.

34. Wade L J, McLaren C G, Quintana L, Harnpichitvitaya D, Rajatasereekul S, Sarawgi A K, Kumar A, Ahmed H U, Singh A K, Rodriguez R., Genotype by environment interactions across diverse rainfed lowland rice environments. *Field Crops Research*, **64**, 35–50, 1999.
35. Yamauchi, T., Colmer, T. D., Pedersen, O., Nakazono, M., Regulation of Root Traits for Internal Aeration and Tolerance to Soil Waterlogging-Flooding Stress, *Plant Physiol.*, 176, 1118– 1130, 2018.

UNDER PEER REVIEW