

Original Research Article

Proximate Composition Analysis of *Spirulina Platensis* in Lab Scale Cultivation: Prospects of Digested Rotten Guava as a Culture Media

Abstract

An experiment was conducted to evaluate the proximate composition of *Spirulina platensis* and growth performance in supernatant of different concentrations (supernatant of 20, 40 and 60 rotten guavas) of digested rotten guava medium (DRGM). The growth rates in terms of optical density, dry cell weight and chlorophyll *a* of spirulina were varied from each other. The cell weight of Spirulina was attained a maximum of $0.818 \pm 0.003 \text{ mg/L}$ in 60% DRGM followed by 0.815 ± 0.0015 and $0.809 \pm 0.0012 \text{ mg/L}$ in supernatant of 20 and 40%, respectively on the 10th day of culture. Similar trend was also observed in the case of optical density of the media contained Spirulina, chlorophyll *a* content (mg/L), total biomass (mg/L), specific growth rates (on the basis of cell weight and chlorophyll *a*) and total biomass of Spirulina. Cell weight of Spirulina grown in these media had highly significant ($P < 0.01$) correlation with the chlorophyll *a* content ($r = 0.746$) and total biomass ($r = 0.742$) of Spirulina. The growth performance of Spirulina in supernatant of 60% DRGM was significantly ($P < 0.01$) higher than that of grown in 20% and 40% DRGM. The percentage of crude protein was found $53.35 \pm 0.32\%$ in supernatant of 40% DRGM. The crude lipid was attained of $10.15 \pm 0.14\%$ in supernatant of 60% DRGM which was significantly ($P < 0.05$) higher than that of grown in 20% and 40% of DRGM.

Keywords: *Spirulina Platensis*, Proximate composition, Growth performance, Digested Rotten Guava Medium (DRGM)

Introduction

With the aim of increased aquaculture production through applying adequate feed large numbers of feed industries are developed in the country (Name of Country?). Due to increased aquaculture practice, demand of good quality feed is increasing day by day. Prime quality feed is essential for fish growth. Maintain feed conversion ratio (FCR) close to 1 is highly depends upon good feed. Feed should have adequate protein content which facilitates high growth. Net protein utilization should be around 27 percent (Review -here?). But fish meal and bone meal are not available in our country (Name of Country?). So, we can find to alternative sources. We can use to alternative fish meal to spirulina. Spirulina is a “superfood” which is the most nutritious, rich in protein and concentrated whole food known to humankind (Review -here?). It has a vibrant history occupied an intriguing biological and ecological niche in the plant kingdom. Spirulina is a spiral-shaped, blue-green microalgae that grows naturally in the wild in freshwater

alkaline lakes, natural springs, and saltwater. Its deep blue green color is what gives the water greenish hue ([Review -here?](#)). Spirulina is also cultivated and harvested in man-made reservoirs around the world. For centuries, civilizations the world over have cultivated and cherished spirulina for its health-improving benefits. The Aztecs harvested the spirulina from Lake Texcoco in Mexico. It grows well in supernatant of different digested agro-industrial wastes available in Bangladesh ([Satter, 2017](#)).

Spirulina may be grown in agro-industrial wastes ([Review -here?](#)), rotten fruits ([Review -here?](#)), chicken wastes ([Review -here?](#)). Among fruits huge quantity of guava spoils (rotten) in different markets in the country. Therefore, market is allowed to digest (aerobic & anaerobic) and supernatant may be used for the growth of spirulina. This inexpensive low-cost medium may be used to produce *Spirulinaplatensis* culture which can contribute significantly for the development of fisheries and fish production. It takes inorganic nutrients for the supernatant and grows. Many factors are important for the production of spirulina at large scale, of which most important factors are nutrient availability, temperature and light. The filamentous cyanobacteria such as spirulina are found to be most compatible microorganisms for the utilization of waste and wastewaters as they are able to produce large quantity of biomass and their harvesting is also relatively easy because of their structure. Also, these wastes reduce the cost of nutrient medium and act as a source of cheap nutrient medium for cultivation of spirulina. The commercial production of spirulina can be made cost effective by reducing the input cost with cheap and readily available materials without sacrificing the production efficiency. They are very small a microscopic and 300-500 micrometer in length. Spirulina contains 50-70% protein, 10-12% carbohydrate, 6% fat, 7% minerals and a lot of vitamins ([Review -here?](#)). However, according to the researchers, one kg of *Spirulina*spp is similar to 1000 kg of other vegetables ([Kato 1991](#)).

Spirulina has been studied for single cell protein (SCP), vitamins, minerals, proteins and polyunsaturated fatty acids (γ -linolenic acid), therapeutic properties, antioxidant activity. Several cultivation methods like open ponds, tubular photo bioreactors, inclined glass panels have been tried. Cost and composition of cultivation media along with growth rate of the algae are challenging factors for commercially viable production. The most convincing trials are of course those conducted among populations which traditionally eat spirulina. Culture and growth performance of *Spirulinaplatensis* in supernatant of digested rotten guava to evaluate the proximate composition of spirulina; to analyze growth parameters in supernatant digested rotten guava; and to find out the suitable concentration of the medium for maximum growth of *Spirulinaplatensis*.

Materials and Method

Study Area

The study carried out in Live Food Aquaculture Laboratory, Department Aquaculture, Faculty of Fisheries, Bangladesh Agricultural University (BAU), Mymensingh-2202, and Bangladesh.

Collection of Rotten Guava

The rotten guava was selected as medium for *Spirulinaplantensis* culture. It was collected from Kamal Ronjit market (K.R.) of Bangladesh Agricultural University, Mymensingh-2202, and Bangladesh. It was thought that the proximate composition of this media might be suitable for the growth of culture species.

Analysis of proximate composition of rotten guava (RG)

Where is Table ?????proximate composition of rotten guava?

Before media preparation, the proximate composition of rotten guava was analyzed to know its nutritional status. The analysis was performed in Fish Nutrition Laboratory, Department Aquaculture, Faculty of Fisheries, Bangladesh Agricultural University (BAU), Mymensingh-2202, and Bangladesh, following standard methods ([Horwitz, 1984](#)).

Moisture

The sample was weighed in a previously pre-weighed small crucible in triplicates. The samples contained in the crucible were dried to moisture from it 105°C for 24 hrs. After drying, the samples with crucible were cooled with the help in a desiccator. Then, cooling at room temperature and weighed in a sensitive balance. The percentage of moisture of the sample was calculated using the following equation:

$$\% \text{ moisture} = \frac{X - Y}{X} \times 100$$

Where, X = Weight of sample before drying; and
Y = Weight of sample after drying.

Crude protein

Kjeldhal Auto 1030 Analyzer was used for determination of crude protein content of samples. A sample of 0.5g and a blank were digested in the digestion tube. 10 ml of concentrated sulphuric acid (H₂SO₄), 2.0 ml of H₂O₂ and one Kjeldhal tablet were added in the tubes and mixed gently by electric mixer. Then, the digestion tubes were set in digestion chamber fixing at 420°C for 45 minutes. The digestion the tubes were allowed to cool and 75 ml of distilled water was added in each tube. 50 ml of 40% NaOH was added before titration. After titration with 1% boric acid and 0.2 N HCl the reading for the samples and blank were recorded. The readings were calculated the following formula:

$$\text{Percentage of nitrogen} = \frac{\text{Milliequivalent wt. of N} \times \text{ml of titrant} \times \text{strength of HCl}}{\text{Samples wt. (g)}} \times 100$$

For animal, % Protein = % Nitrogen x 6.25; and

For plant, % Protein = % Nitrogen x 5.85

Crude lipid

Lipid content was estimated by solvent extraction of lipid using Soxlet apparatus. 2.0 g dried samples were taken into extraction thimbles to place into the extraction unit along with the weighed extraction cups having 50 ml of solvent as acetone. Extraction of first 15 minutes was in the boiling position and the cups were released and dried in the oven for 30 minutes. The percentage of crude lipid was determined using the following equation:

$$\% \text{ Lipid} = \frac{\text{Weight of cup with lipid} - \text{initial weight of cup}}{\text{weight of sample (g)}} \times 100$$

Ash

The pre-weighed crucible containing dried sample from the moisture determination was pre-ashed. The samples were kept into a muffle furnace at 550°C for 6.0 hrs. The crucible containing ash was cooled in a desiccator. The percentage of ash was determined by using the following:

$$\% \text{ Ash} = \frac{\text{Weight of ash with preweighed crucible} - \text{weight of crucible}}{\text{weight of dried sample}} \times 100$$

NFE

NFE were calculated the following formula:

Nitrogen Free Extract = 100 - (Moisture + Crude protein + Crude lipids + Ash).

Crude fiber(Where is Method ??)

Optical density of media contained spirulina(Where is Method ??)

Chlorophyll a(Where is Method ??)

Total biomass of spirulina(Where is Method ??)

Cell weight (mg/L)(Where is Method ??)

Specific growth rates (SGRs) on the basis of cell weight, chlorophyll a and total biomass of Spirulina platensis(Where is Method ??)

Culture and collection of Spirulina platensis

Spirulina platensis was collected from the stock in the live food culture laboratory, Department of aquaculture, Faculty of Fisheries, Bangladesh Agricultural University (BAU), Mymensingh-2202, and Bangladesh. Twelve conical flasks (2 L capacity) were used for the culture of spirulina.

Maintenance of pure stock culture of Spirulina platensis

Pure stock culture of *Spirulinaplatensis* was maintained in the laboratory (Modified after [Zarrouk's, 1996](#)). Growth of *Spirulinaplatensis* were observed at every alternative day and was checked under microscope to confirm its purity following some keys given by [Bold and Wynne \(1978\)](#), [Vymazal \(1995\)](#) and [Phang and Chu \(1999\)](#).

Preparation of digested rotten guava media (DRGM)

According to (Review -here?), compositions of Rotten Guava Medium (RGM) were prepared for culture of *Spirulinaplatensis*. 50 g/L rotten guava was allowed to decompose in 5.0 L glass bottle for 34 days under aerobic condition in the Live Food Culture laboratory, Department of Aquaculture, BAU, Mymensingh. Then a Light reddish white colored supernatant from bottle was diluted and made three concentrations at the rate of 20%, 40% and 60% digested rotten guava. Then the supernatant of three different concentrations were taken in 1.0 L flask with three replications. For the preparation of rotten guava medium, digested and continuous aeration 5 liter volumetric flask was filtered with plankton net after (10.10.18 to 14.11.2018) 34 days left. Then the filtered rotten guava was diluted and added 0.8 g (0.2 g/L) urea according to the above direction with three replications using distilled water. Then the medium was mixed well and sterilized at 115°C for 15 minutes by high pressure bumping water autoclave. After autoclaving, the media were kept 3 days to be sure about any contamination free before culture of micro algae. Mixing, autoclaving and cooling were carried out pursuing the procedure used during the preparation of digested rotten guava media.

Experimental design of Spirulinaplatensis culture

Three types media viz., Rotten guava (RG) were used to culture *Spirulinaplatensis*. Inoculum of *Spirulinaplatensis* was collected from the pure stock culture. Experimental design is shown in (Table 1).

Table 1: Experimental design for *Spirulinaplatensis* culture using supernatant of three different concentrations of digested rotten guava (DRG)

Types of medium	Treatments	Replications	Amounts rotten guava (%)	Duration of culture (days)
Supernatant of DRGM	1	3 (101, 102 and 103)	20	14
	2	3 (201, 202 and 203)	40	
	3	3 (301, 302 and 303)	60	

Analysis of Proximate Composition Spirulina

Best optical density of spirulina (*S. platensis*) was found on 10th day of culture. In that day, spirulina was filtered, collected in petridish and kept in an oven at 40°C for overnight for drying. Then the proximate composition of cultured *S. platensis* was analyzed in the Fish nutrition Laboratory, Faculty of Fisheries, BAU, Mymensingh by standard methods ([Horwitz, 1984](#)). All the procedures for determination of proximate composition, the procedures as described in the Chapter 3.2.2 were followed.

Statistical Analysis

Analysis of variance (ANOVA) of mean cell weight and chlorophyll *a* of *S. platensis* cultured in different media (treatments) were done and to find whether any significant among treatment mean was done by Duncan's Multiple Range Test (DMRT) at 5% level of probability (Zar, 1984).

Results

Proximate composition (%) of spirulina (Spirulinaplatensis)

There was no significant variation among the crude protein of *Spirulina* grown in the supernatant of three different DRG (Table 2). The percentage of crude protein of *Spirulina* was 53.25 ± 0.32 , 53.35 ± 0.34 and $53.28 \pm 0.32\%$ when grown in the supernatant of 20, 40 and 60% DRG media, respectively. There was no significant ($P > 0.05$) variation among the crude protein percent of spirulina cultured in three different media of DRG. There was no significant ($P > 0.05$) difference of crude lipids of spirulina when cultured in the supernatant of 20, 40 and 60% digested rotten guava. Ash (%) of *Spirulina* grown in supernatant of 20% ($10.22 \pm 0.13\%$) 40% ($10.16 \pm 0.17\%$) and 60% ($10.33 \pm 0.21\%$) digested rotten guava. There was no significant ($P > 0.01$) ($P > 0.05$) difference among the ash of spirulina grown in supernatant of 20, 40 and 60% DRG. There was no significant variation among the NFEs of spirulina grown in supernatant of 20, 40 and 60% DRG as the values found for 20% ($16.28 \pm 0.32\%$), 40% ($16.10 \pm 0.18\%$) and 60% ($15.99 \pm 0.22\%$), respectively.

Very small amount of crude fibre (%) was found in spirulina grown in the supernatant of three different digested rotten guava (DRG). However, it was varied from $0.69 \pm 0.0\%$ when *Spirulina* grown in the supernatant of 40% DRG to $0.71 \pm 0.04\%$ and $0.71 \pm 0.03\%$ when cultured in the same of 60% DRG, respectively (Table-2).

Table 2. Proximate composition (% in dry matter basis) of *Spirulinaplatensis*

Treatments	T1 (20% DRG)	T2 (40% DRG)	T3 (60% DRG)
Moisture	9.44 ± 0.05	9.55 ± 0.04	9.53 ± 0.05
Crude Protein	53.25 ± 0.32^b	53.35 ± 0.34^b	53.28 ± 0.32^b
Crude Lipids	10.10 ± 0.16^a	10.14 ± 0.17^a	10.15 ± 0.14^a
Ash	10.22 ± 0.13^b	10.16 ± 0.17^b	10.33 ± 0.21^b
NFE*	16.28 ± 0.32^a	16.10 ± 0.18^a	15.99 ± 0.22^a
Crude Fiber	0.70 ± 0.04	0.69 ± 0.03	0.71 ± 0.04

*NFE (Nitrogen Free Extract) = $100 - (\text{Moisture} + \text{Crude protein} + \text{Crude lipids} + \text{Ash})$. Figures in common letters in the same row do not differ significantly at **5% level of probability**.

Optical density of media contained spirulina

Optical density (OD) of media contained spirulina was found to increased up to 10th day of culture in all the media of digested rotten guava (DRG), and Kosaric medium and then decreased up to 14th day of experiment (Fig.1). However, highest OD of 20% DRG culture contained spirulina was 0.631 ± 0.0023 , where highest OD of 40% DRG culture contained

spirulina was found 0.704 ± 0.0015 . The OD of supernatant of 60% DRG contained spirulina was 0.725 ± 0.0012 .




Fig.1. Mean values of optical density of media contained *Spirulina platensis*

Chlorophyll a of spirulina

Chlorophyll a of spirulina was found also higher on 10th day of culture than other days of culture in supernatant of all the media (Fig.2). Chlorophyll a of spirulina increased from first day (0.00158 ± 0 g/L) up to 10th day (0.770 ± 0.0012 g/L) of culture in 20% digested rotten guava media (DRTM) and then decreased up to 14th day (0.710 ± 0.0012 g/L) of experiment. However, chlorophyll a of spirulina cultured in supernatant of 40% DRG was 0.768 ± 0.0012 g/L on 10th day and then decreased up to 14th day (last day) of culture. Chlorophyll a of spirulina grown in supernatant of 60% DRG was 7.365 ± 0.20 g/L on 10th day from first day (0.0016 ± 0) and then decreased up to 14th day (last day) of experiment (Fig.2).




Fig.2. Mean values of chlorophyll *a* (mg/L) of *Spirulina platensis*

Total biomass of spirulina

Total biomass of spirulina was increased from first day (0.106 ± 0.003) g/L or mg/L up to 10th day (67.77 ± 0.44 g/L g/L or mg/L) in the culture of 20% digested rotten guava media (DRGM) and then decreased up to 14th day (47.57 ± 0.42 g/L g/L or mg/L) of experiment. The highest total biomass of spirulina grown in the culture of 40% DRGM was recorded 51.46 ± 0.28 g/L g/L or mg/L on 10th day of culture and then decreased up to 14th day (42.35 ± 0.19 g/L g/L or mg/L) during the experiment. Total biomass of spirulina cultured in the culture of 60% DRGM was increased from first day (0.107 ± 0.04 mg/L) up to 10th day (57.75 ± 0.20 g/L) and then decreased up to 14th day (41.88 ± 0.14 g/L g/L or mg/L) of experiment (Fig.3).




Fig. 3. Mean values of total biomass (mg/L) of *Spirulina platensis* Where is distribution total biomass(0 to 100)

Comparison of growth parameters of spirulina

Optical density of 20% DRG (0.631 ± 0.002), 40% DRG (0.704 ± 0.0015) and 60% DRG (0.725 ± 0.0012). There was no significant ($P > 0.05$) difference among optical densities of 20, 40 and 60% DRG during the study. There was no significant ($P > 0.05$) difference of cell weight of spirulina grown in 20, 40 and 60% DRG. Chlorophyll *a* (10.53 ± 0.32 mg/L) of spirulina grown in 60% DRG (0.862 ± 0.0012 mg/L) was significantly ($P < 0.01$ or $P < 0.05$) higher than that of spirulina cultured in 40% (0.768 ± 0.0012 mg/L) and 20% DRG (0.770 ± 0.14 mg/L). There was no significant difference among the Chlorophyll *a* of spirulina grown in supernatant of 20, 40 and 60% DRG during the study (Table-3).

Table 3. Comparison of cell weight, chlorophyll *a* and total biomass of *Spirulina platensis*

Parameters	T1 (20% DRG)	T2 (40% DRG)	T3 (60% DRG)
Optical density	0.631 ± 0.002^b	0.704 ± 0.0015^b	0.725 ± 0.0012^b
Cell weight (mg/L)	0.815 ± 0.0015^b	0.809 ± 0.0012^b	0.818 ± 0.0013^b
Chlorophyll <i>a</i> (mg/L)	0.770 ± 0.14^b	0.768 ± 0.0012^b	0.862 ± 0.0012^b
Total biomass(mg/L) *	67.77 ± 0.43^b	51.46 ± 0.28^c	57.75 ± 0.20^{bc}

*Total biomass = Chlorophyll *a* x 67 (Vonshak and Richmond, 1988). Figures in common letters do not differ significantly at **5% level of probability**.

Correlation among the growth parameters of spirulina

Cell weight of spirulina (*Spirulinaplatensis*) had highly significant ($P < 0.01$ (**$P < 0.01$ Or $P < 0.05$**)) direct correlation with chlorophyll *a* ($r = 0.746$) of spirulina grown in the supernatant of different digested rotten guava. Similarly, total biomass of *S. platensis* was highly ($P < 0.01$ (**$P < 0.01$ Or $P < 0.05$**)) and directly correlated with chlorophyll *a* ($r = 0.795$) of spirulina cultured in the supernatant of various digested rotten guava. Again, total biomass of spirulina was found to be highly ($P < 0.01$ (**$P < 0.01$ Or $P < 0.05$**)) and directly correlated with the cell weight ($r = 0.742$) of spirulina grown in the supernatant of different digested rotten guava.

Specific Growth Rate (SGR) in respect to total biomass of spirulina

The SGR in respect to total biomass of spirulina was significantly ($P < 0.05$) varied from that of spirulina grown in the supernatant of 20, 40 and 60% DRG. There was no significant ($P < 0.05$) difference recorded among the SGRs on the basis of total biomass of *S. platensis* grown in the supernatant of 20, 40 and 60% DRG (Table-4).

Table 4. Specific growth rates (SGRs) on the basis of cell weight, chlorophyll *a* and total biomass of *Spirulinaplatensis*

Parameters	T1 (20% DRG)	T2 (40% DRG)	T3 (60% DRG)
SGR of cell weight	0.21 ± 0.010^b	0.22 ± 0.009^b	0.21 ± 0.011^b
SGR of Chlorophyll <i>a</i>	0.22 ± 0.004^b	0.21 ± 0.003^b	0.23 ± 0.004^b
SGR of total biomass	0.54 ± 0.012^b	0.52 ± 0.012^b	0.53 ± 0.013^b

Figures in common letters in the same row do not differ significantly at 5% level of probability.

DISCUSSION

The cell weight of *Spirulinaplatensis* in supernatant of digested rotten guava were found 0.0023 to 0.815 mg/L in 20% digested rotten guava media (DRGM), 0.0024 to 0.809 mg/L in 40% DRGM, 0.0023 to 0.818 mg/L in 60% DRGM. The growth performance of *Spirulinaplatensis* in supernatant of 60% DRGM was found better than 20% and 40% DRGM. This variation might be due to the differences in nutrient concentrations and composition of varied media (**Review - here?**). On the other hand 40% DRGM showed lower growth performance of *Spirulinaplatensis* in relation to 20 and 60% DRGM. This might be due to lower nitrogen and phosphate concentration of the nutrients in the media (**Review -here?**). The concentration of 60% DRGM which were better for the growth of *Spirulinaplatensis* because of the nutrient content **Where is Table ????? proximate composition of rotten guava?**

During the present study, digested organic medium like rotten guava which has the similarity with the findings of **Dineshkumaret al.,(2016) and Sukumaranet al., (2018).** During culture of

Spirulina platensis, the exponential phase was found up to 10th day from the beginning and then the cell weight declined i.e., stationary phase started. **Satter (2017)** recorded the cell weight and chlorophyll *a* content of *S. platensis* was significant ($P < 0.05$) higher in 4.0 g/L digested poultry waste than other media where light intensity, aeration and temperature played significant role to the culture system.

Similarly, **Sharker (2002)** conducted an experiment on the culture of *Spirulina platensis* in various concentrations viz., 0.3, 0.4 and 0.5 g/L of papaya skin powder medium (PSPM) and Kosaric medium (KM) in the laboratory for three months carried out for a period of 12 days. In the present study, the chlorophyll *a* content of inoculated *Spirulina platensis* was 0.0015 mg/L which attained a high content of 0.862 mg/L in 60% DRGM at the 10th day of culture. These findings are not more or less similar with the findings of **Phanget al., (2000), Habibet al., (2003) and Satter (2017)**. This might be due to lower nitrogen and phosphate concentration of the nutrients in the media. **Dineshkumaret al., (2016)** studied that *Spirulina platensis* grew well in natural medium such as Conway medium, Zoarrouic medium, **kosaric medium** and **BGI(What is meaning?)** medium.

In the present study, supernatant of digested rotten guava was used as a media of three concentrations for the culture of *Spirulina platensis*. The supernatant of 60% digested rotten guava showed maximum optical density on the 10th day of culture comparing with 20% and **60% or 40 %** supernatant of DRGM which has the similarity with the findings of **Habibet al. (2003, 2018), Satter (2017)**. From the above discussion, the growth performance of *Spirulina platensis* in supernatant of 60% DRGM was found better than 20% and 40% DRGM. It can be said that the physical properties, chemical parameters and technical facilities used in the present study were more or less similar to those used by other researchers. This study finds out a suitable concentration of digested rotten guava medium (DRGM) as an organic nutrient medium for culture and growth of *Spirulina platensis*.

The experiment revealed the proximate composition, growth performance of *Spirulina platensis* where the initial cell weight was 0.0023 mg/L which attained a maximum cell weight of 0.818 mg/L in 60% DRGM, 0.809 mg/L in 40% DRGM and 0.815 mg/L in 20% DRGM on the 10th day of the culture period. Similarly, the chlorophyll *a* content of inoculated *S. platensis* was 0.0015 mg/L which attained the highest content of 0.862 mg/L in 60% DRGM, 0.768 mg/L in 40% DRGM, 0.770 mg/L in 20% DRGM on the 10th day of culture period.

CONCLUSION

The percentage of crude lipids ($10.15 \pm 0.14\%$) of spirulina cultured in supernatant of 60% DRGM was significantly ($P < 0.05$) and almost two times higher than that of spirulina grown in 20% and 40%, respectively. The percentage of crude protein of *Spirulina* was 53.25 ± 0.32 , 53.35 ± 0.34 and $53.28 \pm 0.32\%$ when grown in the supernatant of 20, 40 and 60% DRG media, respectively. There was no significant ($P > 0.05$) variation among the crude protein percent of spirulina cultured in three different media of DRG. There was no significant ($P > 0.05$)

difference of crude lipids of spirulina when cultured in the supernatant of 20, 40 and 60% digested rotten guava. Ash (%) of *Spirulina* grown in supernatant of 20% ($10.22 \pm 0.13\%$), 40% ($10.16 \pm 0.17\%$) and 60% ($10.33 \pm 0.21\%$) digested rotten guava. There was no significant (**$P > 0.01$ or $P < 0.05$**) difference among the ash of *spirulina* grown in supernatant of 20, 40 and 60% DRGM. There was no significant variation among the NFEs of *Spirulina* grown in supernatant of 20, 40 and 60% DRG as the values found for 20% ($16.28 \pm 0.32\%$), 40% ($16.10 \pm 0.18\%$) and 60% ($15.99 \pm 0.22\%$), respectively. Specific growth rate (SGR) in respect to cell weight of spirulina grown in 60% digested rotten guava (DRG) medium was significantly ($P < 0.05$) higher than that of spirulina cultured in the supernatant of 20, 40%DRGM. This medium may be used commercially as the collection and preparation of these organic media require little cost, less labour and is available throughout Bangladesh. However, it might be suggested that more research and cost-benefit analysis have to be performed to evaluate the grow-out potential of spirulina in lab-based cultivation. The culture techniques may dwindle the cost of production and might be considered as a media for *Spirulinaplatensis* cultivation.

REFERENCES

- Bold, H.C. and Wynne, M.J., 1978: Introduction to the Algae. Structure and Reproduction.**Englewood Cliffs. New Jersey. pp. 706.
- Dineshkumar R, Narendran R, Sampathkumar P 2016: Cultivation of *Spirulinaplatensis* in different selective media. Indian Journal of Marine Science 45 (12) 1749-1754.**
- Habib MAB, Kohinoor AHNM 2018: Culture and production of house fry larvae and spirulina using poultry waste and their use as food for catfish post-larvae. Report on Advanced Research, Ministry of Education, Govt. of People Republic of Bangladesh. chapter-2 66-70.**
- Habib MAB, Yusoff FM, Phang SM, Mohamed S 2003: Growth and nutritional values of *Moinamicrurafed* on *Chlorella vulgaris* grown in digested palm oil mill effluent. Asian Fisheries Science 16 (1-2) 107-119.**
- Horwitz W (Editor) 1984: Official Methods of Analysis of the Association of Official Analytical Chemists. 14th Edition.**Association of Official Analytical Chemists, Washington DC. USA. pp. 1018.
- Kato T 1991: Chemistry of microalgae and their application to food. Food Chemistry 8 30-35.**
- Phang SM, Chu WL 1999: University of Malaya Algae Culture Collection (UMACC). Catalogue of Strain.**Institute of Postgraduate Studies and Research, University of Malaya, Kuala Lumpur, Malaysia.pp. 77.
- Phang SM, Miah MS, Chu WL, Hashim H 2000: Spirulina culture in digested sago starch factory waste water. Journal of Applied Phycology 12 395-400.**

- Sharker MGU 2002: Study of the culture of *Spirulina platensis* in various concentrations using papaya skin powder medium. MS. thesis submitted to the Department of Fisheries Management, Bangladesh Agricultural University, Mymensingh-2202. pp. 58.
- Satter A 2017: Culture and production of housefly larva and *Spirulina* using poultry waste, and their use as food for catfish post-larvae, PhD Thesis, Department of Aquaculture, Bangladesh Agricultural University, Mymensingh. pp. 143.
- Sukumaran P, Nulib R, Halimmon N, Simoh S, Omar H, Ismail A 2018: Formulation of cost-effective medium using urea as a nitrogen source for *Arthrospira platensis* cultivation under real environment. *Annual Research and Review in Biology*. 22(2) 1-12.
- Vonshak, A. & A. Richmond, 1988. Mass production of the blue green alga *Spirulina*: an overview. *Biomass* 15: 233–247.
- Vymazal J 1995: *Algae and Element Cycling in Wetlands*. CRC Press, Inc., Boca Raton, Florida, USA. pp. 689.
- Zar JH 1984: *Biostatistics*. Prentice-Hall Inc., Englewood Cliffs, New Jersey, USA. pp. 718.
- Zarrouk C 1996: Contribution à l'étude d'une cyanobactérie: influence de divers facteurs physiques et chimiques sur la croissance et la photosynthèse de *Spirulina maxima* (Setchell et Gardner) Geitler. PhD thesis, University of Paris, France. pp. 412.