

Immunoglobulin E (IgE) specific to cross reactive carbohydrate determinants (CCDs) in legume allergic cross reactivity and its relevance in clinical allergy

Abstract: A glycol-related IgE reactivity has been demonstrated in most allergen sources, especially in plant kingdom. Recent progress in glycobiology has allowed a clearer classification of these glycan-epitopes. Unlike classical peptide chain-based epitopes, glyco-epitopes can share significant structural homologies beyond the limits of protein families. These glycol epitopes are thus prone to extensive cross products as distinct as pollen and Hymenoptera venoms. Because of a monovalent IgE-binding is reactivity. Many of these glycoepitopes behave as “pan-epitopes” leading to cross reactivity between sufficient in serum-based assays, glycol epitopes and cross reactive carbohydrate (CCD) are classically considered as a potential source of positive *in vitro* results without clinical significance. But some reports recently demonstrated that glycol epitopes could induce a response at the cell level and suggested that they might play a role *in vivo*. So, accurate measurement and specificity assessment of CCD-specific IgE is a crucial and critical factor to understand the allergenic cross reactivity and clinical sensitivity mediated by CCD- specific IgE. Present review considers studies with *in vitro* allergenic cross reactivity due to IgE binding may not be a predictive measure of clinical symptoms and that is very true in case of haptenic or glycan IgE binding of antigens. The CCD and related IgE binding have to be assessed and evaluated for clinical relevance for avoidance by sensitized subject and also it is required for allergenic assessment before the functional proteins are labeled for allergenicity during commercialization of food products containing legume CCD-

glycoproteins. The legume allergenic cross reactivity and biological relevance of CCD-specific IgE will provide information for clinician to review its diagnostic role of IgE to identify allergenicity. It promotes further research to evaluate the clinical relevance of CCD IgE in defining true allergic cross reactivity among glycoproteins.

Key words: Immunoglobulin E (IgE); Food allergy; Cross reactive carbohydrate determinants CCD; glycoproteins, hypsensitive antigen, clinical allergy; mast cell/basophils; glycan-epitopes.


Abbreviations:

CCD: cross-reactive carbohydrate determinant; US: United States; EU: European Union; SPT: skin prick test; MMXF3: two mannose additional on N-glycans with β - 1,2 xylose and α -1,3 fucose residues; MUXF3: deletion of one mannose on N-glycans with β - 1,2 xylose and α -1,3 fucose residues; MMF3: two mannose additional on N-glycans with only α -1,3 fucose residues; MMF3F6: two mannose additional on N-glycans with α -1,3 fucose and α - 1,6 fucose residues deletion of β - 1,2 xylose residue; MMXF3F6: two mannose additional on N-glycans with β - 1,2 xylose, α - 1,3 fucose residues and α - 1,6 fucose residues; SEA: Schistosome soluble Egg Antigen; CAP-RAST: Pharmacia CAP -Radioallergosorbent test; HRP: Horseradish peroxidase; RBL: Rat Basophil Leukemia.

1. Introduction

Allergic diseases are responsible for such serious health problems as asthma, dermatitis, food allergy, bee sting allergy, conjunctivitis, and severe systemic anaphylaxis. [1, 2] These diseases affect a substantial portion of the general population. Asthma and atopic dermatitis, respectively, affect 10% and 15% of the children in some countries. A considerable portion of the health care expenses has been directed at the treatment of allergic diseases. Considerable efforts have been made for identifying factors contributing to these diseases [3, 4]. An allergic reaction is initiated when an allergen interacts with an IgE antibody. Many of the allergens are proteins and the prediction of these allergen proteins facilitate the study of molecular mechanisms and risk assessment of allergic reactions in different individuals. [4, 5] Computational methods have been developed for predicting allergen proteins from sequence segments that show identity, homology, or motif match to a known allergen. These methods achieve good prediction accuracies, but are less effective for novel proteins with no similarity to any known allergen. Common structural and physicochemical features that have been explored for predicting allergenicity from protein sequence. [6, 7] The US, EU, and many other countries have now implemented requirements for evaluating the potential allergenicity of proteins. The primary focus, either elaborated or implied, is on preventing the transfer of a known allergen, or a protein sufficiently similar to a known allergen that it may trigger allergic cross-reactions in food components. The severity of allergic reaction on different organs of human are showed in the (Figure.1)

Figure.1 Allergy Consequences:



2.0. Allergenicity and risk assessment:

An appropriate risk assessment strategy of most novel proteins and glycoproteins include serum IgE testing from allergic serum donors. [8] But based on only these in vitro immunochemical methods it is hard to differentiate between proteins that are likely to cause harm than that are likely to be safe. [9, 10] Because the primary focus for most immunological and bioinformatics approach used to assess the allergenicity of novel proteins will not rule out the non-specific false positive IgE binding with no relevance in clinical sensitivity and it is more likely to result in wrong interpretation in assessing the protein allergenicity. [11, 12] Since it is impossible to suspect based on biochemical assays, that the consumer with positive IgE reactivity to a novel protein might experience a clinically important allergenic reaction after consumption without any

biological assays. There are many reports in the literature which shows a significant *in vitro* IgE binding and evidences of cross reactivity with no clinical symptoms for a particular antigen. [13, 14] Even in assessing cross reactivity for an allergen the consideration of biological relevance in IgE binding assays is essential to predict the possibility of its potential cross reactivity with proteins or glycan chains of proteins of its relative classes which share more than 50% sequence homology.

3.0. Allergenic IgE cross reactivity

The introduction of novel proteins into foods carries a risk of eliciting allergic reactions in individuals sensitive to the introduced protein or of sensitizing susceptible individuals. IgE cross reactivity between a novel protein and a known allergen is an important issue in the evaluation of the allergenic potential of novel proteins. [15] To investigate the potential cross reactivity among proteins, specific IgE screening studies with serum from allergic subjects are generally performed. However, most immunochemical methods are nonfunctional assays and do not provide information on the ability of a protein to induce type I allergic reactions, as in case with *in vivo* tests. [11, 16] However, such *in vivo* tests, including the skin prick test (SPT) and intradermal testing have limitations in regard to evaluating allergenic potential of proteins in foods due to ethical concerns. In order to overcome these problems, *in vitro* cell based degranulation assays with measurement of histamine and beta hexosaminidase mediator was employed for assessment of allergenicity. These can serve as a tool along with other immunochemical based assays in determining the cross reactivity and potential allergenicity of some novel proteins in relation to clinical sensitivity. The functional *in vitro* bioassay using cell lines was compared with immunochemical and basophil histamine release assay methods used

for the analysis of allergen extracts and evaluated with respect to its use in the assessment of biological reactivity of allergenic proteins. [17, 18]

4.0. Legume allergenic cross reactivity:

Legumes are dicotyledonous plants belonging to the Fabales order. The main distinctive characteristic of which is their fruit legumen, seeds contained in pods. This botanical order is formed by three families: Mimosaceae, Caesalpinaceae and Papilionaceae or Fabaceae. The Papilionaceae family includes the most important allergenic species: *Lens culinaris* (lentil), *Cicer arietinum* (chick-pea), *Pisum sativum* (pea), *Arachis hipogea* (peanut), *Phaseolus vulgaris* (bean) and *Glycine max* (soy). Legumes are an important ingredient in the Mediterranean diet. Among children, sensitivity to legumes is the most prevalent dispute of food allergy. Peanut, lentil and chick-pea are the most frequent cause of allergic reactions to legumes. [19, 20] Legumes could be involved in severe allergic symptoms mentioned in Table 1. The different legumes have structurally homologous proteins, but they are not all equally allergenic, thus making it difficult to distinguish *in vitro* and *in vivo* cross-reactivity. [21, 22]

Peanut allergy can be associated with allergy to lentil, chick-pea and pea but less frequently. Contrarily, white bean and overall green bean and soy are well tolerated by children allergic to other legumes. In one study, 82% of the children allergic to legumes had a sensitization to pollen. Pea and bean are the legumes with more *in vitro* cross-reactivity with *Lolium perenne* (perennial ryegrass), *Olea europea* (olive) and *Betula Alba* (birch). [22] This cross-reactivity occurred because of common antigenic determinants or due to the coexistence of pollen and legume allergy. Pan-allergens implication seems to be less probable. [23] It is important to emphasize that in spite of an evident clinical and immunological cross-reactivity,

diagnosis of legume allergy should not be based only on specific IgE tests. The decision to eliminate one legume from the diet should be based on a positive oral food challenge. The peanut, *Arachis hypogea*, is a member of legume family and is related to beans and peas, immediate hypersensitivity reactions to peanut are much more prevalent and severe than other legumes and tend to persist into adulthood. [21] The peanut allergic subjects are known to show in vitro IgE binding to several other legume extracts, which are claim to have cross reactivity. Since many of protein IgE reacting elements are glycoproteins and most of the legumes contain glycoproteins and suspected to cause legume cross-reactivity. CCD-specific IgE is the key factor in recognizing the multiple IgE binding for other legume glycoproteins. [24] But, clear evidence of demonstrating and evidencing possible clinical reactivity of CCD-IgE in legume cross reactivity is lacking in literature. List of some well-known legume allergens were listed below.

(Table.1)

Table.1. Some of well-known Legume allergens list


Sl. no	Legume source	Legume allergen	Molecular weight	Protein family	Protein super-family	References
1	Arachis hypogea	Ara h1	64	7S globulin- vicilin	Cupin	[50]
	Peanut	Ara h2	17	2S albumin- conglutin	Prolamin	[51]
		Ara h3	60	11S globulin- legumins	Cupin	[52]
		Ara h4	37	11S globulin- legumins	Cupin	[53]
		Ara h5	15	Bet v2 -Profilin	Profilin	[54]
		Ara h6	15	2S albumin- conglutin	Prolamin	[55]
		Ara h7	15	2S albumin- conglutin	Prolamin	[53]
		Ara h8	17	Bet v1-family	PR-proteins	[56]
		Ara h9	9.8	Non-specific lipid transfer protein1	Prolamin	[57]
		Ara h10	16	Oleosin- lipid storage bodies	Oleosin	[58]
		Ara h11	14	Oleosin- lipid storage bodies	Oleosin	[59]
2	Glycine max	Gly m1	7	Non-specific lipid transfer protein	Prolamin	[60]
	Soybean	Gly m2	8	Defensin	Scorpion toxin- Knottin	[61]
		Gly m3	14	Profilin	Profilin	[62]
		Gly m4	17	Bet v1-family	Bet v1	[63]
		Gly m Bd28k	28	7S globulin- vicilin	Cupin	[64]
		Gly m Bd30k	34	Thiol protease- papain	Cupin	[64]
		Gly m Lectin	14.5	Lectin-SBA agglutinin	Prolamin	[56]
		Gly m Bd60k	63-67	7S globulin- vicilin	Cupin	[65]
		Gly m glycinin G1	40	11S seed storage protein- Glycinin	Glyinin	[66]
		Gly m glycinin G2	22	11S seed storage protein- Glycinin	Glyinin	[66]
		Gly m glycinin G4	61.2	11S seed storage protein- Glycinin	Glyinin	[66]
Gly m T1	20	Kunitz- trypsin inhibitor	-	[59]		
3	Pisum sativum	Pis s1	44	7S globulin- vicilin	Cupin	[67]
	Pea	Pis s2	63	7S globulin- vicilin	Cupin	[67]
4	Cicer arietinum	Cic a 2S Albumin	10-12	2S albumin-Albumin prolamin	Prolamin	[68]
	Chickpea	Cic a 11S Globulin	46.5 and 39.8	11S Glycinin-legumin type	Cupin	[69]
5	Lens culinaris	Len c1	48	Vicilin- Gamma subunit	Cupin	[70]

6	Lentil	Len c2	66	Biotinylated protein- seed specific	-	[70]
		Len c3	9	Non-specific lipid transfer protein1	Prolamin	[71]
7	Lupinus angustifolius	Lup an1	55-61	Beta conglutin-vicilin	Cupin	[72]
	Lupin	Lup a vicilin	66	Vicilin	Cupin	[73]
		Lup a 11S globulin	20	11S globulin	Cupin	[74]
8	Phaseolus vulgaris	pha v3	8.8-9	Non-specific lipid transfer protein1	-	[75]
	Green bean	pha v3	8.8-9	Non-specific lipid transfer protein	Prolamin	[76]
9	Blackgram	Vig m	28	Homology- Rho specific inhibitor	-	[77]

5.0. Leguminous Allergen properties:

Not one single Characteristic alone is sufficient to recognize a certain protein as allergenic. Especial Characteristics like physical-chemical and immunological properties to identify allergenic proteins which are capable of eliciting allergenic responses in Atopic people. Based on the sensitization pattern IgE-mediated food allergy are classified into two categories namely A) Class-1 food allergens: Stable in gastric fluid (sensitization occurs via gastrointestinal tract). [25] B) Class-2 food allergens: Heat sensitive and susceptible to gastric-digestion (sensitization does not exist via oral tract). They promote allergenic responses by means of IgE-cross reactivity in already sensitized people. Example inhalant allergens like pollen-associated food allergies. [26] Physical-chemical and immunological properties of allergen based on the characteristic of abundance, stability, tertiary structure, modification and glycosylation characteristics of the protein are essential to decide a protein as allergen. Most of the allergens are glycoprotein with of molecular weight 10-70 kDa, with greater stability over acid, heat as well as proteases treatment and water-soluble. Majority of food allergens are stable to pepsin digestion (8-60min) and survive in gastrointestinal tract but the non -allergenic food-proteins are get easily digested

in gastrointestinal tract. [27, 28] Protein glycosylation is an additional criterion in the enhancement of allergenicity of a protein, one such Example- major allergens from legume family namely Ara h1, Gly m Bd28K and Gly m Bd60K are extremely glycosylated. Previous reports further indicated the glycan moiety can also induce allergenicity by cross-linking with IgE. Immunological characteristics not only incorporate binding of IgE but also include awareness about B and T-cell epitopes. Epitopes with greater IgE-binding ability on allergens are accountable for provoking allergenic reactions. These IgE cross-linking epitopes are occur in two forms namely linear or conformational form. During the digestion of legume family, leguminous allergens lose their conformational form and appear as linear epitopes and sensitize individuals in gastrointestinal tract. [29] Some of the physico-chemical and immunological properties of allergen were showed. (Figure.2)




6.0. CCD a cross reactive antigen determinant

The N-linked carbohydrate groups of glycoproteins induce IgE, leading to cross reactivity between several food and pollen allergens. [30, 31] These are called as cross reactive carbohydrate determinants (CCDs) which are present on glycoproteins of vegetables such as peanut, celery, tomato and potato. IgE binding to CCDs have been described in literature for various allergenic extracts and there are several research studies have also been performed to understand and verify presence as well as biological function of the CCD-IgE. [32, 33] Glycosylation has been described for many allergenic and non-allergenic molecules. The glycan structures have been investigated and a heterogeneous IgE reactivity has been reported. [34] *In vitro* demonstration of the glycan allergenic activity has been reported only for a native glycoprotein derivative and for synthetic neo-glycoproteins, whereas no *in vivo* activity of any purified allergenic glycoproteins has ever been reported. From an allergological prospective, most important feature of CCD is the wide distribution among plant-derived proteins, though similar structures can also be present in non-plant glycoproteins. Monoclonal and polyclonal antibodies to CCD have been used to define reactive carbohydrate epitope distribution, whereas limited data are available with on the prevalence and distribution of IgE reactive with CCD. [35] Some of the well-evidenced CCD-glycoprotein epitopes are short-listed from plants, insects, parasites and non-human mammalian oligosaccharides are: MMXF3, MUXF3, MMF3, MMF3F6, MMXF3F6 and alpha-gal. (Shown in figure.3) MMXF3 (MM- indicate the presence of tri-mannose residues)-Horse radish peroxidase -HRP model glycoprotein frequently employed for the study of carbohydrate -based IgE cross-reactivity , MUXF3- (MU-indicates deletion of one Mannose residue) Bromelian commonly used research glycoproteins, these are equally common in pollens, plant foods and rubber latex allergens, MMF3F6,MMF3,MUF3F6(indicates deletion of xylose and addition α 1,6-fucose residue)-yellow jacket venom (Ves v2) [36] honey bee, vespidae and hymenoptera venom [37] MMXF3F6 (indicates addition of α 1,6-fucose residue along with xylose and fucose) - Parasite-Schistome and Schistome soluble Egg Antigen (SEA), Alpha-gal- Non-human mammalian oligosaccharides. These are randomly picked up to rule out leading task of α -1,3 fucose residue in the target of IgE antibody in allergenicity. [38] Preliminary researchers disclosed Xylose as


core element; ulteriorly studies on xylose summarized no IgE or intensively less antibody binding competence observed. [39] These days α -1, 3 fucose surveyed all-important for IgE-reactivity at standpoint of allergy. Structural comparison of plant, insect, parasite and non-human mammalian glycoproteins was mentioned. (Figure.3)

Figure.3 some of well published CCD structures from different sources

A: Capital Nub of N-glycans




B: Core structures of N-linkage and O-linkage




C: Some of Well-known structures of Oligosaccharides that are accepted as Targets of IgE :


a) Horse radish peroxidase (HRP)




b) Bromelain (Brom)




c) Bee Venom



c) Schistomes and Schistome soluble egg Antigen (SEA)



d) Non-Human Mammalian oligosaccharides



D: Monosaccharides present in plant and human Oligosaccharides

- Mannose (Mn)
- N-acetyl glucosamine (GlcNAc)
- Xylose (Xyl)
- Fucose (Fuc)
- Galactose-alpha-1,3 galactose

7.0. CCD-IgE and clinical allergy

The clinical relevance of IgE to cross reactive carbohydrate determinants (CCDs) is still controversial, because of the widespread occurrence of structurally similar CCDs in plants. IgE to carbohydrates often leads to false positive diagnostic results in CAP-RAST, and immunoblot analysis, when only IgE binding to extracts or allergens in relation to the initiated symptoms is considered.[12] Many glycoproteins like bromelain, which carry only one IgE-binding glycan, are unable to cross-link IgE bound receptors of mast cells and basophils, and clinical symptoms might not appear in patients whose IgE response is restricted to CCDs. [11, 23] On the basis of this view several authors regard anti-CCD IgE as specific antibodies without clinical relevance. However there are also many glycoproteins with more than one N-linked glycan, for example; HRP, beta fructofuranoside, polygalacturonase and pectin esterase are potential plant allergens, have got the capability of cross-linking adjacent IgE, on mast cells and basophils resulting in clinical symptoms. [40, 41] Until now, studies to prove whether natural multivalent glycoproteins might be able to induce histamine release or not and thus might also contribute to clinical symptoms of allergic patients are very rare. Most likely due to difficulties in the experimental design, lack of suitable purified multivalent allergenic glycoproteins failed to report their effective role in inducing false positive results. Many reports describe the wide CCD-IgE binding to clinically irrelevant allergenic extracts and defined degree of interference of CCD-IgE in the detection. The presence of CCD-IgE directly correlates with positive assay for several

different allergenic extracts that are unable to trigger an allergic reaction in the skin prick test. [42, 14] However in some studies demonstrated that CCDs are the essential part of IgE epitope in some glycoproteins and show that glycoproteins carrying multiple glycan units can be biologically active in patients sensitized to CCD. [42,43] Plant derived glycoproteins are indeed the most reacting ones, but the higher the concentration of CCD-IgE are more probable chance to detect IgE for any SPT-negative allergenic extract. Moreover, the differences recorded in the sera reactivity, and the presence of sera with CCD-IgE but not reacting with certain extracts, suggest a possible heterogeneity of the CCD-IgE. The use of purified allergenic glycoproteins in comparative studies would certainly increase the information on the specificities of CCD-IgE. From an immunochemical point of view, extended inhibition studies are needed to confirm the statistical correlation of CCD-IgE reactivity. [44]

8.0. CCD glycan epitopes and IgE binding:

The asparagine-linked carbohydrate moieties of plant glycoproteins are the most abundant environmental immuno determinants. They are the structural basis of CCDs despite some structural variation. [33] Two main motifs are xylose and the core 3-linked fucose, which form the essential part of two independent epitopes. Plants contain both epitopes whereas insect glycoproteins contain only fucose [45, 46] about 20% or more of allergic patients generate specific anti-glycan IgE, which is often accompanied by IgG. The clinical relevance of IgE specific for glycan (CCD) has been a matter of controversy. Until now, no convincing experiments have been performed to test the biologic significance of individual multivalent allergens that carry multiple carbohydrate epitopes. [11, 12] We sought to contribute understanding the role of CCD-specific IgE antibodies and to study whether CCD-specific IgE


antibodies are able to activate basophils using different multivalent glycoproteins as allergens. Most glycoproteins in legumes are known to present in multimeric forms and if each subunit contain a single glycan chain, and as multimer it provides more than one interacting glycan chain on IgE recognition which can facilitate the trigger of degranulation. So the addressing of CCD-specific IgE against legume glycoproteins is important to predict the possible cross-reactivity among legume species.

9.0. Legume allergenic cross reactivity, CCD-IgE and its clinical relevance

Legumes are major protein diet of Indian subcontinent. All parts of India consume high amount legumes through various types of foods. Allergic reactions to legumes were reported at different regions of India. Recently some studies identify few allergens from major legume species.[47] Though there is large number clinical report of avoidance of legume by sensitized population a thorough study on legume allergy is lacking in India. Significant cross reactivity among legumes is natural because of the presence of structurally homologous proteins that share common epitopes. This has been demonstrated and observed to have pathological symptoms. [48] In majority of cases, patients were found to be sensitive to more than one legume. Previous reports have demonstrated a great degree of immunological cross-reactivity among legumes and reported that prediction of immunological cross-reactivity should not be based only on allergen-specific IgE antibody determination. [47, 48] However, some reports have been published on cross-reactivity between certain members of leguminosae family due to N-glycosylation. Peanut allergy can also be associated with allergy to lentil, chickpea and pea, but less frequently reported. Lentil, chickpea and pea are widely consumed in Mediterranean area and significant cross-reactivity exists among them whereas low clinical significance of legume cross reactions

reported in earlier reports. A glycan-related IgE reactivity has been demonstrated in most allergen sources, especially in plant kingdom. Recent progress in glycobiology has allowed a clearer classification of these glycon-epitopes. Unlike classical peptide chain-based epitopes, glyco-epitopes can share significant structural homologies beyond the limits of protein families. These glycoepitopes are thus prone to extensive cross reactivity. [48, 49] So, accurate measurement and specificity assessment of CCD-specific IgE is a crucial and critical factor to understand the allergenic cross reactivity and clinical sensitivity mediated by CCD-IgE. The work proposed in this study will be useful to evaluate and assess the possible involvement of CCD-IgE in cross reactivity and clinical sensitivity as they are present in novel and recombinant glycoproteins. We claim that several studies have demonstrated and proven the presence of specific IgE to glycan portion of many dietary glycoproteins. Their clinical relevance is not fully understood yet and are not much reports in the literature to show their possible involvement in clinical cross reactivity except a few preliminary reports. Accurate measurement and specificity assessment of CCD-specific IgE is a crucial and critical factor to understand the allergenic cross reactivity and clinical sensitivity mediated by CCD-IgE. In India, the studies relevant to food allergy were very limited and most focus has been given to pollen and dust allergy compare to food induced hypersensitivity. Presently few groups are working on characterization of legume allergy and few well established reports were published with *in vitro* legume cross reactivity with very low clinical relevance. Many new legume allergens were characterized and reported which were different from the western world. There is variation in the sensitization and tolerance among Indian population to legume diet which ear mark with history of diet and genetic adaptations. The current review aims in understanding the CCD IgE in clinical relevancy which

is more responsible for observed in vitro IgE cross reactivity among similar legume allergens.(Figure.4) shows the overview of allergy occurrence to legume CCD factors.



10. Allergenic risk associated with CCD-IgE

The evaluation of allergenic potential is a key parameter in the safety assessment of novel proteins, including those expressed and used as functional proteins. The majority of allergic reactions to food proteins are immediate type food hypersensitivity reactions in which the

principal biological effector is IgE antibody. [78] The presences of IgE to CCD is well known in many plant extracts and are known to be common in allergic population and are binding to the skin-prick test negative allergenic extracts further state its poor biological activity and clinical sensitivity. [79] CCD-IgE can be recorded for wide range of allergenic unrelated extracts. Allergenic sera to be used in any *in vitro* assay employing native or recombinant glycoproteins should be carefully evaluated for the presence of CCD-IgE to assess its *in vivo* allergenic potential and cross reactivity. The accurate measurement and specificity assessment of CCD-specific IgE is a crucial and critical factor to understand the allergenic cross reactivity and clinical sensitivity mediated by CCD-IgE. [80] So it induced degranulation *in vitro*. This method is also being useful to evaluate and assess the possible involvement of CCD-IgE in cross reactivity and clinical sensitivity as they are present in novel and recombinant glycoproteins. The rat basophilic leukemia (RBL) cells expressed with human FcεRI which is selectively bind to human serum IgE antibodies and can be stimulated by cross-linking with protein allergens to degranulate, releasing granules containing various mediators. This method has been utilized extensively to study the mechanisms of IgE interactions with mast cell FcεR and for the measurement of degranulation induced allergenic proteins mediated by IgE antibodies. The present proposal stands on the prediction of the possibility of IgE to cross reactive carbohydrate epitope and assessment of their specificity in relation to sensitivity resulting in clinical symptoms. Most glycoproteins shows IgE elicitation to its glycan portion that was reported to have poor biological activity but still gives false prediction in allergic assessment using *in vitro* diagnostic methods. [81] The study of specificity and the design of evaluation methods to predict the CCD-specific IgE are essential. In other hand the question of its clinical relevance in terms of its ability to elicit of symptoms and cross-linking of IgE on mast cell to induce

degranulation. The present proposal is focused on the detailed study of CCD-IgE specificity to legume glycoproteins and prediction and assessment of the possible cross reactivity among legumes due to CCD-IgE and evaluation of the clinical significance of CCD-IgE by designing some *in vitro* and *in vivo* assessment methods should be incorporated.

10.1. Conclusion:

We claim that several studies have demonstrated and proven the presence of specific IgE to glycan portion of many dietary glycoproteins. Their clinical relevance is not fully understood yet and no many reports were in the literature to show their possible involvement in clinical cross reactivity except a few preliminary reports. Accurate measurement and specificity assessment of CCD-specific IgE is a crucial and critical factor to understand the allergenic cross reactivity and clinical sensitivity mediated by CCD-IgE. So the work proposed in this study will be useful to understand, evaluate and assess the possible involvement of CCD-IgE in cross reactivity and clinical sensitivity as they are present in novel and recombinant glycoproteins. The use of basophilic degranulation study using rat basophilic leukemia (RBL) cells will set a standard method to assess the possible clinical cross reactivity among the closely related allergenic glycoproteins containing CCD-specific IgE in sensitized population. This proposal also aims at the elucidation of possible molecular mechanism of CCD induced degranulation of basophils/mast cells and also evaluates the potential of CCD-specific IgE to induce clinical symptoms as a function of sensitization to CCD and recognition of its cross reactive glycoproteins of related families.

11. References:

1. Hannah J. Gould, Brian J. Sutton, Andrew J. Beavil, Rebecca L. Beavil, Natalie McCloskey, Heather A. Coker, et al. The Biology of IgE and the basis of allergic disease. *Annu. Rev. Immunol.* 2003; 21:579–628.
2. Hugh A. Sampson. Update on food allergy. *J Allergy Clin Immunol.*2004; 113:805-819.
3. Bruce S. Bochner, MD, William W. Busse, MD Baltimore, M and Madison, Wis. Advances in mechanisms of allergy. *J Allergy Clin Immunol.*2004; 113:5:868–875.
4. Heimo Breiteneder, E.N. Clare Mills. Plant food allergens- structural and functional aspects of allergenicity. *Biotechnology.Advances.*2005; 23:395–399.
5. H. Malandain The clinical use of specific IgE antibody testing – a further word of caution. 2004; 59:3:359–360.
6. Ivanciuc, Ovidiu, Mathura, Venkatarajan, Midoro-Horiuti, Terumi; Braun, et al. Detecting Potential IgE-Reactive Sites on Food Proteins Using a Sequence and Structure Database, SDAP-Food. *Journal of Agricultural and Food Chemistry.* 2003; 51:16: 4830–4837.
7. Gijs A Kleter, Ad ACM Peijnenburg. Screening of transgenic proteins expressed in transgenic food crops for the presence of short amino acid sequences identical to potential, IgE – binding linear epitopes of allergens. *BMC Structural Biology.*2002; 2:1: 8–0.
8. Goodman, Richard E. Hefle, Susan L. Taylor, Steven L. van Ree, Ronald. Assessing Genetically Modified Crops to Minimize the Risk of Increased Food Allergy: A Review. *Int Arch Allergy Immunol.* 2005; 137:2:153–166.
9. Mari Adriano, Iacovacci Patrizia, Afferni Claudia, Barletta Bianca, Tinghino Raffaella, Di Felice Gabriella et al. Specific IgE to cross-reactive carbohydrate determinants strongly affect the in vitro diagnosis of allergic diseases. *J Allergy Clin Immunol.*1999; 103:6:1005–1011.
10. D. G. Ebo, M. M. Hagendorens, C. H. Bridts, L. S. De clerck, W. J. Stevens. Sensitization to cross-reactive carbohydrate determinants and the ubiquitous protein profilin: mimickers of allergy. *Clin Exp Allergy.*2004. 34:1:137–1.

11. Van der Veen, Mauritis J, van Ree, Ronald, Aalberse, Rob C et al. Poor biologic activity of cross-reactive IgE directed to carbohydrate determinants of glycoproteins. *J Allergy Clin Immunol.*1997.100; 3:327–334.
12. Foetisch Kay, Westphal Sandra, Lauer Iris, Retzek Mechthild, Altmann Fritz, Kolarich Daniel et al.. Biological activity of IgE specific for cross-reactive carbohydrate determinants. *J Allergy Clin Immunol.*2003; 111:4:889–896.
13. P. Iacovacci, C. Pini, C. Afferni, B. Barletta, R. Tinghino, E. Schininà, et al. A monoclonal antibody specific for a carbohydrate epitope recognizes an IgE-binding determinant shared by taxonomically unrelated allergenic pollens. *Clinical and Experimental Allergy.*2001; 31:3:458–465.
14. V. Mahler, C. Gutgesell, R. Valenta, T. Fuchs. Natural rubber latex and hymenoptera venoms share Immunoglobulin E-epitopes accounting for cross-reactive carbohydrate determinants. *Clinical and Experimental Allergy.*2006; 36:11:1446–1456.
15. Goodman Richard E, Hefle Sue L. Gaining perspective on the allergenicity assessment of genetically modified food crops. *Expert Rev. Clin. Immunol.*2005; 1:4:561–578.
16. Van Ree Ronald, Aalberse Rob C. Specific IgE without clinical allergy. *J Allergy Clin Immunol.* 1999; 103:6:1000–1001.
17. Aronica Mark A, Vogel Nicola. Pathogens and immunologic memory in asthma: what have we learned? *Expert Rev. Clin. Immunol.* 2005; 1:4:589–601.
18. S. Kaul, D. Lüttkopf, B. Kastner, L. Vogel, G. Höltz, S. Vieths, A. Hoffmann. Mediator release assays based on human or murine immunoglobulin E in allergen standardization. *Clin Exp Allergy.* 2007; 37:1:141–150.
19. M.J. Pereira, M.T. Belver, C.Y. Pascual, M. Martín Esteban. The allergenic significance of legumes. *Allergol ET Immunopathol.*2002. 30; 6:346–353.
20. Martinez San Ireneo M, Ibanez Sandin MD, Fernandez-Caldas E. Hypersensitivity to members of the botanical order Fabales (legumes). *J Investig Allergol Clin Immunol.*2000; 10:4:187-199.
21. Sicherer Scott H., Sampson Hugh A. Peanut allergy: Emerging concepts and approaches for an apparent epidemic. *J Allergy Clin Immunol.*2007; 120:3:491–503.
22. Ibanez M.D., Martinez M, Sanchez JJ., Fernandez-Caldas E. Legume cross-reactivity. *Allergol et Immunopathol.*2003; 31:3:151-161.

23. Stefan Vieths., Stephan Scheurer., Barbara Ballmer-Weber. Current Understanding of Cross-Reactivity of Food Allergens and Pollen. *Annals of the New York Academy of Sciences*.2002; 964:1:47–68.
24. Van Der Veen, Van Neerven, De Jong, Aalberse. Jansen, Van Der Zee. The late asthmatic response is associated with baseline allergen-specific proliferative responsiveness of peripheral T lymphocytes in vitro and serum interleukin-5. *Clin Exp Allergy*.1999; 29:2:217–227.
25. Astwood J.D., Leach J.N., Fuchs R.L. Stability of food allergens to digestion in vitro. *Nature Biotechnology*. 1996; 14:10:1269–73.
26. Egger M, Mutschlechner S, Wopfner N, Gadermaier G, Briza P, Ferreira F. Pollen-food syndromes associated with weed pollinosis: an update from the molecular point of view. *Allergy*. 2006; 61:4:461–76.
27. Thomas K, Aalbers M, Bannon G.A, Bartels M, Dearman R.J, Esdaile D.J et al. A multi-laboratory evaluation of a common in vitro pepsin digestion assay protocol used in assessing the safety of novel proteins. *Regul Toxicol Pharmacol*. 2004; 39:2:87–98.
28. Bannon G.A., Goodman R.E., Leach J.N., Rice E., Fuchs R.L., Astwood J.D. Digestive stability in the context of assessing the potential allergenicity of food proteins. *Comments Toxicol*. 2002; 8:3:271–285.
29. Bannon G.A., Ogawa T. Evaluation of available IgE binding epitope data and its utility in bioinformatics. *Mol Nutr Food Res*. 2006; 50:7:638–644.
30. Jin Chunsheng, Hantusch Brigitte, Hemmer Wolfgang, Stadlmann Johannes, Altmann Friedrich. Affinity of IgE and IgG against cross-reactive carbohydrate determinants on plant and insect glycoproteins. *J Allergy Clin Immunol*.2008; 121:1: 185–190.
31. Kay Foetisch, Dae Son, Friedrich Altmann, Holger Aulepp, Amedeo Conti, Dieter Hausteine et al. Tomato (*Lycopersicon esculentum*) allergens in pollen-allergic patients. *Eur Food Res Technol*. 2001; 213:4:5:259–266.
32. Friedrich Altmann. The Role of Protein Glycosylation in Allergy. *Int Arch Allergy Immunol*. 2007; 142:2:99–115.

33. Malandain H. IgE-reactive carbohydrate epitopes—classification, cross-reactivity, and clinical impact. *European Annals of Allergy and Clinical Immunology*. 2005; 37:4:122-128.
34. Adriano Mari. IgE to Cross-Reactive Carbohydrate Determinants: Analysis of the Distribution and Appraisal of the in vivo and in vitro Reactivity. *Int Arch Allergy Immunol*. 2002; 129:4:286–295.
35. P. Iacovacci, C. Pini, C. Afferni, B. Barletta, R. Tinghino, E. Schinina et al. A monoclonal antibody specific for a carbohydrate epitope recognizes an IgE-binding determinant shared by taxonomically unrelated allergenic pollens. *Clin Exp Allergy*. 2001; 31:3:458–465.
36. Lommerse JP, Kroon-Batenburg LM, Kamerling JP, Vliegenthart JF. Conformational analysis of xylose-containing N-glycan of pineapple stems bromelain as part of the intact glycoprotein. *Biochemistry Journal*. 1995; 34:25: 8196-8206.
37. Aalberse RC, Koshte V, Clemens JG. Immunoglobulin E antibodies with vegetable foods, pollens and Hymenoptera venom. *J Allergy Clin Immunol*. 1981; 68:5:356-64.
38. Kolarich D, Leonard R, Hemmer W, Altmann F. The N-glycans of yellow jacket venom hyaluronidases 21 and the protein sequence of its major isoform in *Vespula vulgaris*. *FEBS Journal*. 2005; 272:20:5182-5190.
39. Lauriere C, Lauriere M, Sturm A, Faye L, Chrispeels MJ. Characterization of β -fructosidase, an extracellular glycoprotein of carrot cells. *International Journal of Biochemistry and Molecular Biology (Biochimie)*. 1988; 70:11:1483-1491.
40. Sandra Westphal., Daniel Kolarich., Kay Foetisch., Iris Lauer., Friedrich Altmann., Amedeo Conti et al. Molecular characterization and allergenic activity of Lyc e 2 (β -fructofuranosidase), a glycosylated allergen of tomato. *Eur. J. Biochem*. 2003; 270:6:1327–1337.
41. B. Fahlbusch., W.D. Müller., O. Rudeschko., L. Jager., O. Cromwell., H. Fiebig. Detection and quantification of group 4 allergens in grass pollen extracts using monoclonal antibodies. *Clin Exp Allergy*. 1998; 28:7:799–807.
42. K. Fotisch., F. Altmann., D. Haustein. S. Vieths. Involvement of Carbohydrate Epitopes in the IgE Response of Celery-Allergic Patients. *Int Arch Allergy Immunol*. 1999; 120:1:30–42.

43. K. Fotisch., J. Fah., B. Wuthrich., F. Altmann., D. Hausteiner. S. Vieths. IgE antibodies specific for carbohydrates in a patient allergic to gum Arabic (Acacia Senegal). *Allergy*.1998; 53:11: 1043–1051.
44. Merima Bublin, Christian Radauer, Iain B.H. Wilson, Dietrich Kraft., Otto Scheiner, Heimo Breiteneder et al.. Cross-reactive N-glycans of Api g 5, a high molecular weight glycoprotein allergen from celery, are required for immunoglobulin E binding and activation of effector cells from allergic patients. *The FASEB Journal*. 2003; 7:0:1697-1699.
45. Verena Tretter., Friedrich Altmann., Viktoria Kubelka., Leopold Marz., Wolf M. Becker. Fucose α 1, 3-Linked to the Core Region of Glycoprotein N-Glycans Creates an Important Epitope for IgE from Honeybee Venom Allergic Individuals. *Int Arch Allergy Immunol* .1993; 102:3:259–266.
46. Malandain H. Allergies associated with both food and pollen. *European Annals of Allergy and Clinical Immunology*.2003; 35:7:253-256.
47. Rinkesh Kumar Gupta, Kriti Gupta., Akanksha Sharma., Mukul Das., Irfan Ahmad Ansari., and Premendra D Dwivedi. Look Insight: Health Risks and Benefits of Chickpea (*Cicer arietinum*) Consumption. *J. Agric. Food. Chem*.2017; 65:1:6-22.
48. Alok Kumar Verma., Sandeep Kumar., Mukul Das., Premendra D Dwivedi. A Comprehensive Review of Legume Allergy. *Clinic Rev Allerg Immunol*. 2013; 45:1:30–46.
49. Alok Kumar Verma., Sandeep Kumar., Mukul Das., Premendra D. Dwivedi. Impact of Thermal Processing on Legume Allergens. *Plant Foods Hum Nutr*.2012; 67:4:430–441.
50. Burks AW, LW Williams, R. M. Helm, W Thresher, JR Brooks, and HA Sampson. Identification of soy protein allergens in patients with atopic dermatitis and positive soy challenges: determination of change in allergenicity after heating or enzyme digestion. *Advances in experimental medicine and biology*. 1991; 289:0:295-307.
51. Burks A. W., L. W. Williams, W. Thresher, C. Connaughton, G. Cockrell, and R. M. Helm. Allergenicity of peanut and soybean extracts altered by chemical or thermal denaturation in patients with atopic dermatitis and positive food challenges. *J Allergy Clin Immunol*. 1992; 90:0:889-97.

52. Jin T, Guo F, Chen YW, Howard A, Zhang YZ. Crystal structure of Ara h 3, a major allergen in peanut. *J. Molecular Immunol.* 2009; 46:8:9:1796–804.
53. Kleber-Janke T, Cramer R, Appenzeller U, Schlaak M, Becker WM. Selective cloning of peanut allergens, including profilin and 2S albumins, by phage display technology. *Int Arch Allergy Immunol.* 1999; 119:265–274.
54. Kleber-Janke T., Cramer R., Scheurer S., Vieths S, Becker WM. Patient-tailored cloning of allergens by phage display: peanut (*Arachis hypogaea*) profilin, a food allergen derived from a rare mRNA. *J Chromatogr B: Biomed Sci Appl.* 2001; 756:1:2:295–305.
55. Van Wijk F., Hartgring S., Koppelman S.J., Pieters R., Knippels LM. Mixed antibody and T cell responses to peanut and the peanut allergens Ara h 1, Ara h 2, Ara h 3 and Ara h 6 in an oral sensitization model. *Clin Exp Allergy.* 2004; 34:9:1422–8.
56. Mittag D., Akkerdaas J., Ballmer-Weber B.K., Vogel L., Wensing M., Becker W.M et al. Ara h 8, a Bet v 1-homologous allergen from peanut, is a major allergen in patients with combined birch pollen and peanut allergy. *J Allergy Clin Immunol.* 2004; 114:6:1410–7.
57. Asero R, Mistrello G, Roncarolo D, de Vries S.C., and Gautier M.F., Ciurana C.L et al. Lipid transfer protein: a pan-allergen in plant-derived foods that is highly resistant to pepsin digestion. *Int Arch Allergy Immunol.* 2000; 122:1:20–32.
58. Pons L., Chery C., Romano A., Namour F., Artesani M.C et al. The 18 kDa peanut oleosin is a candidate allergen for IgE mediated reactions to peanuts. *Allergy.* 2002; 57: 88–93.
59. Burks A.W., Cockrell G., Connaughton C., Guin J., Allen W., Helm R.M. Identification of peanut agglutinin and soybean trypsin inhibitor as minor legume allergens. *Int Arch Allergy Immunol.* 1994; 105:2:143–9.
60. Helm R., Cockrell G., Herman E., Burks A., Sampson H., Bannon G. Cellular and molecular characterization of a major soybean allergen. *Int Arch Allergy Immunol.* 1998; 117:29–37.
61. Wilson S., Blaschek K., de Mejia E. Allergenic proteins in soybean: processing and reduction of P34 allergenicity. *Nutr Rev.* 2005; 63:2:47–58.

62. Ogawa A., Samoto M., Takahashi K. Soybean allergens and hypoallergenic soybean products. *J Nutr Sci Vitaminol (Tokyo)*. 2000; 46:6:271–9.
63. Renaud C., Cardiet C., Dupont C. Allergy to soy lecithin in a child. *Journal of pediatric gastroenterology and nutrition*. 1996; 22:3:328–9.
64. Gonzalez R., Polo F., Zapatero L., Caravaca F., Carreira J. Purification and characterization of major inhalant allergens from soybean hulls. *Clin Exp Allergy*. 1992; 22:8:748–55.
65. Codina R., Lockey R.F., Fernandez-Caldas E., Rama R. Identification of the soybean hull allergens responsible for the Barcelona asthma outbreaks. *Int Arch Allergy Immunol*. 1999; 119:1:69–71.
66. Natarajan S.S., Xu C., Bae H., Caperna T.J., Garrett W.M. Characterization of storage proteins in wild (*Glycine soja*) and cultivated (*Glycine max*) soybean seeds using proteomic analysis. *J Agric Food Chem*. 2006; 54:8:3114–20.
67. Sanchez-Monge R., Lopez-Torrejón G., Pascual C.Y., Varela J., Martin-Esteban M., Salcedo G. Vicilin and convicilin are potential major allergens from pea. *Clin Exp Allergy*. 2004; 34:11:1747–53.
68. Vioque J., Sanchez-Vioque R., Clemente A., Pedroche J., Bautista J., Millan F. Purification and partial characterization of chickpea 2S albumin. *J Agric Food Chem*. 1999; 47:4:1405–9.
69. Karamloo F., Wangorsch A., Kasahara H., Davin L.B., Hausteine D., Lewis N.G et al. Phenylcoumaran benzylic ether and isoflavonoid reductases are a new class of cross-reactive allergens in birch pollen, fruits and vegetables. *Eur J Biochem*. 2001; 268:20:5310–20.
70. Sanchez-Monge R., Pascual C.Y., Diaz-Perales A., FernandezCrespo J., Martin-Esteban M., Salcedo G. Isolation and characterization of relevant allergens from boiled lentils. *J Allergy Clin Immunol*. 2000; 106:5:955–61.
71. Finkina E.I., Balandin S.V., Serebryakova M.V., Potapenko N.A., Tagaev A.A., Ovchinnikova T.V. Purification and primary structure of novel lipid transfer proteins from germinated lentil (*Lens culinaris*) seeds. *Biochemistry (Mosc)*. 2007; 72:4:430–8.

72. Goggin D.E, Mir G, Smith W.B, Stuckey M, Smith P.M. Proteomic analysis of lupine seed proteins to identify conglutin beta as an allergen, Lup an 1. J Agric Food Chem. 2008; 56:15:6370– 6377.
73. Peeters K.A, Koppelman S.J, van Hoffen E, van der Tas CWH, den Hartog Jager C.F., Penninks A.H et al. Does skin prick test reactivity to purified allergens correlate with clinical severity of peanut allergy?. Clin Exp Allergy. 2007; 37:0:108–115.
74. Magni C, Herndl A, Sironi E, Scarafoni A, Ballabio C, Restani P et al. One and two-dimensional electrophoretic identification of IgE binding polypeptides of *Lupinus albus* and other legume seeds. J Agric Food Chem. 2005; 53:11:4567–71.
75. Zoccatelli G, Pokoj S, Foetisch K, Bartra J, Valero A, San Miguel-Moncin M.M et al. Identification and characterization of the major allergen of green bean (*Phaseolus vulgaris*) as a non-specific lipid transfer protein (Pha v 3). Mol Immunol. 2010; 47:7:1561–1568.
76. Kumari D, Arora N, Kasera R, Sridhara S, Kumar R, Singh B.P. Isolation and characterization of a 28 kDa major allergen from blackgram (*Phaseolus mungo*). Immunobiology. 2012; 217:9:895-904.
77. Sandeep Kumar., Alok Kumar Verma., Mukul Das., S.K. Jain., Premendra D. Dwivedi. Clinical complications of kidney bean (*Phaseolus vulgaris* L.) consumption. Nutrition.2013; 29:6:821–827.
78. Xu LL, Chen J, Sun LR, Gao X, Lin H, Ahmed I, Pramod SN et al. Analysis of the allergenicity and B cell epitopes in tropomyosin of shrimp (*Litopenaeus vannamei*) and correlation to cross-reactivity based on epitopes with fish (*Larimichthys crocea*) and clam (*Ruditapes philippinarum*). Food chemistry. 2020; 1:323:126763.
79. Pramod SN, Venkatesh YP. Allergy to eggplant (*Solanum melongena*). Journal of allergy and clinical immunology. 2004; 1:113:1:171-3.
80. Pramod SN, Ofori-Anti AO, Cerny R, Goodman RE. IgE Binding to Cross-Reactive Carbohydrate Determinants (CCD) on Proteins of *Phaseolus vulgaris* using Sera from Peanut Allergic Subjects Demonstrates the Potential for False Diagnosis. Journal of Allergy and Clinical Immunology. 2009; 1:123:2:S24.

81. Ofori-Anti AO, Pramod SN, Goodman RE. In vitro IgE Binding to Cross-Reactive Carbohydrate Determinants or Peptide Epitopes in Legume Extracts?. *Journal of Allergy and Clinical Immunology*. 2008; 1:121:2:S241.

UNDER PEER REVIEW