

Original Research Article

Community Health Workers social demographic factors and roles influencing uptake of Maternal Health services in Musanze District, Northern Province, Rwanda.

Comment [M1]: I suggest to add word care to complete statement community healthcare workers instead of community health workers

ABSTRACT

Community health workers integrate individuals of their communities to provide preventive, habitual, and emergency maternity healthcare requirements. The aim of this study was to assess Community Health Workers social demographic factors and roles influencing uptake of Maternal Health services in Musanze District, Northern Province, Rwanda. **Methods:** The study design used was analytical cross-sectional study design utilizing both quantitative and qualitative methods. In this study, a simple random sample approach was employed to choose 208 CHWs and 16 CHW's supervisors were interviewed for qualitative data (KII). The statistical tool for social sciences (SPSS) Version 26, was utilized to examine the data. Bivariate analysis with Chi-square test of independence was used to check the relationship between the dependent and independent variables. Ordinal logistic regression was used as the most suitable inferential statistic because the predictor variables and the dependent were ordinal variables. Qualitative data were analyzed by INVIVO version 10. 208 CHWs participated in the study and all were female. **Results:** This study found that the uptake of maternal health services was regarded as very high. Among those services, deliveries at the HF were at 73.6%, ANC at 65.9%, PF at 54.3%, and the rate of PNC was at 45.7%. This study assessed the influence of social demographic factors on uptake of maternal health services, level of education with $p^* = .001$, and working experience $p^* = .005$ were associated with uptake of maternal health services. The study recommends improvement of maternal health services through family planning, health care during pregnancy, post-partum care, with particularly scaling up key motherly health services, regular training and supervision for CHWs. This will help to achieve third goal of SDG 2030 as *Good Health and well-being for all People* and end up reducing mortality rates in the community unit. The findings of the study will be helpful to Ministry of Health (at District and national level) in taking strategic steps towards reducing mortality rates in the unit and country at large.

Comment [M2]: Add Background

Comment [M3]: Don't use abbreviation in abstract, please write full statements in all abbreviated words

Key words: Community health workers 'Sociodemographic factors, Role influencing uptake of maternal health services.

1. INTRODUCTION

Community Health workers play an essential role in the community that guarantees key health services are available and accessible to the community members. World Health Organization established a community-based initiative programs and uses community volunteers to provide health care to their people (WHO, 2007; Lauren Crigler,2014). CHWs performs one or many duties linked to the delivery of healthcare and received relevant training and supervision (WHO,2015). Community health workers are unpaid household workers who make home visits to mothers during pregnancy up to labor and delivery (Haines, Sanders, Lehmann, Rowe, &Law, 2007). Several health organizations around the world use CHWs for primary health care.Rwanda as a member of these countries, has primarily adopted community health workers and considered their role to achieve the optimum in different health indicators Bucagu, M., et al., (2012).The criteria for selecting CHWs vary depending on the Ministry of Health in each country; however, the baseline requirements for selecting CHWs through partners in health are that they must be an adult, normally over the age of 18, have reading skills, and be able to adequately read prescription drug packets, fill out simple forms, and take notes all through routine practice sessions Beatrice, U. (2016).According to MoH,2007a CHW has to be mature enough to be allowed to provide maternal health services, the CHWs who have more experience in services are more respected and trustfully in the community than those who are still young. In the study conducted on the performance of CHWs in Kenya on marital status has demonstrated that CHWs majority of CHWs were married with (76.9%)(Mpembeni et al, 2015). Many programs require a certain level of literacy but in Rwanda, the CHWs must complete primary school and be able to read and write as criteria to be selected as CHWs, and training benefits helped them to make a follow-up of CHWs instructions during pregnancy and after Gatwaza, M. H. (2016).The sociodemographic variables (Age, gender, Education, working experience of community health workers all are essential factors while studying how the communities are motivated to uptake the services provided by these CHWs, behavior change toward uptake of CHWs activities; especially young age group, under 30, while age cohort 40-50 had the highest performance of home visits(Mulingwa, 2014; Marura, H. M. (2015)). The health system in the country has recognized the importance of women in obtaining better health in general, as well as maternity and child health in particular, MH-CHWs in their village area are mothers in this regard and women's continued involvement in the implementation of health initiatives in their communities, particularly in the areas of safe delivery and reproductive healthhas achieved major improvements on mother's well-being(Condo et al. 2014; Tuyisenge, G., (2019).The study recommends improvement of maternal health services through family planning, health care during pregnancy, post-partum care, with particularly scaling up key motherly health services This will end up in reducing maternal mortality rate and thereby ensuring the achievement the target of NDGs 2030 and improve the quality of service provided by CHWs.

2. MATERIALS AND METHODS

2.1. Study design

This was a cross-sectional study utilized quantitative and qualitative methods because it examined the exposure and outcome at a time.

2.2. Research area

Research was carried out in Musanze District. The district has 15 Sectors, 68 cells and 432 villages. It has 16 health centers and 432 community in charge of maternal health, Musanze District has a population of 424,572 people and is densely populated of 530, 4 per km².

2.3. Study population

The participants in this study were community health workers and their supervisors appointed at health facilities levels in Musanze District

2.4. Determining the sample size

The researcher utilized Slovin's formula to calculate the sample size, that proposes the random sample is obtained by multiplying the average population by the number of people in the sample, a square of the confidence level. This means that

$$n = \frac{N}{1 + (N \cdot e^2)}$$

Where;

n is the sample size

N represents the total population, in this case, total number of CHWs in Musanze District= 432

e indicates the level of confidence (0.05).

$$n = \frac{432}{1 + (432)(0.05^2)} = 207.69230 \approx 208$$

So, the estimated sample size was 208 CHWs

2.5. Sampling procedures and techniques

A simple random sampling procedure was used to select 208 CHWs. This number of CHWs was proportionally distributed to 16 health facilities. The table below shows how CHWs were proportionally distribute to 16 Health facilities.

Chart 1: The table below shows the distribution of CHWs to the Health facilities.

Health Center's Name	Number of CHWs per Health facility.	Representative sample proportional to sample size	Rounded Sample size
Kimonyi	13	6.2	6
Musanze	18	8.6	7
Busogo	25	12.03	12
Gataraga	16	7.7	8
Kinigi	23	11.07	11
Kabere	25	12.03	12
Nyange	32	15.4	14
Bisate	29	13.9	14
Karwasa	30	14.4	14
Gasiza	27	13	13
Muhoza	49	23.5	24
Nyakinama	51	24.5	25
Rwaza	32	15.4	15
Gashaki	22	10.5	11
Murandi	18	8.6	9
Shingiro	22	10.5	11
TOTAL=16	432	207.33	208

2.6. Quantitative data collection

In this research a semi-structured questionnaire with open and closed-ended questions was used to collect quantitative data. The questionnaire had two sections, A, B, and Section A captured social demographic characteristics of CHWs. Section B covered the influence of community health workers roles on uptake of maternal health services.

2.7. Qualitative data collection

Qualitative data were collected using key informants Interview (KII) with community health supervisors. The data were collected using a conversation guide with pre-prepared questions, the guide extracted information from the participants. The information was recorded using an audio recorder.

2.8. Quantitative and qualitative Data analysis

The statistical tool for social sciences (SPSS) Version 26 was utilized to analyze quantitative data. Categorical variables were summarized with descriptive statistics for frequencies and percentages. Bivariate analysis with Chi-square test of independence was used to check the relationship between the dependent and independent variables. The variables with a significant relationship with the dependent variable were then modeled with ordinal logistic regression to develop a prediction model. This was done with a 95% confidence interval. Ordinal logistic regression was the most suitable inferential statistic as the dependent variable was ordinal. This statistic established further associations for the variables which were significant in bivariate analysis. Data and results were presented using text, tables, and figures and qualitative data were analysed by NVIVO version 10. Data was recorded using audio recorder, the audios were transcribed to text format for analysis. Transcripts were then uploaded to NVIVO. Similar patterns were identified, and themes were generated.

3. FINDINGS

3.1 Quantitative findings

3.1.1. Influence of CHW's Social Demographic Factors on Uptake of Maternal Health Services

This study involved 208 CHWs responded to various metrics of the research questionnaire. All the participants were female, 100%.

3.1.2. Age category and uptake of maternal health services

Results showed that 75.0% of CHWs who were aged between 41-50 years, ranked the uptake of maternal health services as very high, 73.2% who were aged 51 years and above rated the uptake of maternal health services as very high. In this study, uptake of maternal health services is independent of CHWs age. Chi-square test showed no relationship between the uptake of maternal health services and age of CHWs with $p=0.535$.

3.1.3. Education level of CHWs on uptake of maternal health services

The majority of the respondents, 75.6% had completed primary school, rated the uptake of maternal health services as very high. The level of education of CHWs revealed a significant relationship with the uptake of maternal health services with ($X^2 = 27.330$, $df = 8$ $p^* = .001$)

3.1.4. Occupation of CHWs

More than three quarter, 76.0% of CHWs who were farmers ranked the uptake of maternal health services as very high. Chi square test didn't show any significance association between the occupation of CHWs and the uptake of maternal health services in the community where ($X^2 = 28.152$ $df = 6$ $p^* = 1.000$).

3.1.5. Marital status of respondents on uptake of maternal health services

More than three quarter 76.9% of married CHWs ranked the uptake as very high while 3(1.8%) of them ranked the uptake of maternal health services as moderate. The Chi-square test of independence showed there was no association between the uptake of maternal health services in the community and the marital status of CHWs ($X^2 = 7.861$, $df = 6$ $p^* = .462$).

3.1.6. Working Experience of CHWs on uptake of maternal health services

The study also sought to know the period through which CHWs had been providing maternal healthcare services. Majority of CHWs 135(79.9%) with working experience of 7 – 9 years ranked the uptake of maternal health services as very high. Chi-square test of independence showed a significant relationship between CHWs years of experience and the uptake of maternal health service ($X^2 = 17.312$ $df = 6$ $p^* = .005$).

The two significant variables were modeled with the ordinal logistic regression analysis to develop a prediction model for the uptake of maternal health services. From the findings, the Nagelkerke R-square showed that the model explains that 23.4% of the variance in the uptake of maternal health services was due to these variables. Both CHWs' level of education and their working experience significantly contributed to the prediction model, the logistic regression finding showed that CHWs with a University education led to 15 times increase in the uptake of maternal health services compared with those with no education. Similarly, CHWs with more than 10 years of experience led to 3.6 times increase in the uptake of maternal health services compared with those with 6 months to 3 years of experience.

Table 1: Chi-square table for social demographic characteristics of respondents on Uptake of Maternal Health Services

Social-demographic Factors	Category	Uptake of Maternal Health Services			Chi-square p-value
		Moderate	High	Very High	
Age category of respondents	Between 20 - 30 years	0(0.0%)	1(16.7%)	5(83.3%)	$X^2 = 5.135$ df =6 $p^* =.535$
	Between 31 - 40 years	1(2.3%)	6(14.0%)	36(83.7%)	
	Between 41 - 50 years	2(2.3%)	20(22.7%)	66(75.0%)	
	51 years and above	0(0.0%)	19(26.8%)	52(73.2%)	
Level of Education of respondent	None	0(0.0%)	10(71.4%)	4(28.6%)	$X^2 = 27.330$ df =8 $p^* =.001$
	Primary	3(1.9%)	35(22.4%)	118(75.6%)	
	Ordinary Level	0(0.0%)	0(0.0%)	23(100.0%)	
	Secondary	0(0.0%)	0(0.0%)	5(100.0%)	
	University	0(0.0%)	1(10.0%)	9(90.0%)	
Marital Status of CHWs	Single	0(0.0%)	0(0.0%)	2(100.0%)	$X^2 = 7.861$ df =6 $p^* =.462$
	Married	3(1.8%)	36(21.3%)	130(76.9%)	
	Widowed	0(0.0%)	9(25.0%)	27(75.0%)	
	Divorced	0(0.0%)	1(100.0%)	0(0.0%)	
Occupation/profession of respondents	Farmer	3(1.5%)	46(22.5%)	155(76.0%)	$X^2 = 7.842$ df =6 $p^* =1.000$
	Salaried	0(0.0%)	0(0.0%)	1(100.0%)	
	Artisan	0(0.0%)	0(0.0%)	1(100.0%)	
	None	0(0.0%)	0(0.0%)	2(100.0%)	
Working experience of CHWs	6 months to 3 years	0(0.0%)	9(52.9%)	8(47.1%)	$X^2 = 17.312$ df =6 $p^* = .005$
	Between 4 - 6 years	0(0.0%)	4(50.0%)	4(50.0%)	
	Between 7 - 9 years	1(7.1%)	1(7.1%)	12(85.7%)	
	10 years and Above	2(1.2%)	32(18.9%)	135(79.9%)	

3.2. Uptake of Maternal Health Services per Indicator

The researcher sought to find out the individual contribution of the assessed variable to the uptake of maternal health services, the findings showed that 73.6% of participants responded that the deliveries in health care facilities had the most improvement after CHWs started to work in the community. 65.9% of the CHWs who participated in this study responded that there was a very high increase of mothers completing ANC visits. More than half of the CHWs, 54.3% responded that there was a very high uptake of mothers towards using FP after providing health

education. Moreover, 45.7% responded that there was a very high increase in the rate of post-partum mothers' attendance to PNC visits with respect to CHWs' visits.

Table 2. Frequency table showing the uptake of Maternal Health Services per Indicator

Variable	Category	Frequency (N208)	Percentage (%)
Mothers completed ANC visits has increased at the health center after CHWs started to work in the community	Average	4	1.9
	High	67	32.2
	Very high	137	65.9
CHW's record of pregnant mothers has increased deliveries in a health care facility	Moderate	3	1.4
	High	52	25.0
	Very high	153	73.6
The rate of post-partum mothers' attendance to PNC visits with respect to CHWs visit	Very low	2	1.0
	Low	4	1.9
	Moderate	5	2.4
	High	102	49.0
	Very High	95	45.7
The rate of mother's uptake towards using FP after providing health education on FP	Low	1	.5
	Moderate	6	2.9
	High	88	42.3
	Very high	113	54.3

3.2.1. The Influence of Community Health Workers Roles on Uptake of Maternal Health Services.

The results from findings showed that among those who strongly agreed that sending red alert SMS by CHWs towards maternal health services, 157(75.4%) rated the uptake of maternal health services as very high with chi-square test result ($X^2 = 17.493$ df =6 $p^* = .006$).

Majority of respondents ,157(75.4%) confirmed the influence of monthly report submitted by CHWs to the HC, rated the uptake of maternal as very high with chi-square test result with ($X^2 = 13.111$ df =4 $p^* = .008$).

176(84.6%) strongly agreed that CHWs' enrolment of girls and ladies of reproductive age has contributed positively to maternal health, rated the uptake of maternal health as very high with chi-square ($X^2 = 9.788$ df =2 $p^* = .006$).

150(72.1%) of respondents had very high perception that CHWs' follow-up to the pregnant has improved maternal health, rated the uptake of maternal health as very high with chi-square ($X^2 = 39.540$ df =2 $p^* = .001$).

Those who strongly agreed that health education of CHWs to the pregnant women on ANC has improved maternal health were 172(82.6%), rated the uptake of maternal health services as very high, where chi-square test was significant with ($X^2 = 18.317$ df =4 $p^* = .001$). All these variables were then modelled with the ordinal logistic regression analysis to develop a prediction model on uptake of ANC among mothers. The model fitting statistic was significant ($X^2 = 56.841$, $df = 9$, $p < .001$). The model explained 31.4% of the variance in the Wald criterion demonstrates the influence of the five independent variables on the dependent variable. CHWs' follow-up ($p = .001$) and CHWs education to pregnant women ($p = .027$) significantly contributed to the prediction model. Very high CHW follow-up of pregnant women leads to 5.8 times increase in mothers completing ANC visits. Moreover, CHWs' strong acceptance of the positive contribution of education results in 2.7 times increase in the number of mothers completing ANC visits.

Table 3: Chi-square showing the influence of CHWs Roles on ANC Uptake among pregnant Mothers

CHWs Roles	Category	Mothers completed ANC visits has increased at the health centre after CHWs started to work in the community			Chi-square p-value
		Average	High	Very High	
Sending red Alert SMS by CHWs for pregnant mothers' emergencies has improved maternal health	Disagree	0(0.0%)	1(100.0%)	0(0.0%)	$X^2 = 17.493$ df =6 $p^* = .006$
	Neutral	0(0.0%)	0(0.0%)	2(0.0%)	
	Agree	1(2.1%)	25(52.1%)	22(45.8%)	
	Strongly agree	3(1.9%)	41(26.1%)	113(72.0%)	
The rate of monthly report submitted by CHWs to the health center and contribution toward maternal health services	Moderate	0(0.0%)	0(0.0%)	1(100.0%)	$X^2 = 13.111$ df =4 $p^* = .008$
	High	2(4.1%)	24(49.0%)	23(46.9%)	
	Very High	2(1.3%)	43(27.2%)	113(71.5%)	
CHWs' enrolment of girls and ladies of reproductive age has contribute positively to maternal health	Agree	2(6.3%)	16(50.0%)	14(43.8%)	$X^2 = 9.788$ df =2 $p^* = .006$
	Strongly agree	2(1.1%)	51(29.0%)	123(69.9%)	
CHWs' follow-up to the pregnant has improve maternal health	High	1(1.7%)	38(65.5%)	19(32.8%)	$X^2 = 39.540$ df =2 $p^* = .001$
	Very High	3(2.0%)	29(19.3%)	118(78.7%)	
Education of CHWs to the pregnant women on ANC has improve maternal health	Neutral	0(0.0%)	2(100.0%)	0(0.0%)	$X^2 = 18.317$ df =4 $p^* = .001$
	Agree	3(8.8%)	16(47.1%)	15(44.1%)	
	Strongly agree	1(0.6%)	49(28.5%)	122(70.9%)	

Considering the contribution on uptake of PNC period, the findings showed that post-partum mothers' attendance at the HF has increased at 75.4%. The findings also showed that visiting mothers and giving them health education on uptake maternal health services during ANC up to PNC period has also increased the number post-partum mother's attendance at health facility with $X^2 = 26.749$ df =12 $p^* = .026$.

84.6% strongly agreed that CHW's enrolment of girls and ladies did not influenced the uptake of maternal health services with $X^2 = 7.429$ $df = 4$ $p^* = .087$. CHWs' follow-up with the pregnant mothers up to the post-partum period has improved the uptake of maternal health and increased the number of PNC attendance at HFs significantly with $X^2 = 17.271$ $df = 4$ $p^* = .001$.

Lastly, 82.6% of respondents strongly agreed with the statement that health education of CHWs to the pregnant women on the importance of doing postnatal care has improved uptake of maternal health services. The result demonstrated that education provided by CHWs to the pregnant women on ANC didn't show any significant association with uptake of maternal health services $X^2 = 10.842$ $df = 8$ $p^* = .324$

These two variables such as sending red alert and follow-up made by CHWs were modelled with the ordinal logistic regression analysis to develop a prediction model on PNC uptake among post-partum mothers. The model fitting statistic was significant ($X^2 = 19.085$, $df = 4$, $p^* = .001$).

The Wald criterion demonstrates that the two independent variables significantly contributed to the prediction model: Sending red Alert SMS by CHWs ($p^* = .017$) and CHWs' follow-up ($p^* = .007$). Sending red Alert SMS by CHWs results in 2.3 times increase in PNC uptake among post-partum mothers while CHW follow-up of pregnant women leads to 2.4 times increase in PNC uptake among post-partum mothers compared.

Table 4: Chi-square table showing the Influence of CHWs Roles on PNC Uptake among Post-partum Mothers

CHWs Roles	Category	The rate the post-partum mothers' attendance to PNC visit with respect to CHWs visit					Chi-square p-value
		Very Low	Low	Moderate	High	Very High	
Sending red Alert SMS by CHWs for post-partum mothers' emergencies has improved maternal health	Disagree	0(0.0%)	0(0.0%)	0(0.0%)	1(100.0%)	0(0.0%)	$X^2 = 26.749$ df=12 $p^* = .026$
	Neutral	0(0.0%)	0(0.0%)	0(0.0%)	1(50.0%)	1(50.0%)	
	Agree	0(0.0%)	2(4.2%)	2(4.2%)	32(66.7%)	12(25.0%)	
	Strongly agree	2(1.3%)	2(1.3%)	3(1.9%)	68(43.3%)	82(52.2%)	
The rate of monthly report submitted by CHWs to the health center and contribution toward maternal health services	Moderate	0(0.0%)	0(0.0%)	0(0.0%)	1(100.0%)	0(0.0%)	$X^2 = 16.064$ df=8 $p^* = .093$
	High	0(0.0%)	2(4.1%)	1(2.0%)	31(63.3%)	15(30.6%)	
	Very High	2(1.3%)	2(1.3%)	4(2.5%)	70(44.3%)	80(50.6%)	
CHWs' enrolment of girls and ladies of reproductive age has contribute positively to maternal health	Agree	0(0.0%)	1(3.1%)	2(6.3%)	20(62.5%)	9(28.1%)	$X^2 = 7.429$ df=4 $p^* = .087$
	Strongly agree	2(1.1%)	3(1.7%)	3(1.7%)	82(46.6%)	86(48.9%)	
CHWs' follow-up to the pregnant has improve maternal health	High	0(0.0%)	1(1.7%)	2(3.4%)	41(70.7%)	14(24.1%)	$X^2 = 17.271$ df=4 $p^* = .001$
	Very High	2(1.3%)	3(2.0%)	3(2.0%)	61(40.7%)	81(54.0%)	
Education of CHWs to the pregnant women on ANC has improve maternal health	Neutral	0(0.0%)	0(0.0%)	0(0.0%)	1(50.0%)	1(50.0%)	$X^2 = 10.842$ df=8 $p^* = .324$
	Agree	0(0.0%)	0(0.0%)	2(5.9%)	21(61.8%)	11(32.4%)	
	Strongly agree	2(1.2%)	4(2.3%)	3(1.7%)	80(46.5%)	83(48.3%)	

The researcher also sought to find out the influence of contribution of CHWs on uptake of family planning, the majority, 84.7.0% of respondents strongly agreed that CHWs' enrolment of girls and women of reproductive age on uptake of maternal health services has influenced the rate of mother's uptake towards using FP after receiving health education in the community from CHWs as high 55.1% while 15.3% agreed.

To examine the influence of CHW's roles on FP uptake was computed, the findings showed that health education of CHWs on FP ($X^2 = 25.599$ $df = 6$ $p^* = .001$), Contribution of CHW on providing FP and uptake of maternal health with ($X^2 = 15.290$ $df = 3$ $p^* = .001$), and CHWs' follow-up to the pregnant women $X^2 = 14.044$ $df = 3$ $p^* = .001$ were statistically associated with uptake of family planning.

Further analyses were conducted with ordinal logistic regression, the model fitting statistic was less than the alpha value ($X^2 = 25.435$, $df = 4$, $p = .001$). This shows that follow-up of pregnant mothers and health education positively contributed to increased uptake of family planning services. The odds ratio From the $\text{Exp}(B)$ value generated from the parameter estimates, follow-up of pregnant mothers resulted in 2.3 times increase in the uptake of family planning services among women. CHWs contribute positively on providing FP results in 2.1 times increase. Furthermore, health education had 2.9 times increase in family planning uptake.

Table 5: Chi-square test on the influence of CHWs roles toward family Planning Uptake

Comment [M4]: Because manuscript is rich with information I suggest to be two manuscript instead of one

CHWs Roles	Category	The rate of mother's uptake towards using FP after providing health education on FP				Chi-square p-value
		Low	Moderate	High	Very High	
CHWs' enrolment of girls and ladies of reproductive age has contribute positively to maternal health	Agree	0(0.0%)	2(6.3%)	14(43.8%)	16(50.0%)	$X^2 = 2.480$ df =3 $p^* = .461$
	Strongly agree	1(0.6%)	4(2.3%)	74(42.0%)	97(55.1%)	
Education of CHWs to the pregnant woman on ANC has improve maternal health	Neutral	0(0.0%)	1(50.0%)	0(0.0%)	1(50.0%)	$X^2 = 25.599$ df =6 $p^* =.001$
	Agree	0(0.0%)	4(11.8%)	19(55.9%)	11(32.4%)	
Contribution of CHWs in providing FP at community level	Strongly agree	1(0.6%)	1(0.6%)	69(40.1%)	101(58.7%)	$X^2 = 15.290$ df =3 $p^* =.001$
	Agree	0(0.0%)	5(10.9%)	24(52.2%)	17(37.0%)	
CHWs' follow-up to the pregnant has improve maternal health	High	0(0.0%)	3(5.2%)	35(60.3%)	20(34.5%)	$X^2 = 14.044$ df =3 $p^* =.001$
	Very High	1(0.7%)	3(2.0%)	53(35.3%)	93(62.0%)	

Finally, the researcher examined the influence of Community Health Workers' roles on the rate of deliveries in the healthcare facilities. Preliminary analysis with the chi-square test for independence confirmed that all the variables had a significant relationship with the rate of deliveries in healthcare facilities. These were: Sending rapid SMS by CHWs ($X^2 = 34.346$ df =6 $p^* < .001$), rate of the monthly report submitted by CHWs ($X^2 = 11.612$ df =4 $p^* = .022$).

CHWs' enrolment of girls and ladies of reproductive age ($X^2 = 12.510$ df =2 $p^* = .001$), CHWs' follow-up to the pregnant ($X^2 = 22.279$ df =2 $p^* = .001$), and health education of CHWs to the pregnant women ($X^2 = 14.886$ df =4 $p^* = .004$). The ordinal logistic regression analysis to develop a prediction model on the rate of deliveries in health facilities. The model fitting statistic was significant ($X^2 = 56.500$, df =9, $p = .001$). The odds ratio shows Sending red Alert SMS by CHWs resulted in 4.3 times increase in

the rate of deliveries in healthcare facilities. Health education results in 2.8 times increase in the rate of deliveries in healthcare facilities.

Table 6: Chi-square test showing the influence of CHWs Roles on uptake of deliveries at health facilities.

CHWs Roles	Category	CHW's record of pregnant mothers has increased deliveries in a health care facility			Chi-square p-value
		Moderate	High	Very High	
Sending red Alert SMS by CHWs for pregnant mothers' emergencies has improved maternal health	Disagree	0(0.0%)	1(100.0%)	0(0.0%)	$X^2 = 34.346$ df =6 $p^* = .001$
	Neutral	0(0.0%)	0(0.0%)	2(100.0%)	
	Agree	1(2.1%)	26(54.2%)	21(43.8%)	
	Strongly agree	2(1.3%)	25(15.9%)	130(82.8%)	
The rate of monthly report submitted by CHWs to the health center and contribution toward maternal health services	Moderate	0(0.0%)	0(0.0%)	1(100.0%)	$X^2 = 11.612$ df =4 $p^* = .022$
	High	1(2.0%)	20(40.8%)	28(57.1%)	
	Very High	2(1.3%)	32(20.3%)	124(78.5%)	
CHWs' enrolment of girls and ladies of reproductive age has contribute positively to maternal health	Agree	2(6.3%)	14(43.8%)	16(50.0%)	$X^2 = 12.510$ df =2 $p^* = .001$
	Strongly agree	1(0.6%)	38(21.6%)	137(77.8%)	
CHWs' follow-up to the pregnant has improve maternal health	High	1(1.7%)	28(48.3%)	29(50.0%)	$X^2 = 22.279$ df =2 $p^* = .001$
	Very High	2(1.3%)	24(16.0%)	124(82.7%)	
Education of CHWs to the pregnant women on ANC has improve maternal health	Neutral	0(0.0%)	0(0.0%)	2(100.0%)	$X^2 = 14.886$ df =4 $p^* = .004$
	Agree	2(5.9%)	15(44.1%)	17(50.0%)	
	Strongly agree	1(0.6%)	37(21.5%)	134(77.9%)	

3.3. Qualitative findings

Qualitative approaches used key informants' interview on CHW's supervisors, two themes such as Sociodemographic factors of CHWs and influence of Community health workers role on uptake of maternal health services were analyzed and generated.

3.3.1. Influence of sociodemographic factors of CHWs on uptake of maternal health services.

From key informants interview the following sub themes were generated from the theme sociodemographic factors of CHWs and uptake of maternal health services; Age, Education, Occupation, Marital status, Working experience.

.....Age and uptake of maternal health services , the qualitative findings as from interview with CHWs supervisor; *CHWs who are more than 40 years old and served for a long period as CHWS are more trustful than those who are still young, this has been identified by the big number of pregnant women who received health care services at the community level, additional to this, a portion of CHWs with age more than 70 years old have a problem of vision, unable to visit all pregnant women, and make follow up in their respective village* (KII 9).

..... Level of education and uptake of maternal health services, Interviews with CHWs' supervisor 16 stated that *"Education and training provided to the CHWs have influenced uptake of maternal health services as the highly educated are more active in training and reporting compared to those with lower education"*(KII 16).

..... Occupation of CHWs and uptake of maternal health services *"The majority of people in Rwanda including our CHWs are farmers so they are combining their daily activities with providing health services in their respective villages, this could probably negatively affect provision of maternal health services in the country as our CHWs would spend most of their time in erring their livelihood rather than focusing in providing maternal health services"* (KII 10).

.....About Marital status of CHWs and uptake of maternal health services, the findings from the qualitative data were not in agreement with the findings from quantitative data; *married CHWs are more trustful by mothers in the community than those who are still single in the services* (KII 16).

..... The same was also reported in an interview with CHWs supervisor that'' *Experience of CHWs is a more important factor to consider because as well as CHWs are experienced in their work it means the skills also were improved because of training received in different packages*

comparatively to new CHWs recruited on the services and this has influenced the uptake of maternal health services, the reason given by interviewed supervisors (KII 16).

3.2.3. Influence of community health workers roles on uptake of Maternal Health Services.

Four indicators were assessed to find out the level of uptake of maternal health services, these were the number of mothers completing ANC visits, number of mothers delivering at health care facilities, PNC attendance, and rate of FP uptake among mothers. These findings are supported by the CHWs Supervisors in the interview conducted to them.

..... ANC at health facilities have been increased through the effort of CHWs done in health education and the promotion of visiting mothers in their village. This has influenced positively the changes of some miss understanding for certain mother and escorting pregnant women to deliver at health facility has also increased deliveries at a health facility, visit done by CHWs during the ANC period up to delivery this has decreased maternal and child mortality (KII 16).

.....PNC, CHWs know pregnant women in the village and mothers who have delivered, this record facilitated them to make follow up and encourage mothers to respect the postnatal care period as it has given at health facilities, with partnership and collaboration with CHWs and HF, PNC attendance has increased at health facility (KII 16).

.....PF, family planning attendance has increased at health facilities since CHWs started providing services at the community level. Other reasons are that health education is provided on family planning services. Certain mothers are transferred to HF to get the services that are not done by CHWs such as intrauterine devices (IDU), Contraceptive implants, and injectable contraceptives (KII 16).

4. Discussions

The purpose of conducting this research was to assess the influence of CHWs on the uptake of maternal health services in the Musanze district. 208 CHWs participated in the study and all were female. This study found that the uptake of maternal health services was regarded as very high. Among those services, deliveries at the HF were at 73.6%, ANC at 65.9%, PF at 54.3%, and the rate of PNC was at 45.7%. The results showed that Sending red Alert SMS by CHWs for pregnant mothers' emergencies by CHWs towards maternal health services with chi-square test

Comment [M5]: This is not discussion because there is no comparison between previous studies, the paragraph look like results please modified

result ($X^2 = 17.493$ df =6 $p^* = .006$). CHWs' enrolment of girls and ladies of reproductive were significant with ($X^2 = 9.788$ df =2 $p^* = .006$). CHWs' follow-up to the pregnant has improved maternal health services with chi-square ($X^2 = 39.540$ df =2 $p^* = .001$). Health education of CHWs to the pregnant women on ANC has improved maternal health where chi-square test was significant with ($X^2 = 18.317$ df =4 $p^* = .001$). The findings showed that health education of CHWs on FP were statistically significant with ($X^2 = 25.599$ df =6 $p^* = .001$). This study also assessed the influence of social demographic factors on uptake of maternal health services, level of education with $p^* = .001$, and working experience $p^* = .005$ were associated with uptake of maternal health services.

Conclusion

The study found that the uptake of maternal health services in Musanze District was 76.4%, while deliveries at HF, ANC, PF, and PNC were 73.6%, 65.9%, 54.3%, and 45.7% respectively. The study examined the influence of social demographic factors on uptake of maternal health services and results showed that there was a significant association between the level of education of CHWs and uptake of maternal health services and working experience of CHWs.

Recommendations

1. The study recommends improvement of maternal health services through family planning, health care during pregnancy, post-partum care, with particular hard work to scaling up key motherly health services
2. The CHWs who are more than 70 years of age should replace or retire because many of them have vision challenges and they are not able to carry out their activities as expected.

This will help to achieve third goals of SDG 2030 as *Good Health and well-being for all People* and end up reducing mortality rates in the community unit.

Comment [M6]: Not be an italic style

Ethical Considerations

The MKU Institutional Research Ethics and Review Committee (IREC) provided ethical approval for this study. The permission to perform the research was sought from Musanze district after presenting an official authorization letter from MKU. Following receiving written

permission, the investigator went to the health facilities to consent the participants to participate. The participation was voluntary, and confidentiality was assured to the participants by excluding any information that can directly identify the participants.

ACKNOWLEDGEMENTS

I would like first to express my gratitude to the Lord for his mercy, love, and blessing in my existence. I feel obligated to express my heartfelt gratitude to my research supervisors for their consistent help, tireless assistance and constructive idea as well as other lectures from Mount Kenya University's Department of Epidemiology and Biostatistics.

UNDER PEER REVIEW

REFERENCES

Comment [M7]: Please update old references

1. Beatrice, U. (2016). Sicom use and reduction of maternal mortality rate by community health worker in Huye district, Rwanda.
2. Black, R. E., Laxminarayan, R., Temmerman, M., & Walker, N. (2016). Reproductive, maternal, newborn, and child health.
3. Bucagu, M., J. M. Kagubare, P. Basinga, F. Ngabo, K. Barbara, A. C. Lee, M. Bucagu, (2012). Impact of health systems strengthening on coverage of maternal health services in Rwanda: A systematic review. *Reproductive health matters*.
4. Brown A, Malca R, Zumaran A, Miranda J.J. (2006). *On the front line of primary health care: the profile of community health workers in rural Quechua communities in Peru*. *Hum Resour Health* 4(1):11.
5. Condo, J., Mugeni, C., Naughton, B., Hall, K., Tuazon, M. A., Omwega, A., & Binagwaho, A. (2014). Rwanda's evolving community health worker system: A qualitative assessment of client and provider perspectives. *Human resources for health*.
6. Crigler, L. (2014). Rwanda's Community Health Worker Program. Ministry of health. (2018). *Rwanda: Nutrition Profile*.
7. Crispin, N., Wamae, A., Ndirangu, M., Wamalwa, D., Wangalwa, G., Watako, P., & Mbiti, E. (2012). Effects of selected socio-demographic characteristics of community health workers on performance of home visits during pregnancy: A cross-sectional study in Busia District, Kenya. *Global journal of health science*.
8. Declaration of Alma-Ata (1978). International Conference on Primary Health Care, Alma-Ata, USSR. Accessed on 15-01-2021 from http://www.who.int/hpr/NPH/docs/declaration_almaata.pdf
9. Dowswell, T., Carroli, G., Duley, L., Gates, S., Gülmezoglu, A. M., Khan-Neelofur, D., & Piaggio, G. (2015). Alternative versus standard packages of antenatal care for low risk pregnancy. *Cochrane Database of Systematic Reviews*, (7).

10. Gatwaza, M. H. (2016). Adherence of community health workers to dosing and referral national guidelines for the community case management of children under five years, in Gasabo District, Rwanda.
11. Haines, A., Sanders, D., Lehmann, U., Rowe, A. K., Lawn, J. E., Jan, S., ... & Bhutta, Z. (2007). Achieving child survival goals: Potential contribution of community health workers.
12. Martin (2005). Sample size determination and use. *Washington D.C.; Library congress.*
13. Marura, H. M. (2015). Role of community health workers in enhancing maternal health care program in Mombasa county, Kenya (Doctoral dissertation, University of Nairobi).
14. Ministry of Health. (2007). (*Guide for the Implementation of Community Health*). Kigali: Ministry of Health.
15. Mulingwa M.W., (2014). *Performance among Community Health Workers in Njiru District, Nairobi County, Kenya*. School of Public Health of Kenyatta University.
16. Ngabo, F., Nguimfack, J., Nwaigwe, F., Mugeni, C., Muhoza, D., Wilson, D. R., ... & Binagwaho, A. (2012). Designing and implementing an innovative SMS-based alert system (RapidSMS-MCH) to monitor pregnancy and reduce maternal and child deaths in Rwanda. *The Pan African Medical Journal*, 13.