

Original Research Article

TREATMENT SEEKING BEHAVIOR OF PREGNANT WOMEN ABOUT MALARIA IN LAYYAH

[Have a list of authors and their affiliations here](#)

ABSTRACT

Background and Purpose: Malaria during pregnancy and childhood is the major public health problem and about half of the population is at risk of developing malaria in the world. During pregnancy, it causes severe anemia, low birth weight babies, abortion and many other severe complications. Knowledge plays main role to prevent this disease and therefore in this study health seeking behavior of pregnant women is explored to determine frequency of malaria suspected cases and the reasons for non-utilization of Government health facilities by pregnant women.

Methods: A total 381 pregnant women were selected through multistage random sampling. A structured questionnaire was used to know about the knowledge and practice of pregnant women about malaria. A survey was conducted in union council LohanchNashaib of district Layyah. Data was collected and analyzed by using SPSS version 20.

Results: Knowledge was found poor in the majority of the respondents. The major sources of treatment was private health facilities. The major reasons of non-utilization of public health facilities were poor accessibility to find transport, trust on public medical providers, and their bad attitude.

Conclusion:

On the basis of these findings, it was concluded that malaria knowledge has shown to be an important factor for prevention and treatment seeking practices about malaria. Health education and literacy promotion should strengthen the knowledge about malaria prevention and treatment seeking behavior.

Key Words: Malaria, Seeking behavior, public health, pregnant women, Layyah, Pakistan

Comment [Ma1]: Does not make sense

Comment [Ma2]: Do you mean lack of trust?

Comment [Ma3]: patients or medical providers

INTRODUCTION

The health and economy of many developing countries are effected by malaria more; than any other disease. There were 5 million increase of malaria cases reported in 2016 than 2015 (211 million), and 216 million cases of malaria in 2016 (~~According to the latest~~ World Malaria Report, ~~released in~~ November 2017). ~~According to~~ WHO report of 2017, 3.3 billion world population is at risk of malaria . Pakistan is one of the developing countries with low/moderate malaria endemicity. 3/4th of population in the rural areas are at risk (1). Between 2010 and 2016, WHO reported that estimated malaria cases were reduced more than 45%. In Pakistan, Afghanistan, India, Ethiopia, and Indonesia, 85% estimated *vivax* malaria cases occurred in 2016. Over the man's long history, it has been the disease that has most affected his health, economy & policies. It has killed more people than any other disease. It is a serious disease of wide distribution, genus *Plasmodium* has five species (*Plasmodium falciparum*, *P.vivax*, *P.ovale*, *P.malariae*, and *P.knowlesi*) that affect human. *P.falciparum* causes is the most severe form malaria with high fever & chills (2). The risk of premature delivery, spontaneous abortion, stillbirths, ~~severe~~ severe maternal anemia, 1/3rd of preventable low birth weight (LBW) babies & maternal death also increases with malaria in pregnancy (3). *P.vivax* is a wide spread but less dangerous type. Humans are infected with malaria by the bite of more than 30 female mosquitos of *Anopheles* species. ~~Mostly infection during pregnancy in Africa is caused by~~ *P.falciparum*, the most common malarial species in Africa (4). Annually 1 million or more people die of malaria & these are mostly infants, young children & pregnant women in Africa(5). Malaria is a threat for pregnant women & their babies, up to 200,000 newborn deaths each year are as a result of malaria infection in pregnancy(6). Moreover, each year approximately 50 million women living in malaria endemic countries throughout the world become pregnant (7,8).

Now the first-line treatment is AS+SP for all countries in the WHO Eastern Mediterranean Region. Treatment failure rates with AS+SP were less than 10% in Afghanistan, Iran (Islamic Republic of), Pakistan and Yemen.

Layyah is a malaria endemic area, reports reviewed at the Executive District Health Officer Health office showed that there were 6,145 malaria confirmed cases in 2010, ~~&~~ 921 malaria confirmed cases in 2011 & 421 malaria confirmed cases up till 31st August 2012(9).

Aim:-

Comment [Ma4]: Use numbers for reference for consistence

Formatted: Font: Italic

Comment [Ma5]: write in full,

To determine the treatment seeking behavior of pregnant women about malaria.

Objectives:-

To determine frequency of malaria suspected cases in pregnant women.

To ascertain treatment seeking behavior of pregnant women.

To determine the reasons for non-utilization of Government health facility

Comment [Ma6]: Explain this in the introduction

METHODOLOGY

This Study was conducted in Layyah, which is located in the South Punjab in Dera Ghazi Khan Division, in Punjab, Pakistan(10). The District of Layyah ~~is was~~ selected for this study because malaria is endemic ~~due here due~~ to its riverine location & agriculture occupation of its population. All pregnant women in Layyah which were estimated to be as 3.4% of the population which ~~were was~~ 4227. A descriptive Cross Sectional study was carried out. Adopted from WHO's Malaria case management, operations manual. Pregnant women with "fever history in the previous 3 days with no features of other severe diseases" [reference?](#).

Comment [Ma7]: How is this linked to malaria?

Comment [Ma8]: Clarify if this was the population or pregnant women

Inclusion Criteria:-

All pregnant women registered in LHWs register having good & complete records. All pregnant women from the 1st trimester to the last of 3rd trimester in study area. Pregnant women having history of fever were included.

Pregnant women with no history of fever or fever for last 3 days with or without symptoms of severe diseases, having same socio-demographic features were included as comparative group.

Exclusion Criteria:-

Pregnant women in the LHW register whose records were not updated.

Two Stage Sampling was done, estimated population of district Layyah was taken as 1,491,915.

Formatted: Justified

In the 1st stage, 14 BHUs were selected from the 12 union councils situated at the riverine belt of the Layyah which were the suspected breeding sites of the mosquitoes. In the 2nd stage by utilizing simple random sampling one Union Council LohanchNashaib located in the riverine belt was selected which was having a population of 30,040 with estimated expected pregnant women population as 1,021.

It was a 3 month study, carried out from 3rd week of September 2012 to end up in the 2nd week of December 2012. A semi structured standardized questionnaire in Urdu with multiple choices was developed from the questionnaires which have been used in earlier studies & from different

articles related to malaria in pregnancy. The questionnaire was pretested in the pregnant women of nearby union council Ehsan Pur.

Ethical Considerations:-

Written consent was taken on the prescribed Performa. Confidentiality was ensured. All the ethical guidelines regarding research on human subjects were followed. Institutional Review Board (IRB) of the Health Services Academy (HSA) Islamabad approved the Research proposal from the???????

By quantitative methodology Data was analyzed by Statistical Package for Social Sciences (SPSS version 20). Descriptive Statistics (frequency, percentage) primarily summarize the data and make it more graspable.

RESULTS

381 pregnant women were interviewed who gave consent & become eligible for study. For the purpose of analysis we made 3 groups A, B and C, details are given shown in Table 1. Results of microscopy are given shown in Table 2. Age wise data is given shown in Table 3.

Table 1:- Summary of results

Total Sample Size	No of Individuals with Fever	Suspected Malaria	Lab. Confirmed Malaria Cases
	Group A	Group B	Group C
381	314	67	23

Table 2:- Results of Malarial Parasitology

Total No. of positive cases	Individuals with P.vivax infections?? clarify	P.falciparum	Mixed
23	16	4	3

Table 3:- Distribution of respondents according to age

Age (Y)	Total Respondents		Group A (No Fever)		Group B (Suspected Malaria)		Group C (Confirmed Malaria Cases)	
	No.	%	No.	%	No.	%	No.	%
< 20	22	5.8	19	6.1	3	4.5	1	4.3
20 – 39	338	88.7	277	88.2	61	91	21	91.4
40 – 49	19	5	16	5.1	3	4.5	1	4.3
> 50	2	0.5	2	0.6	0	0	0	0
Total	381	100	314	100	67	100	23	100

Good knowledge about malaria was found good in 7.3% in group A, 6% in group B and 4.3 % in group C. It seems gradually decreasing. It showed gradual decrease from group A to group C. On the average only 17.6 % of respondents have good knowledge about malaria. Proportion of good knowledge was found high in group A. Knowledge about malaria was found satisfactory was 43.3 % in group A, 58.2 % in group B and 56.5% in group C. On an average 53% of the respondents have satisfactory knowledge. Proportion of satisfactory knowledge was found high in group B. Poor knowledge about malaria was found 49.4% in group A, 35.5% in group B and 39.1% in group C. Group A was ranked higher in poor knowledge. On an average knowledge about malaria was found poor in 41.4% of the respondents (Figure 1).

Comment [Ma9]: Replace with better word

Comment [Ma10]: How do you measure this??

Comment [Ma11]: Explain how you determine this




Figure 1:- Knowledge distribution among groups about malaria




Figure 2:- Treatment seeking sources of pregnant women among groups




Figure 3:- Reasons of non-utilization of ~~Gov~~ health facilities among group of pregnant women

~~It was found that a~~ Almost one classical symptom of malaria was recognized ~~by-in~~ pregnant women...~~what missing ionformation??~~. The largest recognition in group A was found about 78% for chills and rigors followed by fever 72%. In group recognition for chills and rigor was 81% and for fever was 76% which was far better from ~~other 2 groups~~. In group C it was 78% for chills and rigors and 76% for fever. The knowledge about transmission of malaria was found good among all the groups i.e. 98.4% in group A, 95.5% in group B and 91.3% in group C. On the average about 95% of the respondents recognized mosquitoes as malaria transmitter in human beings. 75.2% in group A, 73.1% in group B and in group C 78.3% use bed net as a measure to prevent from mosquitoes' bite which was found about similar within groups. The average 75.5% of the respondents used bed net.

Comment [Ma12]: Name the groups

In group A the source of treatment seeking about malaria was found ~~Gov.~~government Health Facility after that the source was from private clinic i.e. 37.2% and also a large proportion in this group utilize LHWs as the treatment seeking source of malaria(21.5%). In group B the treatment seeking source of the majority was found from LHWs(27.3%) after that the private clinic was found the other big source of treatment seeking malaria (24.6%). Only 17.5% utilize ~~governmentGov.~~Health Facility as a source of treatment seeking about malaria. In the group C situation was found same as in group B i.e. LHWs (27.3%), private clinic (27.3%) and ~~governmentGov.~~ Health Facility (18.2%). A small proportion of people also utilize the other source like pharmacy traditional healers etc.

In group A the major cause of non-utilization of the ~~governmentGov.~~Health facility was found ~~to be~~ difficulty in transportation (29.3%) ~~after that another the~~ cause was found ~~to be that~~ they don't trust on ~~governmentGov.~~medical workers (21%) and they don't like ~~the attitude of~~ ~~governmentGov.~~medical workers (16.6%). In group B the major cause was found that they don't trust ~~governmentGov.~~medical workers (28.4%) and they don't like the attitude of the ~~governmentGov.~~medical (workers 22.4%) while the difficulty in transportation was found the 3rd major cause (13.4%). In group C the reasons of non-utilization of Gov. health facilities were found similar as in group B i.e., don't trust on ~~governmentGov.~~medical workers (30.4%), don't like the attitude of ~~governmentGov.~~medical workers (21.7%) and difficulty in transportation (8.7%).

Comment [Ma13]: Explain to make it scientific

DISCUSSION

This study revealed that majority of the respondents interviewed were 20-39 (88.2%) years of age. This indicates that the respondents were adult and of the child bearing age having a relative poor (47%) knowledge about malaria in pregnant women. Different results of similar studies conducted among pregnant women in rural area of ~~Ethopia~~Ethiopia were found in which knowledge was found good about malaria in pregnant women (11). The reasons of different result were different social and demographic characteristics of the population and the educational status which was low in our study. 78-81% pregnant women in our study were aware of at least one classic symptom of malaria in our study while it was 92.7% in the similar study~~????~~ ~~incomplete~~ (12). In some other similar studies where knowledge about malaria was found good but poor for the consequences of malaria supported the findings of our study that the poor

knowledge has a serious implication and is likely to affect the utilization of treatment seeking by the pregnant women(13,14).

In our study it was found that only 50% of pregnant women utilize government Gov. health facilities for the treatment of malaria. The major reasons of non-utilization of the government Gov. health facilities were found to be difficulty in transportation (29.3%) followed by lack of trust on government Gov. medical workers (21%) and dis- likeness of attitude of government Gov. medical workers (16.6%) might reflect issues of accessibility and quality in government Gov. health facilities. Health seeking behavior in different countries regarding malaria treatment was found different(15). In another similar study where government Gov. health facilities were found easily accessible to the pregnant women as on ten minutes of walking distance or by public transport on minimal-affordable fare but the utilization is low is likewise our study that the government Gov. health facility was available nearby but utilization was poor, we found that this might be due to distrust on and dislikeness of government Gov. medical workers(16).

Comment [Ma14]: Clarify

Malaria program coordinator should focus on awareness campaign about knowledge of malaria and radio broadcasting health messages. Answers might have been varied in cross sectional design because it gives information at certain point in different malaria seasons and time. Data was competed from the information given by the interviewers.

CONCLUSION

Knowledge about malaria is an important factor for its prevention. Promotion of literacy and health education about malaria transmission and prevention method should be strengthened.

REFERENCES

-
- 1 Bhatti MA, Azharuddin M, Bhatti S, Islam M. and Khan MA. Malaria in Pregnancy: The Perspective in Pakistan. JPMA (2007):57(1):17-25.
 - 2 WHO. World Malaria Report 2011. Geneva: WHO(2011).

-
- 3 WHO. 10 facts on malaria. Geneva: WHO (2012):
<http://www.who.int/features/factfiles/malaria/en/index.html>.(include date accessed)
 - 4 WHO. Roll Back Malaria Partnership: Malaria in Pregnancy. Geneva: WHO(2012):
http://www.rbm.who.int/cmc_upload/0/000/015/369/RBMInfosheet_4.htm (include date accessed)
 - 5Korenromp E, Miller J, Nehlen B, WardlawTe. and Young M. Geneva: WHO; World Malaria Report(2005).
 - 6 WHO. Lives at risk: malaria in pregnancy. Geneva WHO:(2012) Available from:
<http://www.who.int/features/2003/04b/en/index.html>. (include date accessed)
 - 7 Richard W, Stekette RW, Bernard L, Nahlen BL, Parise ME. and Menendez C. The burden of malaria in pregnancy in malaria endemic areas. Am j Trop Med Hyg (2001):64 (1-2):28-35.
 - 8 Murphy SC. and Breman JG. Gaps in childhood malaria. neurological sequelae, anemia, respiratory distress, hypoglycemia & complications of pregnancy. Am j Trop Med Hyg (2001): 64(1-2 Suppl):57 – 67.
 - 9 Chaudhry DTI. Consolidated Malaria Reports, 2010,2011, 2012 Layyah
 - 10 Layyah District - Wikipedia, the free encyclopedia (2012): http://en.wikipedia.org/w/index.php?title=Layyah_District&oldid=522999907. (include date accessed)
 - 11Paulander J, Olsson H, Lemma H, Getachew A. and Sebastain MS. Knowledge, Attitude and Practice about malaria in Rural Tigray, Ethiopia. Am j Trop Med Hyg (2000): 61: 98– 107.
 - 12Yeneneh H, Gyorks T, Joseph L, Pikinging J. AND Tedia S. Antimalarial Drug Utilization by Women in Ethiopia: a Knowledge, Attitude, Practice Study. Bull World Health Organ (1993):71: 63–72
 - 13Enato HFO, Okhamafe AO, Okpere EE. A Survey of Knowledge, Attitude and Practice of Malaria Management among Pregnant Women from two Health Care Facilities in Nigeria. ActaObstetricia(2007): 86:33 – 36
 - 14Nganda RY, Drakely C, Reyburn H. and Marchant T. Knowledge of Malaria influences the use of Insecticide Treated Nets but not Intermittent Presumptive Treatment by Pregnant Women in Tanzania. Malar J(2004):3: 42– 8.
 - 15 Williams HA. and Jones CO. A Critical Review of Behavioral Issues related to Malaria Control in Sub Saharan Africa: What Contributions have Social Scientists made? Soc Sc Med. (2004):59: 501–523.
 - 16Falade CO. and Ajavi LO. "Knowledge and Utilization of Intermittent Preventive Treatment for Malaria among Pregnant Women attending Antenatal Clinics in Primary Health Care Centers

in Rural South West Nigeria: A Cross Sectional Study." BMC Pregnancy Child Birth (2009):
9: 9–28.

UNDER PEER REVIEW