

Attenuation of Potassium Bromate-induced Infertility by African Locust Bean (*Parkia Biglobosa*) Seed

ABSTRACT

Background: *Parkia biglobosa* seed has been reported to enhance sperm quality. Thus, this study sought to assess its ability in attenuating KBrO₃-induced sperm abnormalities.

Methodology: *P. biglobosa* was extracted with soxhlet extractor with ethanol as the solvent. Twenty-four adult male Wistar rats were acclimatized under laboratory conditions and were randomly grouped into A, B, C and D. Group A was given distilled water orally. Animals in groups B, C and D were administered 100 mg/kg body weight of potassium bromate, but groups C and D were also treated with 100 and 200 mg/kg body weight of *P. biglobosa* respectively. Both potassium bromate and *P. biglobosa* were freshly prepared on daily basis and administered to rats by oral gavage. After 28 days of treatment, the animals were sacrificed under mild diethyl ether anaesthetization 24 hours after cessation of last treatment. The cauda epididymis were separated from both testes and tinged with 2 mL of normal saline then teased the cauda epididymis of each rat. The suspension was mixed through a metallic net to avoid any other tissue contamination. This suspension was used for the determination of the sperm parameters.

Results: Exposure of animals to 100 mg/kg body weight of KBrO₃ significantly reduced sperm count, sperm motility, sperm viability and seminal pH but elevated sperm morphology when compared with animals in the control group at $p < 0.05$. These perturbations were attenuated by 100 and 200 mg/kg body weight of *P. biglobosa* seed extract in a dose-dependent manner.

Conclusion: The result of this study showed that potassium bromate induced abnormalities in sperm cells of treated animals, and seed extract of *P. biglobosa* attenuated these abnormalities in a dose-dependent manner. This pharmacological study is a useful tool for further drug development from the natural plant products.

Keywords: Fertility; *Parkia biglobosa* seed; potassium bromate; sperm qualities

1. INTRODUCTION

In many nations around the world, male infertility is steadily rising [1,2]. Infertility among men has reportedly increased to 12% in the USA, according to Louis et al. [3]. Numerous studies have linked exposure to endocrine disrupting chemicals (EDCs), particularly chlorinated compounds that are thought to cause poor sperm quality in males, to the fall in male fertility [4,5]. The EDCs are exogenous substances that disrupt the body's normal processes for

maintaining homeostasis, reproduction, and development by interfering with blood hormone synthesis, secretion, transport, metabolism, binding, and removal [6]. The nuclear and non-nuclear hormone receptors for androgen, estrogen, progesterone, thyroid, and retinoid as well as for nuclear & non-nuclear and neurotransmitter receptors are also thought to be affected by the EDCs [5].

Sequel to digitalization, men now spend a lot of time using laptops in their jobs and are exposed

to radiation from these devices, both of which may have an adverse influence on their ability to conceive [7,8]. The ability to have a fulfilling and safe sexual life as well as reproduce, and the freedom to choose whether and how frequently to do so are all dependent on good reproductive health among people [9].

The halide family of elements, which also contains fluorine, chlorine, and iodine, includes bromides as a member. It is well known that the bromides bind to the same receptors utilized to absorb iodine and interfere with the activity of thyroid hormones, resulting in a low thyroid status [6]. Therefore, bromides can be categorized as an EDC. Due to its oxidizing qualities, potassium bromate (KBrO_3), a member of the bromides group, is employed in the maturation of flour and has been utilized as a dough conditioner in the bread-making process for more than 50 years. For the time being, the WHO, Joint FAO/WHO Expert Committee on Food Additives has suggested a maximum level of 75 ppm of KBrO_3 for treating flour, provided that baking items made from such treated flour have minimal residues of KBrO_3 [10]. According to the same guidelines as for WHO, the level in Japan has been set at 30 ppm. Although KBrO_3 was removed from the list of approved food additives in Canada in 1994, it continued to be identified as an impurity in food packaging sheets [11]. In analytical chemistry, KBrO_3 has also been introduced as an oxidizing agent, a main standard, and a brominating agent. Additionally, the sectors of barley, cosmetics, and water filtration employ KBrO_3 . KBrO_3 has been labeled as potentially carcinogenic in numerous animal experiment studies. Unfortunately, KBrO_3 is carelessly utilized as a bread conditioner in less developed nations like Nigeria in order to increase bakeries' profitability [12,13].

The development and documentation of helpful medicinal plants have been facilitated by the use of herbs and spices in culinary practices [14,15]. The use of herbal remedies and natural products in healthcare has regained popularity across the globe [16]. Numerous studies on the efficacy of

herbal extracts in the treatment of human ailments have been conducted in Africa and approved for usage [16,17]. This interest in medicinal plants may be explained by the known side effects of some conventional medications, their high cost, and the introduction of less expensive traditional medicine or alternative medicine therapies in undeveloped and developing nations [18]. While residents of big African towns engage in the use of both modern medicine and herbs, rural dwellers in some African communities rely primarily on herbs for the treatment of various illnesses [19]. African locust beans are one of these frequently utilized herbs (*Parkia biglobosa*). *P. biglobosa* is a plant from the Leguminosae family that may be found in West Africa's tropical and semi-arid regions, mainly in Nigeria's middle belt and southwest [20]. Due to its high protein, lipid, vitamin, tannin, and mineral content, *P. biglobosa* contains leguminous pods with a tough pericarp; the seeds are used as a food condiment and as an alternative to meat [21]. *P. biglobosa* pods can be harvested and processed into a fermented food item called in the Yoruba, Hausa, and Igbo languages as "iru," "dawadawa," and "ogili," respectively. The fermented pods can be used as a protein supplement because they contain about 40% protein, 32% fat, 24% carbohydrate, and vitamins [22]. There have been suggestions that *P. biglobosa* can improve fertility by improving the quality of male sperm due to its rich nature and popular use in Nigeria [23]. It is possible that this hypothetical study on *P. biglobosa*'s sperm profile effectiveness contributed to perceptions of its rising patronage in Nigeria. A recent study by Obeten et al. [24] showed that *P. biglobosa* seed enhanced sperm quality. Thus, this study sought to assess its ability in attenuating KBrO_3 -induced sperm quality destruction.

2. MATERIALS AND METHODS

2.1 Collection and Extraction of *Parkia biglobosa*

P. biglobosa (African locust bean) seed was purchased from a local market at Orita-

Challenge area of Ibadan, Nigeria and were identified by a botanist. They were sun dried, then a mechanical blender (Moulinex) was used to grind them into powder. According to the procedures outlined by Airaodion et al. [25,26], the extraction was carried out using a soxhlet device with ethanol as the solvent.

2.2 Animal Treatment

The experiment involved twenty-four (24) mature male Wistar rats (*Rattus norvegicus*) weighing between 140 and 160 g. Before the experiment began, they were acclimated for seven (7) days in a laboratory setting. The rats were kept in wire-mesh cages with unlimited access to water and commercial rat food. The animals were housed in conventional temperature and humidity settings with 12-hour light/dark cycles. The Declaration of Helsinki and the regulations set forth by the Committee for the Purpose of Control and Supervision of Experiments on Animals were followed in the conduct of this investigation. Additionally, animal experiments were conducted in compliance with NIH protocol [27]. They were assigned to groups A, B, C, and D at random. Oral distilled water was administered to group A as the normal control. Animals in groups B, C and D were administered 100 mg/kg body weight of potassium bromate, but groups C and D were also treated with 100 and 200 mg/kg body weight of *P. biglobosa* respectively. Rats received daily doses of freshly produced potassium bromate and *P. biglobosa* by oral gavage for 28 days. Twenty-four hours after the last treatment, the animals were sacrificed while being lightly sedated with diethyl ether. The cauda epididymis were separated from both testes and tinged with 2 mL of normal saline then teased the cauda epididymis of each rat. The suspension was mixed through a metallic net to avoid any other tissue contamination. This suspension was used for the determination of the sperm parameters.

2.3 Analyses of Epididymal Sperm Characteristics

2.3.1 Sperm Motility and Count

According to the procedures described by Ogbuagu et al. [28], spermatozoa were retrieved by making small cuts in the cauda epididymis and placed in 0.05 M phosphate buffered saline (pH 7.4). Sperm content and sperm motility were assessed in sperm suspension. Any type of flagellar or head movement, as well as the slow and quick movements of spermatozoa (also known as progressive movements), were used to calculate the percentage of mobility. A Neubauer hemocytometer was utilized to calculate the sperm count.

2.3.2 Sperm Viability

Eosin/nigrosin staining was utilized for the determination of sperm viability. On a microscope slide, the staining was examined using a ratio of one drop of fresh semen to two drops of staining agent. On a different slide, a smear was applied and allowed to dry. Spermatozoa with damaged membranes and unstained spermatozoa were counted at a magnification of 3100. According to Eliasson's research [29], the proportion of undamaged cells correlated with sperm viability.

2.3.3 Sperm Morphology

The procedure outlined by Airaodion et al. [30] was used to determine the morphology of the sperm. To determine the anomalies in spermatozoa, sperm suspension was stained with eosin, and smears were produced on slides, dried by air, and made permanent. With an oil immersion objective and x100 magnification, the slides were examined under the microscope. Spermatozoa that were morphologically aberrant were enumerated, and their proportion was calculated.

2.3.4 Determination of Seminal pH

According to the procedure outlined by Airaodion et al. [31], the pH of the seminal solution was measured using pH paper with a range of 6.0 to 10.0. A drop of the material was uniformly distributed on the pH paper after being fully mixed. After roughly 30 seconds, the impregnated zone's color became consistent,

and the color was compared to the calibration strip to determine the pH, with the matching value being recorded.

2.4 Statistical Analysis

The results are presented as the mean \pm standard deviation. One-way Analysis of Variance (ANOVA) and Tukey's test were used to determine the degree of homogeneity among the groups. P values 0.05 were taken into consideration as statistically significant for all analyses, which were performed using Graph Pad Prism Software.

3. RESULTS

Exposure of animals to 100 mg/kg body weight of KBrO_3 significantly reduced sperm count (fig. 1), sperm motility (fig. 2), sperm viability (fig. 3) and seminal pH (fig. 5) but elevated sperm morphology (fig. 4) when compared with animals in the control group at $p < 0.05$. These perturbations were attenuated by 100 and 200 mg/kg body weight of *P. biglobosa* seed extract in a dose-dependent manner.




Fig. 1: Effect of *Parkia biglobosa* seed extract on the Sperm Count of Potassium bromate-induced Male Rats




Fig. 2: Effect of *Parkia biglobosa* seed extract on the Sperm Motility of Potassium bromate-induced Male Rats




Fig. 3: Effect of *Parkia biglobosa* seed extract on the Sperm Viability of Potassium bromate-induced Male Rats




Fig. 4: Effect of *Parkia biglobosa* seed extract on the Sperm Morphology of Potassium bromate-induced Male Rats




Fig. 5: Effect of *Parkia biglobosa* seed extract on the Seminal pH of Potassium bromate-induced Male Rats

4. DISCUSSION

Food additives are one of the biggest issues facing the food business today [13]. The short- and long-term genetic and toxic consequences of all food additives should be tested because they have not yet been adequately controlled. These appear to be of crucial importance to human health as a result of the daily consumption of food additives through food, which can build in the body over the course of a lifetime due to the extremely lengthy exposure [32]. Numerous investigations have shown that KBrO_3 has genotoxic and carcinogenic effects in both experimental animals and human. A variety of short-term genotoxicity testing methods are currently being employed to assess the genotoxic harm caused by KBrO_3 [33]. Products from medical plants are abundant in secondary

metabolites and therapeutic essential oils, which study the therapeutic effect of different medicinal plants as natural antioxidants to treat chronic diseases [34] as DNA damage, mutagenesis, and carcinogenesis. Semen analysis and testicular biopsies are standard laboratory procedures used to evaluate male fertility. Male infertility has been shown to be caused by semen motility, vitality, and morphology [35], in addition to other factors such as malnutrition, genetic abnormalities, and serious sickness, among others [36]. Another study found that most men are either too shy or ashamed to visit a clinic for proper medical consultations, so they instead turn to herbal or plant-based formulations for self-medication [37]. As a result, only a small percentage of men seek proper medical treatment to definitively resolve their infertility issues. The spermatozoa morphology

test is an accurate way to assess the genotoxicity and germ cell toxicity of various chemicals. In the current study, the impact of African locust bean on KBrO_3 -induced sperm morphology mutation in male rats was evaluated. According to the study's findings, as shown in figs. 1 through 3, KBrO_3 significantly ($p < 0.05$) decreased sperm count, motility, and viability when compared to the control group. According to reports, treating animals with potassium bromate caused a noticeably increased level of free radical activity in the testicles and other bodily tissues [38,39]. Carcinogenesis may result from the unchecked generation of free radicals [40,41]. Reduced sperm count, motility, and morphology are just a few examples of the reproductive dysfunction brought on by the alteration in sperm characteristics [42]. According to a related study, adding KBrO_3 to drinking water inhibited the growth of growing rats, which in turn led to lower testicular and epididymal weights [43]. However, these disturbances were alleviated when rats were given KBrO_3 and 100 mg/kg and 200 mg/kg of *P. biglobosa* at the same time. According to a recent study by Ezirim et al. [38], the antioxidant capacity of *P. biglobosa* seed extract allowed it to potentiate KBrO_3 -induced testicular oxidative stress. Following the findings of another, sperm motility and viability increased at a dose of 300 mg/kg body weight of *P. biglobosa* extract, but decreased at a dose of 500 mg/kg [24]. This shows that a low dose of *P. biglobosa* may improve reproductive health by enhancing sperm motility. This outcome backs up Udo-Affah et al. [44]'s observation that mice subjected to *P. biglobosa*'s extract exhibited significantly improved sperm motility. This rise was noted, and it was linked to the extract's high iron concentration. Sperm vitality is the percentage of healthy, live sperm found in the semen [45].

Sperm morphology is a significant and accurate predictor of the toxicity of chemicals on reproductive system cells [46]. They provide a clear indicator of the caliber of sperm production in chemically treated experimental animals since they can be utilized to assess spermatogenic

damage, fertility, and heritable genetic abnormalities [47]. Genotoxic chemicals have the ability to affect the normal course of gametogenesis, which could result in changes to the shape of sperm cells. This process is not autonomous and is governed by intricate mechanisms that are polygenically controlled. A suitable indication of genetic damage may be any qualitative or quantitative change in the aforementioned trait that can be linked to dietary, pharmaceutical, or environmental alterations [48]. In the current study, the consequences of giving male rats dosages of potassium bromate, a dietary additive, were examined in terms of morphological changes in sperm cells. According to our findings, compared to the control group, potassium bromate supplementation significantly increases sperm morphological abnormalities (Fig. 4). These results corroborated those from Elhaddad et al. [32], who found that treatment of animals with KBrO_3 affected the number and motility of epididymis sperm and increased the incidence of defective sperm. This demonstrates how artificial food additives may have an impact on reproductive health. The impact of KBrO_3 on sperm morphology was lessened by administering *P. biglobosa* seed extract at doses of 100 and 200 mg/kg body weight.

Genetic damage may manifest as defective sperm or other phenotypic manifestations during the process of spermiogenesis if the pre-meiotic, meiotic, or post-meiotic spermatogenic cells are exposed to a mutagenic substance that interferes with DNA. A complicated array of morphological and metabolic processes is involved in the production of sperm heads. Sperm may grow abnormally during the spermiogenesis process. Small deletions or point mutations are the most plausible causes of sperm head defects [48,49]. Additionally, it was thought that an aberrant chromosome was the cause [50]. Two other investigations [51, 52] found that variations in testicular DNA or physiological, cytotoxic, or genetic causes might cause abnormalities in sperm heads, which would then impair the differentiation of spermatozoa [53].

The seminal pH represents the harmony between the alkaline seminal vesicular secretion and the acidic prostatic secretion, which make up the majority of the accessory gland secretions. When $KBrO_3$ treated animals in this study were compared to the control animals after 28 days of treatment, it was observed that seminal pH had decreased (Fig 5). This could be because the food additive changes the pH range that is typical for the treated animals. The medium of seminal plasma becomes acidic if the pH decreases, making sperms extremely fragile and raising the mortality rate [54,55]. The impact of $KBrO_3$ -induced seminal acidity was ameliorated by the administration of various dosages of *P. biglobosa* extract.

The majority of medications contains plant extracts as their active components and is widely used since they have few adverse effects. Because of this, herbal remedies are now more commonly used in traditional African homes than conventional medications. Plants have given mankind a wide range of effective medicines to lessen or eliminate infections and disease-related suffering. Some medications produced from plants continue to be significant and important despite advances in synthetic pharmaceuticals [56, 57]. Despite the progress made in contemporary medicine, there are still many illnesses or infections (diseases) for which effective medications have not yet been developed. In order to treat inflammatory disorders like diabetes, liver diseases, gastrointestinal disorders, and infertility, safer drugs from the environment are urgently needed [58, 59]. Research on herbal plants or medicines has been promoted by the interdisciplinary approach used to treat some disorders, and there appears to have been significant progress in recent years in the pharmacological evaluation of diverse plants employed in traditional systems of medicine [60].

The results of the semen parameters seen in this investigation support a study by Hussain et al. [61] that showed a polyherbal formulation could have a synergistic efficacy leading to an improvement in the key semen parameters in

oligospermic males. The findings of the semen analysis indicate that the experimental animals' semen parameters were improved by our herbal formulation of *P. biglobosa* seed extract. Prior to this, medicinal plants were divided into two primary categories: dismutagenicity, which refers to a plant's ability to work outside the cell membrane, and antimutagenicity, which refers to a plant's ability to act in numerous places [32]. Additionally, certain medicinal plants have the capacity to function both outside and inside cells. It's possible that our medicinal plant, *P. biglobosa*, has antimutagenic and disantimutagenic properties. These observations point to novel antimutagenic functions of *P. biglobosa* as well as mutagenic effects of potassium bromate.

5. CONCLUSION

The result of this study showed that potassium bromate induced abnormalities in sperm cells of treated animals, and seed extract of *P. biglobosa* attenuated these abnormalities in a dose-dependent manner. This pharmacological study is a useful tool for further drug development from the natural plant products.

CONSENT

It is not applicable.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests or non-financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. Airaodion AI, Ekenjoku JA, Ogbuagu EO, Okoroukwu VN, Ogbuagu U. *Carica papaya* leaves might cause miscarriage. Asian Research Journal of Gynaecology and Obstetrics. 2019;2(2): 1-9.
2. Airaodion AI, Ekenjoku JA, Ngwogu KO, Ngwogu AC. Consumption of coconut (*Cocos nucifera* L.) water improved fertility parameters in male Wistar rats. Asian

- Journal of Pregnancy and Childbirth. 2019;2(3):1-7.
3. Louis JF, Thoma ME, Sorensen DN, McLain AC, King RB, Sundaram R, et al.. The prevalence of couple infertility in the United States from a male perspective: evidence from a nationally representative sample. *Andrology* 2013; 1: 741-748.
 4. Toft G. Persistent organochlorine pollutants and human reproductive health. *Dan.Med J* 2014; 61:B4967.
 5. Toft G, Jonsson BA, Lindh CH, Giwercman A, Spano M, Heederik D, et al. Exposure to perfluorinated compounds and human semen quality in Arctic and European populations. *Human Reproduction* 2014; 27: 2532-2540.
 6. Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM. Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocrine Review* 2009; 30:293-342.
 7. Eisenberg ML, Kim S, Chen Z, Sundaram R, Schisterman EF, Buck Louis GM (2014). The relationship between male BMI and waist circumference on semen quality: Data from the life study. *Human Reproduction* 29(2):193-200.
 8. Gaskins AJ, Mendiola J, Afeiche M, Jørgensen N, Swan SH, Chavarro JE (2015). Physical activity and television watching in relation to semen quality in young men. *British Journal of Sports Medicine*. 49(4):265-270.
 9. Hajizadeh MB, Tartibian B, Chehrizi M (2017). The effects of three different exercise modalities on markers of male reproduction in healthy subjects: a randomized controlled trial. *Reproduction*. 153(2):157-174.
 10. WHO. Evaluation of Certain Food Additives and Contaminants. Geneva: WHO Press; 2007.
 11. Environment Canada, Health Canada. Proposed risk management approach for bromic acid, potassium salt (potassium bromate). Chemical Abstracts Service Registry Number (CAS RN): 7758-01-2. [Online]. Available from: <https://www.ec.gc.ca/ese-ees/default.asp?lang=En&n=D1613C58-1#top>. [Accessed in August 2022].
 12. Airaodion AI, Ewa O, Ogbuagu EO, Ogbuagu U, Agunbiade AP, Oloruntoba AP. Evaluation of potassium bromate in bread in Ibadan metropolis: Fifteen years after ban in Nigeria. *Asian Food Science Journal*. 2019;7(4):1-7.
 13. Airaodion AI, Awosanya OO, Ogbuagu EO, Ogbuagu U, Njoku OC, Esonu C, Airaodion EO. Assessment of bread in ogbomoso metropolis for the presence of potassium bromate. *Asian Journal of Research in Biochemistry*. 2019;4(2):1-6.
 14. Airaodion AI, Ogbuagu EO, Ekenjoku JA, Ogbuagu U, Airaodion EO. (2019). Therapeutic effect of methanolic extract of *Telfairia occidentalis* leaves against acute ethanol-induced oxidative stress in Wistar rats. *International Journal of Bio-Science and Bio-Technology*. 11(7):179-189.
 15. Airaodion AI, Adeniji AR, Ogbuagu EO, Ogbuagu U, Agunbiade AP. Hypoglycemic and hypolipidaemic activities of methanolic extract of *Talinum triangulare* leaves in Wistar rats. *International Journal of Bio-Science and Bio-Technology*. 2019;11(5):1-13.
 16. Airaodion AI, Akaninyene IU, Ngwogu KO, Ekenjoku JA, Ngwogu AC. Hypolipidaemic and antidiabetic potency of *Allium cepa* (onions) bulb in alloxan-induced diabetic rats. *Acta Scientific Nutritional Health*. 2020;4(3): 1-8.
 17. Ogbuagu EO, Nweke IN, Unekwe PC, Airaodion AI, Ogbuagu U. Weight gain reduction and hypoglycemic effects of *Xylopia aethiopica* fruit extract on Wistar rats. *International Journal of Research and Reports in Hematology*. 2020;5(3):1-8.
 18. Airaodion AI, Ogbuagu U, Ogbuagu EO, Airaodion EO, Agunbiade AP, Oloruntoba AP, Mokelu IP, Ekeh SC. Investigation of aqueous extract of *Zingiber officinale* root potential in the prevention of peptic ulcer in

- albino rats. *International Journal of Research and Innovation in Applied Science*. 2019;4(2):64-67.
19. Airaodion AI, Ogbuagu EO, Airaodion EO, Ekenjoku JA, Ogbuagu U. Pharmacotherapeutic effect of methanolic extract of *Telfairia occidentalis* leaves on glycemic and lipidemic indexes of alloxan-induced diabetic rats. *International Journal of Bio-Science and Bio-Technology*. 2019;11(8):1-17.
 20. Airaodion AI, Airaodion EO, Ogbuagu EO, Ogbuagu U, Osemwowa EU. Effect of oral intake of African locust bean on fasting blood sugar and lipid profile of albino rats. *Asian Journal of Research in Biochemistry*. 2019;4(4), 1-9.
 21. Airaodion AI, Ogbuagu EO. Ameliorative effect of *Parkia biglobosa* (African locust bean) against egg-yolk induced hypertension. *International Journal of Bio-Science and Bio-Technology*. 2020; 12(5):17-25.
 22. Diawara B, Sawadogo L, Amoa-Awua WF, Jakobsen M (2000). Capability building for research and development in quality assurance and fermentation technology for African fermented foods HACCP system for traditional African fermented foods: soumbala. Soumbala. WAITRO.
 23. Soetan KO, Lasisi OT, Agboluaje AK (2011). Comparative assessment of *in-vitro* anthelmintic effects of the aqueous extracts of the seeds and leaves of the African locust bean (*Parkia biglobosa*) on bovine nematode eggs. *Journal of Cell and Animal Biology* 5(6):109-112.
 24. Obeten KE, Isaac UE, Archibong VB, Oshie PA, Etukudo E, Onanuga IO, Ochieng JJ, Monima LA. Histomorphology and sperm profile of animal models administered with aqueous seed extract of *Parkia biglobosa*. *Journal of Cell and Animal Biology*, 2022; 15(1): 1-8.
 25. Airaodion AI, Ngwogu AC, Ekenjoku JA, Ngwogu KO. Hepatoprotective potency of ethanolic extract of *Garcinia kola* (Heckel) seed against acute ethanol-induced oxidative stress in Wistar rats. *International Research Journal of Gastroenterology and Hepatology*. 2020;3(2):1-10.
 26. Airaodion AI, Ogbuagu EO, Ogbuagu U, Awosanya OO, Airaodion EO. Effect of Methanolic extract of *Corchorus olitorius* leaves on hypoglycemic and hypolipidaemic activities in albino rats. *Asian Plant Research Journal*. 2019;2(4):1-13.
 27. National Research Council. Guide for the Care and Use of Laboratory Animals, 8th ed.; The National Academies Press: Washington, DC, USA; 2011.
 28. Ogbuagu EO, Airaodion AI, Ogbuagu U, Ezirim EO, Nweke IN, Unekwe PC. Ethanolic extract of *Xylopiya aethiopica* (African negro pepper) fruit adversely perturbed semen qualities in male Wistar rats. *International Journal of Research and Reports in Gynaecology*, 2022;5(2): 44-55.
 29. Eliasson R. Supravital staining of human spermatozoa. *Fertil Steril*, 2017; 28, 1257.
 30. Airaodion AI, Ayanleke IA, Agunbiade AP, Ogbuagu, EO, Airaodion EO, Ogbuagu U. Antifertility Propensity of *Jatropha curcas* Linn. Leaves on Male Wistar Rats. *International Journal of Research and Reports in Gynaecology*. 2020;3(2): 21-29.
 31. Airaodion AI, Ogbuagu EO, Ekenjoku JA, Okoroukwu VN, Ogbuagu U, Airaodion EO. Antifertility effect of ethanolic leaf extract of *Carica papaya* in male Wistar rats. *Merit Research Journal of Medicine and Medical Science*. 2019;7(10):374-381.
 32. Elhaddad NS, Efkerine EM, Khatab HA, Eldurssi IS, Belkasem EM. Anti-mutagenic Action of *Ruta chalepensis* against Rat Sperm Cell Abnormalities-induced by Potassium Bromate. *Annual Research & Review in Biology* 2020;35(2): 41-51.
 33. Ahmed M, Mahmood R. Protective effect of taurine against potassium bromate-induced hemoglobin oxidation, oxidative stress, and

- impairment of antioxidant defense system in blood. *Environmental toxicology*. 2016;3:304-313.
34. Airaodion AI, Ekenjoku JA, Ngwogu KO, Ngwogu AC. Ethanolic extract of *Garcinia kola* (Heckel) seed possesses antiplasmodial properties against *Plasmodium berghei*. *Asian Journal of Medical Principles and Clinical Practice*. 2020;3(1):33-39.
 35. Mishra RK, Verma HP, Singh N, Singh SK. Male infertility: lifestyle and oriental remedies. *Journal of Scientific Research*, 2012;56:93-101.
 36. Biswas TK, Pandit S, Mondal S, Biswas SK, Jana U, Ghosh T, Tripathi PC, Debnath PK, Auddy RG, Auddy B. "Clinical evaluation of spermatogenic activity of processed Shilajit in oligospermia. *Andrologia* 2010;42(1):48-56.
 37. Vital and Health Statistics, no. 26, CDC (2002). Available online at: <http://www.cdc.gov>.
 38. Ezirim EO, Uche CL, Abali IO, Iwuoha CE, Chika-Igwenyi NM, Onyeaghala CA, Orji SF, Ugwu CN, Ugwu NI, Igwenyi C, Airaodion AI. Therapeutic Potential of *Parkia biglobosa* Seed against Potassium Bromate-induced Testicular Toxicity. *International Journal of Research and Reports in Gynaecology*. 2022;5(3):78-89.
 39. Ugwu CN, Abali IO, Iwuoha CE, Chika-Igwenyi NM, Onyeaghala CA, Orji SF, Igwenyi C, Uche CL, Onyekachi OIN, Nwobodo MU, Airaodion AI. Ameliorative effect of *Parkia biglobosa* (African locust bean) seed against potassium bromate-induced oxidative stress. *Merit Research Journal of Medicine and Medical Science*. 2022;10(8):213-219.
 40. Kurokawa Y, Aoki S, Matsushima Y, Takamura N, Imazawa T, Hayashi Y. Dose-response studies on the carcinogenicity of potassium bromate in F344 rats after long-term oral administration. *Journal of the National Cancer Institute*. 1986;77:977-982.
 41. Sai K, Tyson CA, Thomas DW, Dabbs JE, Hasegawa R, Kurokawa Y. Oxidative DNA damage induced by potassium bromate in isolated rat renal proximal tubules and renal nuclei. *Cancer letters*. 1994;87:1-7.
 42. De Souza Predes F, Diamante M.A, Dolder H. Testis response to low doses of cadmium in Wistar rats. *International Journal of Experimental Pathology*. 2010; 91:125-131.
 43. Elsheikh AS, Fadul TF, Aboagla EM, Gameel AA. Effects of potassium bromate on male rat growth and testicular histology. *Asian Pacific Journal of Reproduction*. 2016;5:376-380.
 44. Udo-Affah G, Obeten KE, Adewale AO, Okim AA. Studies on the effect of aqueous seed extracts of *Parkia biglobosa* on the histology and glycogen profile of the epididymis of Wistar rats. *Chemical & Pharmaceutical Research*. 2021; 3(1): 1-5
 45. Guzick ASM, Overstreet JW, Factor-Litvat P, Brazil CK, Nkajima ST. (2001). National cooperative reproductive medicine Network. Sperm morphology, motility and concentration in infertile men. *National England Journal of Medicine* 345(19):1388-1393.
 46. Isidori AM, Pozza CD, Gianfrilli IA (2006). Medical treatment to improve sperm quality. *Journal of Reproduction Biomedicine* 12(6):704-714.
 47. Devi SA, Umasankar ME, Babu S (2011). A Comparative Study of Antioxidant Properties in Common Indian Spices. *International Research Journal of Pharmaceuticals* 3(5):465-468.
 48. Narayana K, D'souza UJ, Rao KS. Ribavirin-induced sperm shape abnormalities in Wistar rat. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*. 2002; 513: 193-196.
 49. Wyrobek A, Bruce W. Chemical induction of sperm abnormalities in mice. *Proceedings of the National Academy of Sciences*. 1975;72:4425-4429.
 50. Bruce W, Furrer R, Wyrobek A. Abnormalities in the shape of murine sperm after acute testicular X-irradiation, *Mutation Research/Fundamental and Molecular*

- Mechanisms of Mutagenesis. 1974;23:381-386.
51. Agunbiade S, Okonko I, Alimba C, Folarin A, Anugweje K. Effects of a carbonaceous bottling plant effluent on albino mice sperm morphology and testes histopathology. *Nat Sci*. 2012;10:154-60.
52. Odeigah P, Nurudeen O, Amund OO. Genotoxicity of oil field wastewater in Nigeria. *Hereditas*. 1997;126:161-167.
53. Giri S, Prasad S, Giri A, Sharma G. Genotoxic effects of malathion: an organophosphorus insecticide, using three mammalian bioassays *in vivo*. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*. 2002;514:223-231.
54. Ekenjoku JA, Airaodion AI, Okoroukwu VN, Ogbuagu EO, Ogbuagu U. Oral administration of ethanolic extract of *Vernonia amygdalina* leaves might impact negatively on fertility in male Wistar rats. *Asian Journal of Medical Principles and Clinical Practice*. 2019;2(3):1-8.
55. Ogbuagu EO, Airaodion AI. Tiger Nut (*Cyperus esculentus* L.) Boosts Fertility in Male Wistar Rats. *Asian Research Journal of Gynaecology and Obstetrics*. 2020;3(3): 8-18
56. Airaodion AI, Olayeri IM, Ewa AO, Ogbuagu EO, Ogbuagu U, Akinmolayan JD, Agunbiade AP, Oloruntoba AP, Airaodion EO, Adeniji AR, Obajimi OO, Awosanya OO. Evaluation of *Moringa oleifera* leaf potential in the prevention of peptic ulcer in wistar rats. *International Journal of Research*. 2019;6(2):579-584.
57. Airaodion AI, Ogbuagu U, Ogbuagu EO, Airaodion EO, Agunbiade AP, Oloruntoba AP, Mokelu IP, Ekeh SC. Investigation of aqueous extract of *Zingiber officinale* root potential in the prevention of peptic ulcer in albino rats. *International Journal of Research and Innovation in Applied Science*. 2019;4(2):64-67.
58. Airaodion AI, Ogbuagu EO, Airaodion EO, Ogbuagu U, Ekenjoku JA. Antidiabetic effect of ethanolic extract of *Carica papaya* leaves in alloxan-induced diabetic rats. *American Journal of Biomedical Science & Research*. 2019;11(8):93-109
59. Airaodion AI, Ayita EB, Oluba SO, Emaleku SA, Osunmuyiwa OJ. Chemical composition and nutraceutical potential of velvet tamarind fruit pulp. *Asian Journal of Biochemistry, Genetics and Molecular Biology*. 2021;3(5):22-31.
60. Wuthi SP. Antifungal Activities of *Senna alata* extracts using different methods of Extraction. *Journal of Health Resources*, 2010;24(3):117-122.
61. Hussain SA, Ahsan H, Furqan N, Yang W, Hafiz Ansar RS, Yuanda S. Evaluation of the Spermatogenic Activity of Polyherbal Formulation in Oligospermic Males. *BioMed Research International*, 2018;20(7):08-15