

---

# Marijuana and Its Effects in Brain

---

## ABSTRACT

*Cannabis sativa* (hemp) plant is the source of cannabis-based products which comes from any part of the plant such as seeds, stems and flowering tops. The products of this plant are used for recreational purpose. It can also be used to sooth the pain and other medical conditions. Not all products of this plant is mind altering but some of them posses the psychotic effect. The active ingredient of the plant is called as cannabinoids. Almost 120 active chemicals are present in cannabis. According to the scientific reports these cannabis can be used to chronic pain in adults, nausea and vomiting resulting from chemotherapy treatment, some symptoms of multiple sclerosis (MS). These cannabis can be ingested in various forms such as smoking, taking it as supplements, topical treatment, eating either raw or with candies, can be added in tea as a brew. This chapter will address the effect of these cannabis in human brain and its mechanisms of action.

*Keywords: Cannabinoids; cannaboid receptors; dopaminergic neuron; endocannabinoids;  $\Delta^9$ -tetrahydrocannabinol (THC).*

## 1. INTRODUCTION

From the plant *Cannabis sativa*, Marijuana is synthesized, especially from a shredded, green-brown mix of dried flowers, stems, and leaves. It is the most addictive drug apart from alcohol and tobacco. Marijuana is used as hand rolled cigarettes or in pipes or water pipes. Marijuana use is higher in younger people. More than 11.8 million young adults have abused marijuana in 2018 [1,2]. Delta-9-tetrahydrocannabinol (THC) is a cannabinoid molecule in marijuana from *Cannabis sativa* (hemp) plant. In rats exposed to THC from gestational day 15 to postnatal day 9, with THC at a daily dose (5 *per os*) has caused subtle neurofunctional deficits [3]. THC, can also affect the rodent ontogeny process, especially on dopaminergic neuron development and enkephalinergic neuron. Cannaboid receptors (CB1) are seen during prenatal development and there are higher chances of these receptors to be transferred from mother to the offspring through placenta in both human and rodents [4]. Young pregnant women tend to abuse this drug, in western culture. In 2014, approximately 183 million people used marijunana and in 2016, 22 million people abused cannabis [5,6].

## 2. ANIMAL STUDIES

During active brain development, cannaboid receptors are expressed in high level in areas

such as hippocampus, amygdala, and prefrontal cortex (PFC). The age of exposure to cannabis affects the prenatal, perinatal and adolescent periods of life [7]. In rats, THC pretreated animals has shown worse performance with a defect in spatial working memory when tested in radial arm maze [8]. This clearly indicates that in experimental models, adolescent chronic exposure in rats to THC has serious impact on learning and memory. In adolescent rats exposed to THC, it caused defects in synaptic plasticity in hippocampus [9]. Endocannabinoid system is a neuromodulatory network in brain plays a major role in cognitive and physiological process. Heavy marijuana use alters the brain tissue analysed by Voxel-based morphometry [10].

$\Delta^9$ -tetrahydrocannabinol is the primary chemical binds with cannabinoid receptors. Anandamide (N-arachidonylethanolamine) and 2-arachidonoyl glycerol (2-AG) in animals are the main endocannabinoids. CB1 and CB2 receptors, mediate the binding of phyto as well as endocannabinoids. Inside the brain those areas with higher cognitive functions, motor control regions, hindbrain regions express high level of CB1 receptors [11].

## 3. CANNABIS USE AND COGNITIVE FUNCTIONS

---

Regular cannabis usage affects various functions of the brain such as working memory, attention, and cognitive control processing [12]. Functional MRI has shown there is an interaction between recreational cannabis use and cognitive function [13]. Adult chronic cannabis users have reduced attention, cognitive control, poor memory, improper decision-making, motor performance and affective processing [14-22]. A meta-analysis on CANN+ patients with CANN- has revealed several cognitive dysfunctions when tested in premorbid IQ, current IQ, verbal learning, verbal working memory, motor inhibition [23]. CBD induces alterations in brain activity in both the healthy volunteers and people with psychiatric disorder [24].

#### **4. SCHIZOPHRENIA-LIKE PSYCHOSIS**

The *CNR1*, *AKT1*, *DBH* and *5-HTT/SLC6A4* genes may also be involved in modulating the neurological effects of cannabis use. All these genes are also known to be linked with schizophrenia [25]. 'psychotomimetic' effects of  $\Delta 9$ -tetrahydrocannabinol (THC)- haloperidol and their combination was investigated in a double-blind, placebo-controlled, partial three-way crossover ascending dose study in 35 healthy, male mild cannabis users. THC significantly decreased many 'psychotomimetic' effects of CNS [26]. Schizophrenia-like psychosis is reported from heavy use of cannabis as per the recent findings. In a study conducted in twenty two adult males it was found that THC can induce acute psychotic reaction in psychiatrically well individuals [27]. Executive function and inhibitory motor control are also affected by high potency marijuana [28]. And MRI study has revealed that orbitofrontal gyri volume was significantly lower in marijuana users followed by orbitofrontal cortex and forceps minor innervations was higher [29].

#### **5. ALTERNATIONS IN NEURAL CONNECTIVITY**

Apart from natural source from cannabis plant this also is produced synthetically in names such as spice, K2 or Kronic being widely misused . Although cannabis is used in various treatments still its danger can not be overlooked. Consuming delta-9-tetrahydrocannabinol (THC) has lot of effects on synaptic plasticity as a short term change and neuronal structural alternations as the long term change and it affects the endocannabinoid system Exposure to THC at the early life leads to the alternations in neural

connectivity in areas such as cognition and behaviour. Marijuana use not only affects the brain but it also effects the sympathetic nervous system, that controls blood pressure, platelet activation, and electrophysiological effects. During periods of aactive neuronal development, cannabinoid receptors are highly expressed in areas that control cognitive and behaviour. Animal studies have shown that exocannbenoids can damage the processes that control cognition and behaviour [30].

#### **6. ROUTE OF ENTRY**

Marijuana is smoked and enters blood stream through lungs, if taken orally through digestive system it enters the blood. Though blood it reaches brain where it binds with cannebinoid receptors causing alterations. From childhood if it is taken then through midlife it causes neuropsychological decline [31]. In adolescent period if marijuana is used it leads to long term effects on the how the neural connections are made within the brain [32]. These changes in the neural network can impact attention, memory, and learning. THC cause "high" feeling in people. Cannabinoid hypermesis syndrome are characterised by vomiting. Nausea and dehydration [33]. The other names of cannabis are pot, weed and ganja that is the most widely used recreational drug globally. USA and Canada has delegailised this drug that increasing the usage in teen population [34,35].

#### **7. STROKE AND MARIJUANA USE**

Ischemic stroke can occur in individuals who chronically use marijuana [36]. Cerebral infract is another effects of cannabinoids noted through inhalation and combustion [35,37]. CB-1 receptor agonist WIN 55,212-2 is a synthetic cannabis upon injection in Adult male Sprague–Dawley rats for 22 days it caused significant hippocampal morphological alternations [38,39]. Early cannabis exposure causes psychotic like experience in adolescents [40]. THC alters the inhibitory interneurons in cerebellum [41]. Mariuana smoking can increase stroke , a case study who used marijuana regular end up with having the stroke in occipital lobe , the subject is also positive for heterozygous for factor V Leiden associated with thrombosis [42,43]. THC can impact prenatal life which is a time for normal signalling pathways and also it affects the adolescence time by altering the brain maturation. In USA marijuana use is higher and

approximately 3 in 10 patients express marijuana use disorder [44].

## 8. CONCLUSION

Though marijuana is said to be harmless our understanding on its effects in brain has revealed the presence of exiting evidence on abnormal changes in brain and behaviour, so this drug must carefully be used by pregnant mothers and adolescents. Genetic variations also impact the use of cannabis, in subjects with Val *COMT* allele there was a reduced functions of working memory, verbal and visual memory. Adult chronic cannabis users have reduced attention, cognitive control, poor memory, improper decision-making, motor performance and affective processing. Cortical gene expression and neurotransmission are also affected in rats due to delta-9-tetrahydrocannabinol. Visual-attention network and cerebellar hypoactivation is also noted in marijuana users.

## CONSENT AND ETHICAL APPROVAL

It is not applicable.

## COMPETING INTERESTS

Author has declared that no competing interests exist.

## REFERENCES

1. National institute on Drug Abuse. How does marijuana produce its effects?; 2019.
2. National institute on Drug Abuse. What are marijuana's long-term effects on the brain?; 2019.
3. Campolongo P, Trezza V, Cassano T, Gaetani S, Morgese MG, Ubaldi M, et al. Perinatal exposure to delta-9-tetrahydrocannabinol causes enduring cognitive deficits associated with alteration of cortical gene expression and neurotransmission in rats. *Addict Biol*. 2007;12(3-4):485-95.
4. Dalterio SL. Cannabinoid exposure: effects on development. *Neurobehav Toxicol Teratol*. 1986;8(4):345-52.
5. United Nations Office on Drugs and Crime. World drug report [cited Feb 10, 2021]. Available: <https://www.unodc.org/unodc/en/index.html>; 2015.
6. GBD 2016 Alcohol and Drug Use Collaborators. The global burden of disease attributable to alcohol and drug use in 195 countries and territories, 1990-2016: A systematic analysis for the global burden of disease study 2016. *Lancet Psychiatry*. 2018;5(12):987-1012. DOI: 10.1016/S2215-0366(18)30337-7
7. Hurd YL, Manzoni OJ, Pletnikov MV, Lee FS, Bhattacharyya S, Melis M. Cannabis and the developing brain: Insights into its long-lasting effects. *J Neurosci*. 2019;39(42):8250-8. DOI: 10.1523/JNEUROSCI.1165-19.2019
8. Rubino T, Realini N, Braidà D, Guidi S, Capurro V, Viganò D, et al. Changes in hippocampal morphology and neuroplasticity induced by adolescent THC treatment are associated with cognitive impairment in adulthood. *Hippocampus*. 2009;19(8):763-72. DOI: 10.1002/hipo.20554
9. Le AA, Quintanilla J, Amani MA, Piomelli D, Lynch G, Gall CM. Persistent sexually dimorphic effects of adolescent THC exposure on hippocampal synaptic plasticity and episodic memory in rodents. *Neurobiol Dis*. 2022;162(January):105565. DOI: 10.1016/j.nbd.2021.105565
10. Matochik JA, Eldreth DA, Cadet JL, Bolla KI. Altered brain tissue composition in heavy marijuana users. *Drug Alcohol Depend*. 2005;77(1):23-30. DOI: 10.1016/j.drugalcdep.2004.06.011
11. Glass M, Dragunow M, Faull RL. Cannabinoid receptors in the human brain: A detailed anatomical and quantitative autoradiographic study in the fetal, neonatal and adult human brain. *Neuroscience*. 1997;77(2):299-318. DOI: 10.1016/S0306-4522(96)00428-9
12. Kim DJ, Schnakenberg Martin AM, Shin YW, Jo HJ, Cheng H, Newman SD et al. Aberrant structural-functional coupling in adult cannabis users. *Hum Brain Mapp*. 2019;40(1):252-61. DOI: 10.1002/hbm.24369
13. Sagar KA, Gruber SA. Interactions between recreational cannabis use and cognitive function: Lessons from functional magnetic resonance imaging. *Ann N Y Acad Sci*. 2019;1451(1):42-70. DOI: 10.1111/nyas.13990
14. Chang L, Yakupov R, Cloak C, Ernst T. Marijuana use is associated with a reorganized visual-attention network and

- cerebellar hypoactivation. *Brain*. 2006;129 (5):1096-112.  
DOI: 10.1093/brain/awl064
15. Jager G, Kahn RS, Van Den Brink W, Van Ree JM, Ramsey NF. Long-term effects of frequent cannabis use on working memory and attention: An fMRI study. *Psychopharmacol (Berl)*. 2006; 185(3):358-68.  
DOI: 10.1007/s00213-005-0298-7
  16. Kanayama G, Rogowska J, Pope HG, Gruber SA, Yurgelun-Todd DA. Spatial working memory in heavy cannabis users: a functional magnetic resonance imaging study. *Psychopharmacol (Berl)*. 2004;176 (3-4):239-47.  
DOI: 10.1007/s00213-004-1885-8
  17. Bechara A, Tranel D, Damasio H. Characterization of the decision-making deficit of patients with ventromedial prefrontal cortex lesions. *Brain*. 2000;123 (11):2189-202.  
DOI: 10.1093/brain/123.11.2189
  18. Nestor L, Hester R, Garavan H. Increased ventral striatal bold activity during non-drug reward anticipation in cannabis users. *NeuroImage*. 2010;49(1):1133-43.  
DOI: 10.1016/j.neuroimage.2009.07.022
  19. Vaidya JG, Block RI, O'Leary DS, Ponto LB, Ghoneim MM, Bechara A. Effects of chronic marijuana use on brain activity during monetary decision-making. *Neuropsychopharmacology*. 2012;37(3): 618-29.  
DOI: 10.1038/npp.2011.227
  20. Wesley MJ, Hanlon CA, Porrino LJ. Poor decision-making by chronic marijuana users is associated with decreased functional responsiveness to negative consequences. *Psychiatry Res*. 2011;191 (1):51-9.  
DOI: 10.1016/j.psychresns.2010.10.002.
  21. King GR, Ernst T, Deng W, Stenger A, Gonzales RMK, Nakama H et al. Altered brain activation during visuomotor integration in chronic active cannabis users: relationship to cortisol levels. *J Neurosci*. 2011;31(49):17923-31.  
DOI: 10.1523/JNEUROSCI.4148-11.2011
  22. Gruber SA, Rogowska J, Yurgelun-Todd DA. Altered affective response in marijuana smokers: An FMRI study. *Drug Alcohol Depend*. 2009;105(1-2):139-53.  
DOI: 10.1016/j.drugalcdep.2009.06.019
  23. Bogaty SER, Lee RSC, Hickie IB, Hermens DF. Meta-analysis of neurocognition in young psychosis patients with current cannabis use. *J Psychiatr Res*. 2018;99:22-32.  
DOI: 10.1016/j.jpsychires.2018.01.010
  24. Batalla A, Bos J, Postma A, Bossong MG. The impact of cannabidiol on human brain function: A systematic review. *Front Pharmacol*. 2020;11:618184.  
DOI: 10.3389/fphar.2020.618184
  25. Coskera E, Schwitzerab T, Ramozcd N, Ligierae F, Lalanef L, Gorwoodcdg P et al. Laprévoteabh. The effect of interactions between genetics and cannabis use on neurocognition. A review. *Prog Neuropsychopharmacol Biol Psychiatry*. 2018;82:95-106.
  26. Liem-Moolenaar M, te Beek ET, de Kam ML, Fransen KL, Kahn RS, Hijman R et al. Central nervous system effects of haloperidol on THC in healthy male volunteers. *J Psychopharmacol*. 2010;24 (11):1697-708.  
DOI: 10.1177/0269881109358200
  27. Morrison PD, Zois V, McKeown DA, Lee TD, Holt DW, Powell JF et al. The acute effects of synthetic intravenous  $\Delta^9$ -tetrahydrocannabinol on psychosis, mood and cognitive functioning. *Psychol Med*. 2019;39(10).
  28. Ramaekers JG, Kauert G, van Ruitenbeek P, Theunissen EL, Schneider E, Moeller MR. High-potency marijuana impairs executive function and inhibitory motor control. *Neuropsychopharmacology*. 2006; 31(10):2296-303.  
DOI: 10.1038/sj.npp.1301068
  29. Filbey FM, Aslan S, Calhoun VD, Spence JS, Damaraju E, Caprihan A et al. Long-term effects of marijuana use on the brain. *Proc Natl Acad Sci USA*. 2014;111(47): 16913-8.  
DOI: 10.1073/pnas.1415297111
  30. Testai FD, Gorelick PB, Aparicio HJ, Filbey FM, Gonzalez R, Gottesman RF et al.; Council on Arteriosclerosis. Thrombosis and vascular biology; Council on Cardiovascular and Stroke Nursing. Stroke; and Council on Peripheral Vascular Disease. 2022;53(4):e176-87.  
DOI: 10.1161/STR.0000000000000396
  31. Meier MH, Caspi A, Ambler A, Harrington H, Houts R, Keefe RS et al. Persistent cannabis users show neuropsychological decline from childhood

- to midlife. *Proc Natl Acad Sci USA*. 2012;109(40):E2657-64.  
DOI: 10.1073/pnas.1206820109
32. National institute on Drug Abuse. Marijuana; 2019.
  33. Galli JA, Sawaya RA, FriedenberG FK. Cannabinoid hyperemesis syndrome. *Curr Drug Abuse Rev*. 2011;4(4):241-9.  
DOI: 10.2174/1874473711104040241
  34. Thomas G, Kloner RA, Rezkalla S. Adverse cardiovascular, cerebrovascular, and peripheral vascular effects of marijuana inhalation: What cardiologists need to know. *Am J Cardiol*. 2014;113(1):187-90.  
DOI: 10.1016/j.amjcard.2013.09.042
  35. Wolff V, Jouanjus E. Strokes are possible complications of cannabinoids use. *Epilepsy Behav*. 2017;70(B):355-63.  
DOI: 10.1016/j.yebeh.2017.01.031
  36. Thanvi BR, Treadwell SD. Cannabis and stroke: Is there a link? *Postgrad Med J*. 2009;85(1000):80-3.  
DOI: 10.1136/pgmj.2008.070425
  37. Garrett CP, Braithwaite RA, Teale JD. Unusual case of tetrahydrocannabinol intoxication confirmed by radioimmunoassay. *Br Med J*. 1977;2(6080):166.  
DOI: 10.1136/bmj.2.6080.166
  38. Lawston J, Borella A, Robinson JK, Whitaker-Azmitia PM. Changes in hippocampal morphology following chronic treatment with the synthetic cannabinoid WIN 55,212-2. *Brain Res*. 2000;877(2):407-10.  
DOI: 10.1016/s0006-8993(00)02739-6
  39. Jager G, Van Hell HH, De Win MML, Kahn RS, Van Den Brink W, Van Ree JM et al. Effects of frequent cannabis use on hippocampal activity during an associative memory task. *Eur Neuropsychopharmacol*. 2007;17(4):289-97.  
DOI: 10.1016/j.euroneuro.2006.10.003.
  40. Denissoff A, Mustonen A, Miettunen J, Alakokkare AE, Veijola J, Scott JG et al. Trajectories of adolescent psychotic-like experiences and early cannabis exposure: Results from a Finnish Birth Cohort Study. *Schizophr Res*. 2022;246:95-102.  
DOI: 10.1016/j.schres.2022.06.014
  41. Ma YL, Weston SE, Whalley BJ, Stephens GJ. The phytocannabinoid  $\Delta^9$ -tetrahydrocannabinol modulates inhibitory neurotransmission in the cerebellum. *British Journal of Pharmacology*. 2008;154(1):204-15.  
DOI: 10.1038/bjp.2008.57.
  42. Marinella MA. Stroke after marijuana smoking in a teenager with factor V Leiden mutation. *South Med J*. 2001;94(12):1217-8.
  43. Mateo I, Infante J, Gómez Beldarrain M, García-Moncó JC. Cannabis and cerebrovascular disease. *Neurologia*. 2006;21(4):204-8.
  44. Hasin DS, Saha TD, Kerridge BT, Goldstein RB, Chou SP, Zhang H, et al. Prevalence of marijuana use disorders in the United States between 2001-2002 and 2012-2013. *JAMA Psychiatry*. 2015;72(12):1235-42.  
DOI: 10.1001/jamapsychiatry.2015.1858