

The Baby Blues and Perinatal Psychic Disorders: About Prevention and Management

Abstract:

The baby blues is a precocious psychic impairment that may occur within the first two weeks following delivery. It is considered a trigger point, and sometimes a decompensation towards more serious postpartum psychiatric disorders. It is as well a hazard, predisposing to abnormal psycho-affective development in infants, otherwise intimately dependent on maternal bonding. Although there have been advances in understanding the psychopathology, the clinical presentations, and the inter-relationship of the condition with other disorders of perinatality, the management however remains unclear and poorly described. There seems to be confusion between indications for mere observation with monitoring on one hand, and the need for psychotherapy or pharmacotherapy on the other hand. This is maintained by the ambiguous distinction between primary baby blues which is milder, and severe baby blues with immediate and late complications that are sometimes neglected. Whereas, intervention may be necessary in a number of cases to prevent adverse outcomes in affected mothers and their infants. In this short paper, we review the management of baby blues according to the severity and we give clues for prevention, based on known protective factors.

Keywords: Baby blues, Perinatality, postpartum depression, psychosis

Introduction:

The effective prevention and management of the baby blues should take place and span throughout the duration of the pregnancy and beyond the immediate postpartum period. In effect, adverse events that are likely to impact the course of the pregnancy, the delivery process, and maternity in postpartum may occur very early after conception, and so should be avoided as much as possible to ease perinatality [1-6]. This process is known as the “carry-over phenomenon” and needs to be prevented through a number of measures. More than 80% of cases with baby blues may be described as primary blues with mild transient or benign presentation. This form of postpartum blues is closer to a normal physiological process than to a psychiatric disorder. However, 20% with more severe signs and symptoms may be real mental impairments, presenting as a “signal” or “trigger point” towards more serious psychiatric disorders of postpartum, and with related complications [1-6].

As a general rule, the primary baby blues to the sense of Kennerley and Gath is of precocious onset and fleeting, requiring only the mother's specific counseling, assistance, monitoring, and support from the entourage. It is such a mild condition that it may be unnoticed by the medical staff [1-6]. Whereas, the severe blues with atypically serious symptomatology and lasting for more than ten days necessitates psychiatric assessment, psychotherapy, and/or medication for treatment. This is because of the fact that severe baby blues are very likely to progress towards delivery-bound post-traumatic stress disorder, postpartum depression, or psychosis [1-6].

Documented protective factors against the onset of the baby blues include women having from average to high socioeconomic situation, a peaceful family, a good and satisfying couple relationship, benefiting adequate financial support from the partner, family, or friends, adequate emotional support from the partner, family or friends, being able to rely on the partner, family or friends when need be, having confidence in the partner and friends, having planned the pregnancy, desired and done prenatal check-up. These factors are thought to procure a strong psychosocial support network that is reassuring and psycho-prophylactic for mothers during the perinatal period [1-6].

As a matter of fact, besides the biological theory with hormonal imbalances of pregnancy justifying the blues, the psychodynamic theory of mental representations and maternal infantile conflicts have a central role in psychogenesis. Given the reluctance of physicians to intervene in physiological endocrinologic changes of perinatality, and for presumed mild affections such as the blues, the management relies mainly on the reinforcement of psychosocial support and/or psychotherapy. Moreover, the effective management of concomitant maternal or neonatal health issues including infections, injuries, and pains, that are likely to induce, maintain or worsen the blues through the summation effect, is important as well [5-12].

Prevention and Management

During pregnancy, antenatal visits should be observed by women with targeted counseling and information about the progress of pregnancy, the development of the fetus, preparation for delivery, and maternity [12-22]. This contributes to reassuring the mother and her partner, preventing worries and anxiety. It also helps to anticipate problems and challenges, so as to better manage them and avoid last-minute surprises [22-30]. In effect, in case a congenital birth defect or malformation is early diagnosed, parents' assistance with a psychologist is indispensable and should absolutely be recommended by the physician. This will facilitate gradual acceptance of the diagnosis, with the understanding of embryo-fetopathy, conditioning and coping, motivation to continue visits, and treatment observance [15-33].

In the course of the delivery process, trauma should be avoided as much as possible and labor facilitated through optimal obstetrical practices. Dystocic deliveries should be made less traumatic as possible with analgesia and effective anesthesia when a cesarean section is indicated. The presence of the husband or partner should be allowed in the delivery room and/or operating theatre when possible, for sympathetic assistance and mothers reassurance. Optimal obstetrical practices, just as safe delivery and effective analgesia have the double advantage of preventing pain-induced depression and enabling the mother to take care of the neonate immediately, following delivery [6-22].

After childbirth, postpartum blues management requires mothers' support with counseling and education on the benign and transient character of their physical and mental discomforts [1-6]. They should be reassured of the restoration of their body organs including uterine involution, cessation of lochia, wound healing, and weight loss, just to name some. This is cheering and recomforting for mothers and helps to break anxiety too. Furthermore, the medical staff should emphasize to the mother's entourage and relatives, her needs for enough rest and emotional support, assistance with baby care, balanced feeding with enough hydration, and breastfeeding practice and techniques [7-14]. Through such measures, mothers are relieved from a number of unnecessary preoccupations, so as to appease their minds and strengthen confidence, while enhancing optimism.

The mother's responsibilities as well as the necessity for good interactions with the baby are discussed with the medical staff. This helps mothers to understand the importance of the mechanisms of bonding establishment, and its "psycho-prophylactic" effects for both mother and infant, in the short and long term. Analgesia should be maintained or increased if necessary, and mothers should be thought or assisted with perineal care [7-22].

All these interventions may be complementary to cognitive and behavioral therapy, aimed at improving patients' thoughts and behaviors for a more comfortable psychological adaptation to the postpartum period. Family and relatives' support through sympathizing and reassurance help to calm down the mother, thereby favoring contending, with the additional benefit of not administering drugs [1-6].

After the mother's discharge from the hospital, positive interactions with loved ones should be maintained. In particular cultures such as in Africa, newly delivered women are fed with dishes that contain natural uterotonics and lactogenic galactagogues which are respectively responsible for uterine involution and increased breastmilk production [7-22]. Postnatal visits should be regularly observed with reassessment and

close follow-up of women with severe maternity blues to prevent postpartum depression. Nurse visits to newly delivered mothers' homes may serve a great purpose as well when proximity healthcare practices are possible. However, newly delivered women are often given the following recommendations before their discharge:

- To avoid extreme fatigue. Sleeping when the baby is sleeping. Allowing their partners or other support people to take care of the baby occasionally, so they can get extra rest
- Concentrate on themselves and the baby, letting others help with housework, laundry, cooking, shopping, and other stressful activities
- Get out of the house every day or when possible
- Go for a walk and/or meet a friend
- To treat themselves to something they like to do
- Share their feelings and frustrations with their partners or close friend.

They are also advised to notify their health care provider in case feelings of sadness, anxiety, resentment or guilt last longer than 2 weeks and interfere with their ability to care for themselves and/or the baby [34-44].

Given that severe maternity blues can easily progress towards post-partum depression and other psychic disorders of perinatality if poorly managed, newly delivered women are advised to keep in contact with their health care providers. They are also encouraged to contact their physician in case they experience any of the following symptoms of depression and anxiety disorders:

- Inability to sleep or sleeping all the time
- Loss of appetite
- Feeling down most of the time
- Feeling even the smallest task seems to take too much effort
- Feeling very critical of themselves or others
- Worrying constantly about the baby
- Having thoughts of harming themselves or the baby.

While psychotic manifestations will be reported by family members and partners if present. In such cases with baby blues complications, a mental reassessment should be done by the medical psychologist or the psychiatrist, and the need for intense psychotherapy or drug prescription appreciated.

The administration of progesterone in postpartum to prevent severe baby blues mental impairments is often recommended by most clinicians to quench progression

towards postpartum depression. This therapeutic intervention is based on the endocrinological theory of baby blues onset, with an estimated progesterone drop after delivery [34-44]. Following psychiatric reassessment, complications with postpartum depression which are the most frequent will be managed through sessions of psychotherapy as outpatients and the prescription of anti-depressants. Such medications mainly include serotonin reuptake inhibitors and mood or thymoregulators. In addition to this, treatment for any other medical issues related to depression will be administered as well. Although these mothers could be allowed to continue caring for their babies with breastfeeding, the drugs are known to be secreted in breastmilk, but in low doses that are harmless to the baby [7-14]. The treatment of postpartum depression may last up to six months so as to prevent chronic depression. Contrarily to postpartum depression, post-traumatic stress related to delivery is best managed through cognitive behavioral therapy consisting of debriefing, counseling, and positive interactions during a few weeks or months [34-44]. Generally, no medication is required. Bad habits such as doping and illicit psychotropic drug consumption including alcohol are prohibited, as they may worsen the condition.

In the case of postpartum psychosis, mothers should be separated from infants according to the severity of their condition. They may be hospitalized with physical or pharmacological contention in case of acute delirium with agitation or aggressiveness. Antipsychotic drugs are required in combination with mood stabilizers or thymoregulators and benzodiazepines to control signs and symptoms [34-44]. Breastfeeding is not possible because of mother-infant separation with decreased breastmilk secretion, and also due to antipsychotic drugs which are harmful to the baby. In the meantime, the baby can be fed with human donor milk or formula milk. The infant may be adopted and cared for by a family member, with the advantage of being safe from nosocomial infections, or it may be kept in the hospital's maternity and close to the mother, but with risk of hospital borne-infection.

Conclusion and recommendations

The psychic disorders of perinatal and postpartum are quite common, starting with the baby blues. When diagnosed, they should be further assessed by the mental health specialist for clinical distinction and precision. Most cases with primary blues and post-traumatic stress may uniquely be managed by cognitive behavioral therapy. While more serious conditions such as postpartum depression and psychosis respectively require antidepressants and antipsychotics in addition. Although the management of the later conditions may last for months, mother-infant bonding should be preserved as much as possible to prevent complications with the baby's psycho-affective development.

Conflict of Interest

The authors declare that they have no competing interests.

Acknowledgments

Hospitals authorities, all collaborators to this project.

References

1. Moyo GPK. Epidemio-clinical Profile of the Baby Blues in Cameroonian Women. *Journal of Family Medicine and Health Care*. 2020;6(1):20-23. doi: 10.11648/j.jfmhc.20200601.14
2. Moyo GPK, Djoda N. The Emotional Impact of Mode of Delivery in Cameroonian Mothers: Comparing Vaginal Delivery and Caesarean Section. *American Journal of Psychiatry and Neuroscience*. 2020;8(1): 22-25. doi: 10.11648/j.ajpn.20200801.15
3. Moyo GPK. Perinatal and Childbirth as a Factor of Decompensation of Mental Illness: The Case of Depressive States in Newly Delivered Cameroonian Women ABEB 2020: 4(4). ID.000592. DOI: 10.33552/ABEB.2020.04.000592
4. Moyo GPK, Djoda N. Relationship Between the Baby Blues and Postpartum Depression: A Study Among Cameroonian Women. *American Journal of Psychiatry and Neuroscience* 2020; 8(1): 26-29. Doi: 10.11648/j.ajpn.20200801.16
5. Foumane P, Olen JPK, Fouedjio JH, GPK Moyo, Nsahlai C, Mboudou E. Risk Factors of Maternity Blues after Caesarean Section. *Int J Reprod Contracept Obstet Gynecol*. 2016 Dec;5(12):4424-4427. DOI: <http://dx.doi.org/10.18203/2320-1770.ijrcog20164357>
6. Moyo GPK, Mendo RM, Batibonack C, Thérèse Mbang AT. Neonatal Determinants of Mothers' Affective Involvement in Newly Delivered Cameroonian Women. *Journal of Family Medicine and Health Care* 2020;6(2):125-128. doi: 10.11648/j.ajp.20200602.21
7. Hermann ND, Moyo GPK. Neonatal Determinants of Inadequate Breastfeeding: A Survey among a Group of Neonate Infants in Yaounde, Cameroon. *Open Access Library Journal* (2020); 7: e6541. DOI: 10.4236/oalib.1106541
8. Hermann ND, Georges Pius Kamsu Moyo GPK, Ejake L, Félicité N, Evelyn M, Linda M. Determinants of Breastfeeding Initiation Among Newly Delivered Women in Yaounde, Cameroon: a Cross-Sectional Survey *Health Sci Dis* 2020; 21:20-24. Doi: <https://www.hsd-fmsb.org/index.php/hsd/article/view/2265>
9. Moyo GPK, Dany Hermann ND. Clinical Characteristics of a Group of Cameroonian Neonates with Delayed Breastfeeding Initiation. *Am J Pediatr* 2020;6(3):292-295. doi: 10.11648/j.ajp.20200603.28
10. Moyo GPK, Hermann ND. The Psycho-Sociocultural Considerations of Breastfeeding in a Group of Cameroonian Women with Inadequate Practices. *J Psychiatry Psychiatric Disord* 2020; 4 (3): 130-138. doi: 10.26502/jppd.2572-519X0100
11. Moyo GPK. Breast Pathologies and Inadequate Breastfeeding Practices: A Survey Among a Group of Newly Delivered Women in Yaounde, Cameroon. *Journal of Family Medicine and Health Care*. 2020;6(3)87-90. doi: 10.11648/j.jfmhc.20200603.16
12. Moyo GPK, Ngwanou DH, Sap SNU, Nguefack F, Mah EM. The Pattern of Breastfeeding among a Group of Neonates in Yaoundé, Cameroon. *International Journal of Progressive Sciences and Technologies* 2020;22(1):61-66. DOI: 10.52155

13. Nguefack F, Ngwanou DH, Kamsu Moyo GP, Ejake L, Mah, EM, Wafeu G, Fodoung Wamba DS, Chiabi A. Reasons for Delayed Breastfeeding Initiation among Newly Delivered Women in Two First-Category Hospitals in Yaoundé, Cameroon. *Open J Pediatr* 2020;10:474-485. <https://doi.org/10.4236/ojped.2020.103048>
14. Moyo GPK, Hermann ND (2020) Caesarean Delivery as a Predictor of Inadequate Breastfeeding among a Group of Neonates in Yaoundé, Cameroon. *J Perina Clin Pediatr*: 2: 105. DOI: 10.29011/JPCP-105.100005
15. Moyo GPK*, Tidang CT, Zebaze S, Mendomo RM, Makowa LK, Mbang AT, Batibonak C, Essomba AA. Fetal and Neonatal Complications of Gestational Diabetes: A Survey in Two Referral Hospitals of Yaoundé, Cameroon *Pediatr Stud Care*, Volume 2(1): 1–4, 2022
16. Moyo GPK, Sobguemezing D, Adjifack HT. Neonatal Emergencies in Full-term Infants: A Seasonal Description in a Pediatric Referral Hospital of Yaoundé, Cameroon. *Am J Pediatr*. 2020;6(2):87-90. doi: 10.11648/j.ajp.20200602.13
17. Georges Pius K M, Aurore Albane E, Marie-Paul B, et al. Neonatal Sepsis: Highlights and Controversies. *J Pediatr Neonatal*. 2022; 4(1): 1-5. Doi: 10.33425/2689-1085.1035
18. Moyo GPK, Kuate L, Foe Mba M, Komba D, Tchougene AF, Bege MP, Tchinda CT, Awa HDM. Why Diazepam More than Other Benzodiazepines is Unsuitable for Neonates? *Pediatr Stud Care*, Volume 2(1): 1–3, 2022
19. Chiabi A, Kago DA, Moyo GPK, Obadeyi B. “Relevance and Applicability of the Apgar Score in Current Clinical Practice”. *EC Paediatrics* 2019;8(11):01-07.
20. Moyo GPK, Tetsiguia JRM. Discussing the “First Cry” as an Initial Assessment for Neonates. *Am J of Pediatr*. 2020;6(2): 129-132. doi: 10.11648/j.ajp.20200602.22
21. Moyo GPK, Um SSN, Awa HDM, et al. The pathophysiology of neonatal jaundice in urosepsis is complex with mixed bilirubin!!! *J Pediatr Neonatal Care*. 2022;12(2):68–70. DOI: 10.15406/jpnc.2022.12.00458
22. Moyo GPK, Nguedjam M, Miaffo L. Necrotizing Enterocolitis Complicating Sepsis in a Late Preterm Cameroonian Infant. *Am J Pediatr*. Vol. 6, No. 2, 2020, pp. 83-86. doi: 10.11648/j.ajp.20200602.12
23. Moyo GPK, Sap Ngo Um S, Awa HDM, Mbang TA, Virginie B, Makowa LK et al. An Atypical Case of Congenital and Neonatal Grave’s Disease. *Annal Cas Rep Rev: ACRR*-314. 2022;2. doi: 10.39127/2575-8253/IAPN:1000114
24. Moyo G P K, Um S S, Awa H B M, Kana C N, Makowa L K, Mbang T A, Virginie B, Ndombo P O K. Paroxysmal Dyskinesia, an unusual but Particular Presentation of Congenital and Neonatal Hyperthyroidism: A Case Report with Literature Review. *Clin Res Pediatr* 2022;4(1):06-11. DOI: 10.33309/2638-7654.040102
25. Ngwanou DH, Ngantchet E, Moyo GPK. Prune-Belly syndrome, a rare case presentation in neonatology: about one case in Yaounde, Cameroon. *Pan Afr Med J*. 2020;36:120. doi: 10.11604/pamj.2020.36.102.24062
26. Tague DAT, Evelyn Mah, Félicitee Nguefack, Georges Pius Kamsu Moyo, Lionel Loic Kago Tcheyanou, Faustin Mouafo. Beckwith-Wiedemann Syndrome: A Case Report at the Gynaeco-Obstetric and Pediatric Hospital in Yaounde, Cameroon. *Am J Pediatr* 2020;6(4):433-436. doi: 10.11648/j.ajp.20200604.17
27. Moyo GPK, Djomkam IFK. Epidemio-clinical Profile of Stunting in School Children of an Urban Community in Cameroon. *Am J Pediatr* 2020;6(2): 94-97. doi: 10.11648/j.ajp.20200602.15

28. Moyo GPK, Djomkam IFK. Factors Associated with Stunting in School Children of an Urban Community in Cameroon. *Am J Pediatr*. 2020;6(2):121-124. doi: 10.11648/j.ajp.20200602.20.
29. Moyo GPK, Ngapout OD, Makowa LK, Mbang AT, Binda V, Albane EA et al. Exogenous Cushing's Syndrome with Secondary Adrenal Insufficiency in an Asthmatic Infant: "Healing Evil with Evil". *Arch Pediatr* 2022; 7: 203. DOI: 10.29011/2575-825X.100203
30. Moyo GPK. Children and Adolescents' Violence: The Pattern and Determinants Beyond Psychological Theories. *Am J Pediatr* 2020;6(2):138-145. doi: 10.11648/j.ajp.20200602.24
31. Moyo GPK, Mbang AT, Makowa LK, Mendomo RM, Zebaze S, Awa HDM. Neurocysticercosis in a Child Living in the Urban Community of Yaoundé, Cameroon: A Case Report in a Low Resource Setting. *Am J Pediatr*. 2020; 6(2):91-93. doi: 10.11648/j.ajp.20200602.14
32. Essomba AA*, Zicfried K, Kouam JE, Edongue M, Nzotsa S, Engama MN et al. OEIS Complex: A Case Report and Literature Review of a Rare Polymalformative Syndrome. *Pediatr Res Child Health* 2022; 6(1)
33. Zicfried K*, Essomba AA, Makowa LK, Mendomo RM, Zebaze SZ, Mbang AT, Batibonak C, Ngapout OD. Cloacal Exstrophy: Documenting a Particular Case in Yaoundé, Cameroon. *J Pediatr Neonatal Biol* 2022;7(1): 69-72.
34. Kennerley H, Gath D. Maternity blues. *Br J Psychiatry* 1995; 153(5): 337-40
35. Kendell RF, Chalmers JC, Platz C. Epidemiology of puerperal psychoses. *Br J Psychiatry* 1987; 150(2): 662-673.
36. Pitt B. Maternity blues. *Br J Psychiatry* 1973; 122(569):431-3.
37. Davidson JTR. Postpartum mood changes in American women and a discussion on its significance. *Br J Psychiatry* 1972; 121(1): 659.
38. Handley SL, Dunn, T L Waldron and Baker. Tryptophan, cortisol and puerperal mood. *Br J Psychiatry*, 1980; 136(3): 496-508.
39. Kennerley H, Gath D. Maternity blues. *Br J Psychiatry* 1995; 153(5): 337-40.
40. O'Hara MW, Schlechte JA, Lewis DA, varner MW. Controlled prospective study of postpartum mood disorders: Psychological, environmental and hormonal variables. *J Abnorm Psychol* 1991;100(1):63-73
41. Zilborg G. The dynamic of schizophrenia reactions related to pregnancy and childbirth. *Am J Psychiatry* 1929; 85(20):733-741.
42. Oates M. Normal emotional changes in pregnancy and the puerperium. *Baillères Clin Obstet Gynaecol* 1989; 3(6):791-804.
43. Guedeney A, Bungener C, Widlöcher D. Le postpartum blues: une revue critique de la littérature. *Psychiatr Infant* 1993; 36(1): 329-354.
44. Adewuya A O. Prevalence and risk factors of maternity blues in western Nigerian women. *Am J Ostet Gynaecol*, 2005; 193(4): 1522-5.