

## TREES SPECIES DIVERSITY, ABUNDANCE AND SOIL PHYSICOCHEMICAL PROPERTIES OF UKPON RIVER FOREST RESERVES, CROSS RIVER, NIGERIA.

**Comment [H1]:** Title is same with Akendele et al., (2021). Author should not copy the same sentence or context.

### Abstract

Current status of species diversity, composition and abundance provides guidance for their management and assessment of their ecological usefulness. In this study trees species diversity, abundance and soil properties of Ukpon River Forest Reserve was assessed. Line transect method was using to establish 4 sample plots of 50 x 50 m. Data on vegetation were collected using the appropriate tool and soil samples from the plot were collected with the aid of soil auger. Tree species composition, abundance and diversity indices were estimated using the appropriate formulae and soil samples were analyzed following the by using standard methods. A total of 194 individuals in 60 species belonging to 26 families were encountered in the study area. The dominant families are Leguminaceae, Moraceae, Fabaceae, Burseraceae, Apocynaceae, Calsalpinaceae and Euphorbiaceae. The total basal area (tBA) estimated for tree species was 147.615m<sup>2</sup> with *Bombax bounpozen* having the highest of 11.09m<sup>2</sup> and relative dominance (RDo) of 7.51% while *Piptadeniastrum africana* has the highest relative density (RD) of 7.73% and importance value index (IMI) of 15.63%. Based on their relative density, 68.34% of the trees were Rare, 18.33% Threatened or Endangered, 5% Abundant/ Occasional and 3.33% Frequent. The study had a high Shannon – Weiner index value of 3.04 and Margalef richness of 20.67 and low dominance index of 0.06. The soil properties such as Clay, Organic matter, Phosphorus, Calcium, Cation Exchangeable capacity and Base Saturation were high. The pH value of 5.77 shows the soil were moderately acidic. These properties have been shown to improve soil fertility status and moisture content needed plant growth. Although the tree species diversity in the study area was high, some species appears to have been threatened while majority were rare, sustainable conservation effort should be geared towards ensuring their continuous existence.

**Formatted:** Superscript

Keywords: Tree species, Abundance, Diversity Indices, Soil Properties, Forest Reserve

### Introduction

The tropical rainforest has been identified as the most biologically diverse terrestrial ecosystem on earth (FAO 2010, and IUCN 2010). In terms of tree composition and species diversity, tropical rain forests are Earth's most complex ecosystems (Gebreselasse, 2011). This high species diversity is partly responsible for the intense pressure under which rainforests are subjected to the effects of forest degradation and deforestation. Forest degradation is usually accompanied by species extinction, reduction in biodiversity, and decrease in primary productivity. About ten million hectares of rainforests are degraded annually worldwide and according to WRM (1999) 200,000 hectares of rainforest in Cameroon and Nigeria are degraded annually, with over 40 trees and wildlife species being threatened with extinction. The Nigerian rainforest ecosystems occupy only 9.7% of the country's land mass of 983,213 km<sup>2</sup> but it is the most densely populated and source of the bulk of Nigeria's timber (FAO 2010). The acceptance of Nigerian timber species in the international market in the early 20th century, coupled with rising domestic demand, led to the

**Comment [H2]:** Better to rewrite sentences marked with yellow highlight

**Formatted:** Highlight

**Comment [H3]:** Add reference to the statement.

situation where exploitations became unregulated. This led to serious forest degradation that left less than 5% of the country's rainforest ecosystems as undisturbed (Gillespie *et al.*, 2004). Recent global forest resources assessment revealed that Nigeria is one of the five countries in the world with the highest annual rate of deforestation for the period 2000 –2010 (FRA, 2010). According to Adekunle *et al.*, (2010), a total of 111,377 timber stems, belonging to 62 different indigenous hardwood species of tropical rainforest ecosystem, distributed among 16 families, were exploited from Ondo State forest ecosystem between 2003 and 2005. Maliyat and Datt (2010) reported that the expansion of biotic activities is also responsible for overexploitation of natural resources, and this has subsequently disturbed the delicate equilibrium that exists between living organisms and their environment also disturb the ecological balance of a community. Many rainforests are severely threatened and persist as forest fragments leading to a reduction in species richness, thereby resulting in biodiversity loss and establishment of Forest Reserves. Forest Reserve is one of the in-situ methods of conservation that are required to restrict human activities and degradation of the forest. The process of biodiversity in protected areas begins with the assessment of the status of organism's species composition and abundance, and a According to USAID/ Nigeria (2008) the problem is lack of adequate documented data on the status in the country. These changes have caused the loss of some plant species and a decline in the biodiversity conservation status of the forest and environmental quality.

Formatted: Highlight

Comment [H4]: What does it mean?  
Anthropogenic activities?

Formatted: Highlight

Comment [H5]: Follow the sequence of writing.  
May reove the sentence.

Due to unprecedented increase in human population over the years, trees species have undergone different levels of disturbance which has adversely impacted on their abundance, composition, diversity and conservation (Omoro *et al.*, 2010). Trees species Species composition and abundance will provide guidance for their management and valuable reference for assessment as well as improve our knowledge in identification of ecologically useful species (Suratman, 2012). Higher numbers of species increases the number of ecological niches as well as the numbers of associated species (Kanowski *et al.*, 2003). According to Adeyemi *et al.* (2015) a total of 125 tree species belonging to 36 families and 96 genera were recorded in the area with Margalef's index of species richness of 2.2754 and almost (99) of the tree species encountered were threatened/endangered. High trees species composition in Okwango forest, Cross River, Nigeria with dominant species to include *Terminalia ivorensis*, *Pterocarpus soyauxii*, *Melicia excelsa*, *Bailonella toxisperma* and *Afzelia bipindensis*. Edet *et al.* (2011) reported a total of 102 species belonging to 35 families in Afi Mountain Wildlife Sanctuary. Most tropical forest is characterized by abundance of trees with small diameter at breath high (dbh). Jimoh *et al* (2012)

noted that Oban Division of Cross River National Park was characterized by dominance of tree species in lower diameter classes. Ogwu *et al.* (2016) recorded ~~a total of~~ 214 individual ~~of~~ 20 trees species from 12 families with Arecaceae and Fabaceae as the most abundant in Ugbowo, Benin City, Nigeria. Two hundred and ten (210) trees distributed among 54 species in 25 families with Fabaceae and Euphorbiaceae as the most dominant enumerated in Akure Forest Reserve, Nigeria (Akindele *et al.*, 2021). Iheyen *et al.* (2009) reported Fabaceae family as the most abundant in Ehor Forest Reserve.

The relationships between vegetation and fertility are closely linked to the quality and quantity of litter and soil organic matter ~~quality and quantity~~ (Maro *et al.*, 1991). According to Van Bremen (1993), organisms appear to affect soil fertility, soil moisture content and other soil properties in such a way that ~~with time~~ the substrate becomes more favorable for the growth of plants and soil organisms with time. Availability of Soil moisture ~~availability has been reported~~ to be the main factors affecting habitat associations of tropical trees, shrubs and herbs (Chaturvedi and Raghubanshi 2018). Studies on the relationship between species diversity and ~~soil~~ physicochemical properties of soil are the basis of community ecology and biodiversity conservation and management (Peng *et al.*, 2015). Nirmat-Kumar *et al.*, (2010) investigated nutrients content in different sites of an Indian rainforest and found that there was a strong positive relationship between tree species richness and the concentration of nitrogen, phosphorus and organic carbon. In Cross River, past researches on Forest Reserves have focus more on assessing tree species diversity, abundance and composition ignoring its relationship with soil properties (Edet *et al.*, 2012; Jimoh *et al.*, 2012; Adeyemi *et al.*, 2015). Akindele *et al.*, (2021) stated that Potassium, Cation Exchangeable Capacity, Sodium, Phosphorus, Clay and pH were fundamental soil properties that determines the quality and diversity of trees species in Akure Forest Reserve.

Trees provide many ecosystem services such as species conservation, prevention of soil erosion and preservation of habitat for plants and animals (Armenteras et al 2009). Overexploitation has led to rapid loss of trees diversity which is the major environmental and economic problem worldwide (Mani and Parthasarathy, 2006). Sustainable management and use of forest resources is essential for the nation's economic and environmental security (Akinsanmi, 1999). The need to provide adequate quantitative and qualitative ecological data to guide forest owners and managers in fashioning out realistic and effective management strategies is imperative. There is still

Formatted: Highlight


Comment [H6]: Not clearly understood. Rewrite the sentence.

insufficient information on trees species diversity, abundance and the physicochemical properties of Ukpon River Forest Reserve. Therefore, this study is aimed to determine the trees species diversity, abundance and physicochemical status of Ukpon River Forest Reserve.

## Materials and Methods

### Study Area

The study was carried out at the Ukpon River Forest Reserve located on latitude  $5^{\circ} 78' - 5^{\circ} 46' N$  and longitude  $8^{\circ} 32' - 8^{\circ} 19' E$ . The Reserve forest which is located in Obubra L.G.A has a total area of 12,950 hectares and is managed by the Cross River State Forestry Commission. The mean annual rainfall ranges from 2400 – 4000mm and is fairly distributed throughout the months of April to October with an annual temperature range of 27.6 - 33.1°C. The Reserve forest lies within the lowland rainforest with fresh water swamp at the fringes of Ukpon River and Derived Savannah north of the Reserve with heterogeneous floristic composition. The topography varies from undulating land on the south and southwestern parts to rugged and hilly land on the North and Northeastern part of the Forest reserve. The principal occupation of the communities around the Forest Reserve is farming, hunting and lumbering.



**Figure 1: Map of Cross River State showing the location of the Forest Reserve**

**Comment [H7]:** Image should be edited properly. It looks more shrunk. Correct it in copy editing time.

### Method of sampling and data collection

A random sampling Techniques was used to established 4 Temporary Sample plots of 50m x 50m in using a systematic line transect method according to Akindele *et al.* (2021). Global Positioning System (GPS) was used to locate the sample plots.

**Comment [H8]:** Period of sampling and number of sampling per year is missing. Please provide the details.

All trees species on each sample plot were tagged, measured, identifies and classified into families. The number and scientific names of all the tree species encountered in each plot were recorded. Leaves of trees not identified on the field were collected and taken to the herbarium for proper identification using botanical identification manual (Okpeke, 1987). Tree growth variables such as diameter at breast height (DBH)  $\geq 10$  cm and the total height was measured using Spiegel relaskop.

Diameter at breast height (DBH) and Height (H) were used to calculate Basal Area, Frequency, Relative Frequency, Relative Density, Relative Dominance, and Importance Value Index (IVI)

**Comment [H9]:** Mention all the sources for using formula below.

$$\text{Basal Area (BA)} = \frac{\pi D^2}{4} \dots\dots\dots (1)$$

Where D = Diameter at breast height (cm) and  $\pi$  = Pie (3.142)

$$\text{Where; Frequency} = \frac{\text{No of quadrat in which species occurred}}{\text{Total No of quadrat studied}} \times 100 \dots\dots\dots (2)$$

$$\text{Relative Frequency} = \frac{\text{Frequency of occurrence of species}}{\text{Total frequency of occurrence of all species}} \times 100 \dots\dots\dots (3)$$

$$\text{Relative density} = \frac{\text{No of individual species}}{\text{Total No of individual of all species}} \times 100 \dots\dots\dots (4)$$

$$\text{Relative Dominance} = \frac{\text{Total basal area of a species}}{\text{Total basal area of all species}} \times 100 \dots\dots\dots (5)$$

**Comment [H10]:** Don't copy and paste here. Authors must mention all the sources of this formula.

$$\text{Relative abundance} = \frac{\text{Relative Density (RD)}}{100} \dots\dots\dots (6)$$

The various trees species were classified according to their relative densities (RD) as abundant (RD ≥ 5.00), Frequent (4.00 ≤ RD ≤ 4.99), occasional (3.00 ≤ RD ≤ 3.99), rare (1.00 ≤ RD ≤ 2.99) and threatened (0.00 < RD ≤ 1.00) according to Ogwu et al (2016)

$$\text{Importance Value Index (IVI)} = \frac{\text{RD} + \text{RF} + \text{RDo}}{3} \dots\dots\dots (7)$$

Where RD = Relative Density; RF = Relative Frequency, RDo = Relative Dominance (Panwar and Bhardwaj, 2012)

Species diversity index was calculated using Shannon-Weiner diversity index

$$H' = - \sum_{i=1}^S P_i \ln(P_i) \dots\dots\dots (8)$$

Comment [H11]: Follow the above comment

Where H' = the Shannon-Wiener index

p<sub>i</sub> = the proportion of individuals belonging to species i

ln = the natural log (i.e., 2.718)

Margalef species richness index (D) used to measure the species richness according to Margalef (1948)

$$D = \frac{(S-1)}{\ln N} \dots\dots\dots (9)$$

Where S = Number of species and N = Number of individuals

The species Evenness (E<sub>H</sub>): Shannon's equitability equation was adopted to obtain the species evenness in each plot (Kent and Coker, 1992).

$$E_H = \frac{H'}{H_{Max}} = \frac{\sum_{i=1}^S P_i \ln(P_i)}{\ln(S)} \dots\dots\dots (10)$$

Comment [H12]: Don't copy, write in running letter

### Physicochemical Properties Analysis

Soil samples at the depths of 0 – 15cm, 15 – 30cm and 30 – 45cm were collected along the diagonal using the soil auger. The samples collected within each sample plot and depth were bulked together and air dried, sieved with a 2mm net and transferred to the laboratory for physicochemical analyses. The selected soil properties investigated were sand, clay, silt, phosphorus, calcium, magnesium, potassium, aluminum, sodium, pH, total nitrogen, organic carbon, organic matter, exchangeable cation capacity (ECC) exchangeable acidity (EA), electric

Comment [H13]: Exactly copied from Akindede et al., 2021

conductivity (EC), base saturation (BS) and bulk density. The chemical properties were analyzed using various standard laboratory methods. Total nitrogen was determined using micro Kjeldahl apparatus according to Martins et al. (2015). Sodium (Na) and potassium (K) concentrations were done using the flame photometer while calcium (Ca) and magnesium (Mg) were determined using the Atomic absorption spectrophotometer (AAS). Soil pH was determined using glass electrode pH meter while conductivity will be determined following the procedure of Clay et al (2012).

### Statistical data analysis

Data obtained from the soil physicochemical properties were subjected to descriptive statistics to compute for means and standard deviation of each parameter. The analyses were computed using statistical package for social sciences (SPSS) version 20.

**Comment [H14]:** Author must calculate ANOVA and multiple regression (R value) data to correlate the changes in diversity and abundance with physico-chemical parameters

### Results

The result presented in Table 1 shows the list of identified tree species and classification into their families in the study area. A total of 194 individual trees distributed among 62 different tree species belonging to 26 families were identified documented during the study. The family leguminaceae shows dominance with 11 tree species followed by Moraceae and Fabaceae with 7 and 6 species respectively. The family Burseraceae had 4 while Apocynaceae, calsalpinaceae and Euphobiaceae recorded 3 tree species each. Other such as the Bombacaceae, Flacourtaceae ,Myristicaceae, Sapindaceae and Sapotaceae has 2 tree species while Anisophilleacea, Cercropaceae , Compretaceae, Malvaceae, Saccoglobiaceae and Ulmaceae have 1 tree species each. The family Leguminaceae had the highest individuals with a total number of 44 followed by Fabaceae, 31 and Moraceae 24.

**Table 1: Families and their number of species and individualFamily wise distribution of plant species-trees within the study area.**

Family	Number tree species	Number of individual trees
<i>Anisophylleacea</i>	1	1
<i>Apocynaceae</i>	3	11
<i>Bombacaceae</i>	2	6

<i>Burseraceae</i>	4	6
<i>Caesalpiniaceae</i>	3	12
<i>Cluciaceae</i>	1	2
<i>Cercropiaceae</i>	1	2
<i>Compretaceae</i>	1	3
<i>Eupobiaceae</i>	3	14
<i>Fabaccae</i>	6	31
<i>Flacourtiaceae</i>	2	4
<i>Irvingiaceae</i>	2	4
<i>Leguminoeseae</i>	11	44
<i>Meliaceae</i>	1	1
<i>Malvaceae</i>	1	1
<i>Merbebaseae</i>	1	2
<i>Moraceae</i>	7	24
<i>Myristicacene</i>	2	4
<i>Rutiaceae</i>	1	5
<i>Rubiaceae</i>	1	1
<i>Sapindaceae</i>	2	4
<i>Saccogobiaceae</i>	1	1
<i>Sapotaceae</i>	2	4
<i>Sterculiaceae</i>	1	4
<i>Simaroubaceae</i>	1	1
<i>Ulmaceae</i>	1	3
<b>Total</b>	<b>26</b>	<b>194</b>

The ~~result in Table 2 represents~~ result in Table 2 represents the trees species and their basal area, relative densities (RD), relative dominance (RDo), relative abundance (RA) and importance value index (IVI). The basal area ranged from 0.031 -11.089 m<sup>2</sup> for diverse species of trees in the study area, the total basal area of tree species recorded was 147.615m<sup>2</sup>. *Bombax bounopozense*, belonging to the family *Bombacaceae* recorded the highest basal area of 11.089m<sup>2</sup> followed by *Ricinodendron heudelotii*, 8.301m<sup>2</sup> and *Piptadeniastrum africana*, 7.318m<sup>2</sup>. Other include *Hylodendron garbonenses* with 7.104m<sup>2</sup>, *Pentaclethra macrophylla* had 7.054m<sup>2</sup> while *Triplochiton scleroxylon* recorded the lowest basal area of 0.031m<sup>2</sup>. The relative density ranged from 0.05-7.73% with *P. africana* which belong to the family *Fabaceae* recorded the highest density of 7.73% while, *C. gabonensis*, *S. gabonensis*, *T. triplochitin* and *L. trichiloides* recorded the lowest of 0.05% each. The relative dominance varied between 0.02 (*Triplochiton scleroxylum*) and 7.51 (*Bombax bounopozens*). The highest relative abundance of 0.08 was recorded in *P. africana* followed by *R. heudelotii* with 0.06 while, *C. gabonensis*, *S. gabonensis*, *T. triplochitin* and *L. trichiloides* had the lowest value of 0.001 each. Importance value index (IVI) shows that, *Piptadeniastrum africana* has the highest (IVI) of 15.65 followed closely by *Ricinodendron*

*heudelotii* which recorded 13.99 while *Tryplochiton scleroxylon* has the least dominant value of 0.85 in the study area. The IVI also shows that *P. africana*, *R. heudelotii*, *B. bounopozens*, *A. ptercarpoides* and *P. macrophylla* were the most dominant tree species with IVI value above 10.00.

**Table 2: Tree species, their basal area, relative density, dominance, abundance and importance value index in the study area.**

S/N	Species	No of individual	Basal Area (Mm <sup>2</sup> )	Relative density (RD)	Relative dominance (RD <sub>0</sub> )	Relative Frequency (RF)	Relative Abundance (RA)	Importance Value Index (IVI)	Classification based on (RD)
1	<i>Piptadeniastrum africana</i>	15	7.318	7.73	4.96	3.13	0.08	15.63	Abundant
2	<i>Ricinodemdron heudelotii</i>	10	8.301	5.15	5.62	3.13	0.06	13.99	Abundant
3	<i>Bosqueia angolensis</i>	10	1.368	5.15	0.93	3.13	0.03	9.3	Abundant
4	<i>Gosweilorodendron balsalmiformis</i>	5	6.440	2.58	4.36	2.34	0.02	9.32	Rare
5	<i>Blighia sapida</i>	3	2.910	1.55	1.97	3.34	0.03	5.88	Rare
6	<i>Daniella ogea</i>	3	3.309	1.55	2.24	1.56	0.02	5.37	Rare
7	<i>Bombax bounopozens</i>	4	11.087	2.06	7.51	3.13	0.02	12.73	Rare
8	<i>Baphia gracilypis</i>	8	3.462	4.12	2.35	3.13	0.04	9.67	Frequent
9	<i>Picalima nitida</i>	6	2.985	3.09	2.02	3.13	0.03	8.29	Occasional
10	<i>Allanblankia floribunda</i>	2	1.040	1.03	0.7	0.78	0.01	2.53	Rare
11	<i>Terminalia superba</i>	3	5.782	1.55	3.91	3.13	0.03	8.61	Rare
12	<i>Alstoea boonei</i>	3	2.772	1.55	1.88	2.34	0.02	5.79	Rare
13	<i>Nuclea diderachi</i>	2	1.229	1.03	0.83	0.78	0.01	2.66	Rare
14	<i>Antiaris africana</i>	2	0.622	1.03	0.42	1.56	0.01	3.03	Rare
15	<i>Stautia stipitata</i>	2	1.232	1.03	0.83	1.56	0.01	3.44	Rare
16	<i>Mitrigyna cyliata</i>	2	1.213	1.03	0.82	1.56	0.01	3.43	Rare
17	<i>Mammea africana</i>	4	3.157	2.06	2.14	2.34	0.02	6.57	Rare
18	<i>Hylodendrom gabonensis</i>	4	7.104	2.06	4.81	1.56	0.02	8.46	Rare
19	<i>Celtis philipense (spp)</i>	3	1.901	1.55	1.29	1.56	0.02	4.42	Rare
20	<i>Vitex doniana</i>	2	0.984	1.03	0.67	1.56	0.01	3.28	Rare
21	<i>Lanmea welwiitchii</i>	1	0.245	0.52	0.17	0.78	0.01	1.47	Threatened
22	<i>Sterculia oblonga</i>	4	0.188	2.06	0.13	1.56	0.02	3.78	Rare

23	<i>Amphimas ptericarpoides</i>	9	3.880	4.64	2.63	3.13	0.05	10.47	Frequent
24	<i>Albizia ferruginea(Spp)</i>	3	1.756	1.55	1.19	2.34	0.02	5.1	Rare
25	<i>Musanga cercropioides</i>	2	6.284	1.03	4.26	1.56	0.01	6.87	Rare
26	<i>Fagara Macrophylla</i>	5	2.454	2.58	1.66	3.13	0.03	7.41	Rare
27	<i>Cryptosepalum pellegriniamin</i>	3	2.837	1.55	1.92	1.56	0.02	5.05	Rare
28	<i>Treulia affricana</i>	3	1.339	1.55	0.91	1.56	0.02	4.04	Rare
29	<i>Dedelota africana</i>	2	0.490	1.03	0.33	1.56	0.02	2.94	Rare
30	<i>Dialum guniensis</i>	1	0.113	0.52	0.08	0.78	0.01	1.38	Threatened
31	<i>Pycnanthus angolensis</i>	2	1.961	1.03	1.33	0.78	0.01	3.16	Rare
32	<i>Deterium microcarpum</i>	6	1.087	3.09	1.74	2.34	0.03	7.22	Occasional
33	<i>Brachystegia eurycoma</i>	3	5.574	1.55	3.78	1.56	0.02	6.91	Rare
34	<i>Azelia Africana</i>	3	1.138	1.55	0.78	1.56	0.02	3.91	Rare
35	<i>Belinia auriculata</i>	2	0.313	1.03	0.21	1.56	0.01	2.82	Rare
36	<i>Anthonatha macrophylla</i>	5	2.781	2.58	1.88	2.34	0.02	6.84	Rare
37	<i>pentaclethra macrophylla</i>	6	7.054	3.09	4.78	2.34	0.03	10.26	Occasional
38	<i>Gambeya albida</i>	2	3.036	1.03	2.06	1.56	0.01	4.67	Rare
39	<i>Pterocarpus osun</i>	3	4.123	1.55	2.79	2.34	0.02	6.7	Rare
40	<i>Eriocolum macrocarpum</i>	1	0.044	0.52	0.03	0.78	0.01	1.33	Threatened
41	<i>Dacryoides edulis</i>	2	0.283	1.03	0.19	1.56	0.01	2.8	Rare
42	<i>Irvingia garbonensis</i>	3	1.998	1.55	1.35	1.56	0.02	4.48	Rare
43	<i>Garcinia cola</i>	1	0.987	0.52	0.67	0.78	0.01	1.97	Threatened
44	<i>Anisophyllia purpurasens</i>	1	1.910	0.52	1.21	0.78	0.01	2.51	Threatened
45	<i>Homalium letistus</i>	3	0.588	1.55	1.4	0.78	0.02	3.75	Rare
46	<i>Millettia mecrophylla</i>	2	0.666	1.03	0.45	1.56	0.01	3.06	Rare
47	<i>Lophyria alata</i>	2	2.316	1.03	1.57	1.56	0.01	4.18	Rare
48	<i>Poga Oleosa</i>	2	2.542	1.03	1.76	1.56	0.01	4.37	Rare
49	<i>Ceiba pentandra</i>	2	1.913	1.03	1.3	1.56	0.01	3.91	Rare
50	<i>Parkia bicolor</i>	2	1.948	1.03	1.32	1.56	0.01	3.93	Rare
51	<i>Millicia etelsa</i>	3	3.142	1.55	2.13	0.78	0.02	3.43	Rare
52	<i>Distemanthus bentamianus</i>	2	1.791	1.03	1.21	0.78	0.01	3.04	Rare
53	<i>Discordis cliamiaper</i>	1	1.530	0.05	1.04	0.78	0.001	1.87	Threatened
54	<i>Santira trimera</i>	2	0.952	1.03	1.64	0.78	0.01	3.47	Rare
55	<i>Honoa Klaineana</i>	1	0.386	0.05	0.26	0.78	0.001	1.09	Threatened
56	<i>Cylincodiscus gabonensis</i>	1	0.987	0.05	0.67	0.78	0.001	1.5	Threatened

57	<i>Parinaris Africana</i>	2	2.111	1.03	1.43	0.78	0.01	2.26	Rare
58	<i>Sacoglotis gabonensis</i>	1	0.364	0.05	0.25	0.78	0.001	1.08	Threatened
59	<i>Triplochiton scleroxylon</i>	1	0.031	0.05	0.02	0.78	0.001	0.85	Threatened
60	<i>Lovoa trichiloides</i>	1	0.257	0.05	0.17	0.78	0.001	1	Threatened
<b>Total</b>		<b>194</b>	<b>147.615</b>	<b>100.0</b>	<b>100.00</b>	<b>100.00</b>	<b>1.07</b>	<b>300.00</b>	

The result of the tree growth variables and biodiversity indices as presented in table 3 revealed that in 26 families with 62 tree species and 194 individual trees their basal area was 147.62m<sup>3</sup>. The result also showed that only 5% of the tree species were categorize as Abundant and Occasional. Majority of the trees representing 68.34% were Rare species and 18.33% were either threatened or endangered species while 3.33% of the trees were Frequent species.

**Table 3: Summary of tree growth variables and biodiversity indices of the study area.**

Growth variables and Biodiversity indices	Values
Number of families	26
Number of species	62
Number of trees	194
Total basal area (m <sup>3</sup> )	147.62
Abundant (Relative density; RD ≥ 5)	3 (5%)
Frequent (4.00 ≤ RD ≤ 4.99)	2 (3.33%)
Occasional (3.00 ≤ RD ≤ 3.99)	3 (5%)
Rare (1.00 ≤ RD ≤ 2.99)	41 (68.33%)
Threatened/ Endangered (0.00 < RD ≤ 1)	11 (18.33%)
Shannon – wiener index (H)	3.04
Margalef Species Richness (M)	20.67
Species Evenness (E <sub>H</sub> )	2.19
Index of Dominance	0.06

The result of the physicochemical properties of soil of Ukpon River Forest is presented in Table 4. The properties showed 30.46% sand, 60.16% clay and 9.30% silt. The soil is slightly acidic (5.77) with higher available phosphorus (25.05 Cmol/kg) followed by Calcium (17.15 Cmol/kg) with the lowest value of sodium (0.12 Cmol/kg). The result also recorded a base saturation level of 91.41%, organic matter of 4.32% and a bulk density of 1.62 g/cm<sup>3</sup>.

Table 4: The mean values of the physicochemical properties of soil in the study area

Parameter	Mean	Standard deviation
Sand (%)	30.46	7.66
Silt (%)	9.30	2.34
Clay (%)	60.16	5.62
pH	5.77	1.77
Electric conductivity ( $\mu\text{S}/\text{cm}$ )	0.23	0.19
Organic matter (%)	4.32	1.99
Total nitrogen (%)	0.11	0.05
Phosphorus (Cmol/kg)	25.05	8.99
Calcium (Cmol/kg)	17.15	4.27
Magnesium (Cmol/kg)	5.79	1.49
Sodium (Cmol/kg)	0.12	0.04
Potassium (Cmol/kg)	0.20	0.11
Aluminum (Cmol/kg)	0.71	0.33
Effective Cation Exchange Capacity (Cmol/kg)	26.41	5.24
Exchange acidity (Cmol/kg)	2.24	0.89
Base saturation (%)	91.41	4.72
Bulk density ( $\text{g}/\text{cm}^3$ )	1.62	0.37

## Discussion

Forest play important role in maintaining fundamental ecological processes by providing livelihood and supporting economic growth. There is a growing concern for developing new global, regional and national programme for conserving and managing forest biodiversity. The creation of protected areas is considered the most effective means to stop and reverse degradation of our forest (Jekins and Joppa, 2009). The result obtained from this study show that trees are diverse in Ukpon River Forest Reserve of Cross River, Nigeria. A total of 194 individual trees in 62 species belonging to 26 families encountered in this study was higher than the 20 tree species in 12 families encountered by Ogwu et al. (2016) in University of Benin campus and 54 species in 25 families reported by Akindele et al (2021) in Akure Forest Reserve. Ikaagba et al. (2015) and Edu, (2017) have both reported a lower trees species of 52 in University of Agriculture, Makurdi and selected forest reserves in cross river respectively. The slight high in population recorded in the study area may be attributed to favourable microclimate within the forest and availability of viable seeds of trees to sustain regeneration. However, Aigbe and Omokhua (2015) recorded a much higher tree species 72 in 30 families in Oban Forest Reserve of Cross River. The dominance

**Comment [H15]:** Author discuss correlation of soil physic-chemical parameters with tree species diversity, but the present study results does not show any data on the correlation and significant variation in between diversity and physic chemical parameters. Please justify the reason.

of the tree species of the families *Leguminaceae*, *Moraceae*, *Meliaceae* and *Fabaceae* in this study is in line with those of Edu (2017) and Edet *et al.*, (2012) at Ukpon River and Afi Forest Reserves respectively. Akindele *et al* (2021) also reported these families as dominant except Leguminaceae which was not encountered in Akure Forest Reserve. The representation of most families by more than (5) species is typical of the rainforest vegetation of Nigeria (Edet *et al.*, 2012; Aigbe and Omokhua 2014). Ihenyen *et al.* (2009) and Akindele *et al* (2021) reported the Fabaceae family as the most abundant in Ehor and Akure Forest Reserves respectively as against Leguminaceae reported in this study. Dominance of a family may be attributed to their efficiency in seed dispersal mechanism and soil fertility status (Udo *et al.*, 2007). However, some families such as *Ebenaceae* and *Malvaceae*, had less than (3) species each which is also typical of West African families (Abbey 2006; Richard 2007). This low species representation in these families could be due to poor regeneration abilities and/or anthropogenic activities (Zhigila *et al.*, 2016). *Piptadeniastrum africana* (15.63%) of the family Leguminaceae and *Ricinodendron heudeltii* (13.99%) of the family Euphorbeaceae had the highest importance value indices. Akindele *et al.* (2021) also reported *Ricinodendron heudeltii* (10.49%) as the second highest after *Celtis zenkeri* (15.49%) of the family Ulmaceae. According to Zhigila *et al.* (2016) high importance value index of a species indicated its dominance and ecological success, its good power of regeneration and greater ecological amplitude. Some such as *Triplochiton scleroxylon* (0.85%), *Lovoa trichiloides* (1%), *Eriocolum macrcarpum* (1.33%) and *Cylindrodiscus gabonensis* (1.5%) have low importance value index indicating their low potential for regeneration hence require high conservation efforts. Most species had relative density of less than 1 about 18.33% and are considered a Rare and threatened or endangered species within the study area. These species may soon be absent in Ukpon Reserve if sustainable management practices are not adopted. According to Iroko *et al* (2008) over exploitation and replacement of natural forest ecosystems with human amenities results in the decimation of trees species.

The diversity index values of Shannon – Werner (3.04), Species Richness (20.67) and Species Evenness (2.19) was considered as high compare to those reported by Sundaranpandian and Swamy (2000) tropical forest of Kodayar in the western Ghats of Southern India and Edu (2017) in selected Forest Reserves of Cross River. These were however, similar to those obtained by Adekunle *et al* (2013) and Akindele *et al.* (2021) in Akure Forest Reserve; Aigbe and Omokhua, (2015) and Edet *et al.* (2012) in Oban and Afi mountain Reserves respectively in Cross River State. Species richness index measures the variety of species. It takes into

consideration the total number of a particular species in relation to the total number of individuals within the forest stand (Gebreselasse, 2011). The species richness obtained in this study was higher than the 10.444 reported for Afi Forest Reserve (Aigbe *et al.*, 2014) and 13.26 in Akure Forest Reserve (Akindele *et al.* (2021) all in Nigeria and 6.36-8.08 for Kibale Forest and 7.54-8.20 for kasyoha-kitomi Forest in the rift of Uganda (Eihe *et al.*, 2004). Species evenness is a measure of the relative abundance of species that make up the richness of an area. The higher the values of evenness the more even the species are in their distribution (Kent and Corker 1992). In this study the evenness index values of 2.19 was higher than the 0.887 recorded Aigbe and Omokhua, (2014) in Oban Forest Reserve, 0.907 obtained by Aigbe *et al.*, (2014) in Afi River Forest Reserve and 1.90 for Cross River South Forest Reserve by Edu, (2017). Akindele *et al.* (2021) obtained a lower value of 0.86 in Akure forest Reserve. Index of dominance shows, that two individuals drawn at random from a population belong to the same species. The higher the index of dominance values the lower the diversity of the species Aigbe *et al.*, 2014). The values of 0.06 obtained in this study is considered as low compared with the 0.78 reported by Akindele *et al.* (2021) in Akure Forest Reserve, indicating, that one particular tree species is dominating more than the others. This also implies that tree diversity is higher in the study area than in Akure Forest Reserve. This corroborates the findings of Egom and Umerod (1990) and Aigbe *et al.* (2014), who reported that species diversity varies across vegetation in tropical rainforest which may either be abundance or scanty in a particular location.

The influence of soil property on the distribution of flora has been documented by some scholars in the past. Aweto (1981) identified high organic matter and clay proportion as soil variables that exerted marked influence on the distribution and abundance of tree species. This study recorded high values of clay (60.16%), organic matter (4.32%) and pH (5.77) indicating high fertility. Ukpong (1994) identify nutrients and salinity as factors influencing species variation in mangrove swamps whereas John *et al.*, (2007) identify soil pH as the strongest soil factor that influenced the distribution of species in the tropical forest. According to Zare *et al.* (2011) soil texture, Salinity, effective depth, available Nitrogen, Potassium, Organic matter, lime and moisture are major factors influencing variation in the pattern of vegetation. Brady and Weil (2008) reported that Clay have specific surface area giving them a tremendous capacity to absorb water and nutrients. However Sandy soil has low specific surface area with little capacity to hold water and nutrients, making them infertile and to leaching (Eynaw *et al.*, 2004). Edu (2017) reported that a low pH of the some selected soils of Cross River Forest Reserve was responsible for the lower species

diversity, dominance index and high evenness in the study areas. This finding also agrees with the assertion of John *et al.* (2007) and Zare *et al.* (2011) that tree species distribution, dominance and evenness correlates with soil chemical properties and possibly topography. Long *et al.* (2018) found a positive significant correlation between tree species diversity and Potassium content in a tropical secondary forest. The results from this study compares favourably with those of Nadeau *et al.* (2015) and Onyekwelu *et al.* (2008) on the effects of soil properties on trees diversities, abundance and composition.

### **Conclusion**

The result in this study revealed a fundamental information of tree species diversity and soil physicochemical properties of Ukpon River Forest Reserve. The family Leguminaceae shows dominance and the species *P. africana* belonging to the family Fabaceae have the highest relative abundance and importance value index. The highest basal area was recorded in *Bombax bonnopoens* while *Triplochiton scleroxylon* had the lowest area. About 68.34% of the trees were Rare while 18.33% were Threatened or Endangered. The diversity indices showed that tree species composition was high with low Dominance index (0.06). Soil properties such as Clay, Organic matter, Phosphours, Calcium, Cation Exchangeable capacity and Base Saturation were high with mildly acidic pH. This may have been responsible for the high species diversity recorded in this study although some species are endangered. Therefore conservation measures should be recommended to checkmate the disappearance of these species by safeguarding the areas from anthropogenic activities to avoid degradation and deforestation.

### **References**

- Abbey, M. L (2006): Evaluation of flora resources, and utilization in Ikwe Game Reserved. Unpublished B. Forestry project in Department of Forestry University of Agriculture Makurdi, 10-29pp.
- Adekunle, V. A. J., Olagoke, A.O. and Ogundare, L.F. (2010). Rate of timber production in a tropical rainforest ecosystem of southwest Nigeria and its implications on sustainable forest management. *Journal of Forestry Research* **21**:225-230
- Adeyemi, A. A., Ibe A. E., and Okedimma, F. C, (2015), Tree structural and diversities in Okwangwo forest, Cross River State, Nigeria, Department of forestry and wildlife Technology, Federal University of Technology, Owerre, Nigeria. Ifewr publication, 3Pp., *Journal of Research in Forestry and Wildlife and environment* Vol 7.

- Adeyemi, A.A., Jimoh, S.O. and Adesoye, P.O. (2013). Assessment of Tree Diversities in Oban Division of the Cross River National Park (CRNP), Nigeria. *Journal of Agriculture, Forestry and the Social Sciences*. 11(1): 216-230. *Agriculture*. 4(1): 13-18.
- Aigbe H.I , Akindele S.O, Onyekwelu J, C. (2014). Tree Species Diversity And Density Pattern In Afi River Forest Reserve, Nigeria, *Journal of Scientific & Technology Research Volume 3*, Issue 10, October 2014 ISSN 2277-8616. 4Pp.
- Aigbe, H. I. and Omokhua, G. E. (2015).Tree Species Composition and Diversity in Oban Forest Reserve, Nigeria, *Journal of Agricultural Studies* ISSN 2166-0379 2015, Vol. 3, No. 1. 13Pp.
- Akindele T. F., Adekunle, V. A. J and Lawal, A. (2021). Tree species diversity, abundance, and Soil Physicochemical Status of PSP 29, Akure Forest Reserve, Ondo State, Nigeria. *Journal of Research in Forestry, Wildlife & Environment* Vol. 13(4): 120 - 129
- Akinsanmi, F.A.(1999). Updating Environmental Information in Nigeria. Invited Lead paper presented at the 26th Annual Conference of the Forestry Association of Nigeria held in Maiduguri, Borno. November 7
- Armenteras, D., Rodriguez, N. and Retana, J. (2009). Are conservation strategies effective in avoiding the deforestation of the Colombian Guyana Shield? *Biological Conservation* 42: 1411–1419
- Aweto, A.O. (1981). Secondary succession and soil fertility restoration in south-western Nigeria: Soil and vegetation interrelationships. *J Ecology*, 69: 957-963.
- Brady, N. C. and Weil R. R. (2008). The nature and properties of soils. Fourteen Edition. Pearson Education Inc. 1046Pp. Condit R *et al.*, (2002). Quantifying the deciduousness of tropical forest canopies under varying climate *Journal veg. Science* (5) 649-658.
- Chaturvedi, R.K, and Raghubanshi, A S. (2018). Effect of Soil Moisture on Composition and Diversity of Trees in Tropical dry Forest. *MOJ Eco Environ Sci* 3(1): 00059. DOI: 10.15406
- Clay, D.E, Carlson, C.G., clay, S.A. and murrell, T.S. (2012). Mathematics and calculations for agronomists and soil by the international plant nutrition institute and South Dakota state university, USA, page122.
- Edet, D. I., Ijeomah, H. M. and Augustine, U. O. (2012).Preliminary assessment of tree species diversity in Afi Mountain Wildlife Sanctuary, Southern Nigeria. Department of Forestry and Wildlife Management, University of Port Harcourt, P.M.B 5323, Choba, Port Harcourt, Nigeria, *Agriculture and Biology Journal of North America*. P.M.B 1115 Calabar, Nigeria. 2Pp.
- Edu, F. E. (2017). Effects of Natural Sites and Soil Types on Tropical Tree Species Distribution of Selected Forest Reserves in Cross River State, Obubra Campus. *MSc. Thesis (Unpublished)*.

- Egom I. P and Umerod S. J (1990). "The Conhng challenges of testing species distribution models" journal of Applied Ecology 42(4) 720 – 730Pp.
- FRA, (2-010): Global Forest Resources Assessment 2010, Main report. FAO Forestry Paper, 163, FAO Rome, 340 pp.
- Gebreselasse, G. V. (2011). Plant Communities Species Diversity seedling bank and re-sprouting in Nandi Forest, Kenya. PhD Thesis, Universitat Koblenz-Landau.
- Gillespie, T. W., Brock, J. and Wright, C.W. (2004). Prospects for quantifying structure, floristic composition and species richness of tropical forests. *Int. J. Remote Sensing* 25(4), 707–71Pp.
- Iheyen, J., Okoegwale, E.E., and Mensah, J.K. (2009). Composition of tree species in Ehor Forest Reserve, Edo State, Nigeria. *Nature and Science* 7(8): 8 – 18.
- Ikyaagba, T. E., Tee, T. N., Dagba, B.I., Ancha, U. P., Ngibo, K. D., and Tume, C. (2015). Tree Composition and Distribution in Federal University of Agriculture Makurdi, Nigeria. 3Pp. Inc., New Jersey, USA
- Iroko, O. A., Kareem, A. A., Adio, A. F. and Gbadebo, J. O. (2008). Impact of Human Activities on the Forest and their effects on Climate Change. In: Popoola, L (ed). Proceedings of the 32<sup>nd</sup> Annual Conference of the Forestry Association of Nigeria (FAN) held in Umuahia, Abia State between 20th and 24th October, 2008. Pp 208-214Pp.
- IUCN (2010). *Plants under pressure –a global assessment. The first report of the IUCN Sampled Red List Index for Plants*. Royal Botanic Gardens, Kew, UK, Natural History Museum, London, and International Union for Conservation of Nature.
- Jenkins, C. N., and Joppa, L. (2009). Expansion of the global terrestrial protected areas system. *Biological Conservation*, 142: 2166-2174.
- Jimoh S. O., Adesoye P. O., Adeyemi A.A., &Ikyaagba E.T., (2012). Forest Structure Analysis in the Oban Division of Cross River National Park, Nigeria. *Journal of Agricultural Science and Technology B* 2,510-518.
- John, R., Dalling, J. W., Harms, K. E., Yavitt, J.B., Stallard, R. F., Mirabello, M., Hubbell, S.P., Valencia, R., Navarrete, H., Vallejo, M., Foster, R. B. (2007). Soil Nutrients influence spatial distributions of tropical tree species. *PNAS*, 104 (3): 864–869.
- Kanowski, J., Catterall, C.P., Wardell-Johnson, G.W., Proctor, H. and Reis, T. (2003). Development of forest structure on cleared rainforest land in eastern Australia under different styles of reforestation. *Forest Ecology and Management* 183: 265-280.
- Kent, M. and Coker, P., (1992). *Vegetation description and analysis: a practical approach*. Belhaven press London, 363Pp.
- Long, W., Long, C., Yang, X., Li, D., Zhou, W. and Zhang, H. (2018). Soil Nutrients Influence Plant Community Assembly in Two Tropical Coastal Secondary Forests. *Tropical Conservation Science*, 11: 1-9.

- Maliyat, S. D. and Datt, B.(2010). A contribution to the flora of Katarnia Ghats Wildlife Sanctuary, Baharaich District, Uttar Pradesh. *Journal of Economy and Taxonomy Botany* **34**: 42-68.
- Mani, S. and Parthasarathy, N. (2006). Tree diversity and stand structure in inland and coastal tropical dry evergreen forests of peninsular India. *Current Science* 90: 1238–1246.
- Margalef, R. (1958). Temporal succession and spatial heterogeneity in phytoplankton. In: Buzzati-Traverso (ed.). *Perspectives in Marine biology*. University of California Press, Berkeley, 323 – 347pp.
- Martin B. Nadeau and Thomas P. Sullivan (2015). Relationships between Plant Biodiversity and Soil Fertility in a Mature Tropical Forest, Costa Rica. *Hindawi. Vol. 1 pp13*.
- Nadeau and Sullivan, (2015). Species diversity in ecological communities Univeristy of Chicago press. Chieago III U.S.A.
- Nirmal, J.I., Kumar, R.N., Kumar, R.K., Bhoi and Sajish P.R. (2010). “Tree species diversity and soil nutrient status in three sites of tropical dry deciduous forest of Western India,” *Tropical Ecology*, vol. 51, no. 2, Pp. 273-279.
- Ogwu, M.C., Osawaru, M.E. and Obayuwana, O.K. (2016). Diversity and Abundance of Tree Species in the University of Benin, Benin City, Nigeria. *Applied Tropical Agriculture*, 21(3): 46-54.
- Omor, L.M.A., Pellikka, P.K.E. and Rogers, P.C. (2010). Tree species diversity, richness, and similarity between exotic and indigenous forests in the cloud forests of Eastern Arc Mountains, Taita Hills, Kenya. *Journal of Forestry Research* 21(3): 255–264
- Onyekwelu, J.C., Mosandl, R., &Stimm, B.(2008).Tree species diversity and soil status of primary and degraded tropical rainforest ecosystems in South-Western Nigeria. *Journal of Tropical Forest Science*, 20 (3),193 -204.
- Pankaj Panwar and S. D. Bhardwaj, (2012). *Handbook of Practical Forestry*: Pp. 65 – 73.
- Peng, J., Cao, F., Liu, Z., Cao, J., Wu, L., Li, M., and Dong, X. (2015). A Correlation Analysis of Rocky Desertification Grades, Plant Diversity and Soil Factors in Central Hunan of China. *Acta Scientiae et Intellectus*, 1(2): 45-57
- Richard S.W, (2006). *An Introduction to Tropical Rainforest* second edition Oxford University press, Oxford 285Pp.
- Shannon, C. E. and Wiener, W. (1949). *The mathematical theory of communication*. Urbana, University of Illinois Press. 177 p.
- Sundarapandian, S. M. and Swamy, P. S. (2000). Forest ecosystem structure and composition along an altitudinal gradient in the Western Ghats, South India. *J. Trop. For. Sci.*, 12, 104-123.

- Suratman, M. N. (2012). Tree Species Diversity and Forest Stand Structure of Pahang National Park, Malaysia. In: Biodiversity Enrichment in a Diverse World. Chapter 18. INTECH. 473-492pp
- Udo, E. J, Ibia, O. T., Ogunwale, A.J., Ano, O. A. and Esu, E. I. (2009). Manual of soil, plant and water analysis. Sibon books Fourth Avenue Festac Lagos, 183pp
- Ukpong, I.E. (1994). Soil-vegetation interrelationships of mangrove swamps as revealed by multivariate analyses. *Geoderma*, 64: 167-181.
- USAID/Nigeria. (2008). Nigeria Biodiversity and Tropical forestry assessment: Maximizing Agricultural Revenue in Key Enterprises (Markets). Chemonics International Inc.
- Van Breemen, N. (1993) Soils as biotic constructs favouring net primary productivity. *Geoderma* 57, 183-211.
- World Rainforest Movement (WRM), (1999). Africa: Background document. Workshop on underlying causes of deforestation and forest degradation, 18-22 January, 1999, Costa Rica.
- Zare S, Jafari M, Tavili A, Abbasi H, Rostampour M 2011. Relationship between environmental factors and plant distribution in arid and semi-arid area (Case Study: Shahriyar Rangelands, Iran). *American-Eurasian J Agric and Environ Sci*, 10(1): 97-105
- Zhigila, D. A.; Abdul S. D. and Sawa, F. B. (2016). Plant Species Diversity, Abundance and Distribution in Communities of Zamfara State, Nigeria: Implications for Conservation. 6Pp.