

Biochemical and Thermodynamic Characterization of β amylase from *Dioscorea alata L.*

ABSTRACT

β -amylase (E.C.3.2.1.2) is a starch hydrolyzing enzyme fondly used in foods, pharmaceuticals, and brewing industries to convert starch into maltose. The aim of this study was to determine the physicochemical, kinetic, and thermodynamic properties as well as the potential industrial use of β -amylase from *Dioscorea alata L.* Studying the enzyme stability with Arrhenius methods showed that the enzyme was stable at a temperature range of 20–50 °C, and had good pH stability by retaining over 50 % of its initial activity over a wide range of pH from 4 – 8 and kinetic stability by increasing the half-life of the enzyme. The activation energy (E_a) for catalysis by water yam β -amylase at 25 °C was 6.45 kcal/mol. The activation energy (E_a), half-life, free energy change (ΔG^\ddagger), enthalpy change (ΔH^\ddagger), and entropy change (ΔS^\ddagger) for inactivation at optimum temperature (40 °C) and pH 5 were 13.92 kcal/mol, 41.25 min, 20.89 kcal/mol, 13.30 kcal/mol and -24.25 cal/mol/K respectively. K_m and V_{max} values were reduced from 2.25 to 2.13 mg/ml and 2.95 to 1.48 $\mu\text{mol}/\text{min}/\text{ml}$ respectively. The optimum pH shifted from 5 to 6, while the Optimum temperature increased from 40 to 50 °C after immobilization. Enzyme retained up to 67 % activity after 5 cycles. The enzyme would be of importance in manufacturing companies based on the kinetics and application features reported in this study, since it is a cheap and readily available source.

KEYWORDS: *Dioscoreaalata*, β -Amylase, Thermo-Stability, Half-Life, Immobilization, Industrial Application

1. INTRODUCTION

α -1,4-glucan bonds from the non-reducing ends of starch and other carbohydrate polymers are attacked by β -amylase, which converts them to maltose units. Maltose is a two-glucose disaccharide with numerous applications in the food and pharmaceutical industries [1-4,2,3,4].

Amylases are found in a wide range of microbes, plants, and mammals, and have found use in a variety of industries, including starch liquefaction. [5-7 5,6,7]. Although chemical methods can be used to liquefy starch, enzymatic hydrolysis can be done under mild circumstances, avoiding the harsh conditions

required by chemical treatments. Furthermore, unlike chemical procedures, the enzymatic approach did not result in water pollution [8]. The β -amylase reaction is important in the formation of maltose and the creation of fermented foods and alcoholic beverages [9]. β -amylase is used as a biocatalyst in a variety of systems, including packed bed reactors and fluidized-bed reactors.

In the tropics, yam is a staple food with nutritional value. It is one of the most widely cultivated and eaten plants. About 75-84 percent of the dry weight of yam is starch [10]. This might be used to determine the expression of numerous enzymes in the crop's subcellular regions, catalysing the conversion of starch into

its many derivatives. It could be a suitable source for industrial starch hydrolysing enzymes like β -amylase, which are in great demand. Although neglected compared to the popular type of *Dioscorea rotundata*, water yam (*Dioscorea alata*) is known for its bulkiness, high moisture, and starch content [11]. *Dioscorea alata* is one of the six economically important yam species, although it is underutilized in Nigeria for main food products due to traditional prejudice, which fails to identify the species' distinctive qualitative traits and agronomic adaptability.

The purpose of this research was to look at the kinetics of chemical reactions, isothermal conditions, and the behaviour of phase transition processes using classical kinetic models. The major benefit was to look at the mechanism of interaction between kinetic models and activation energies, both of which are responsible for the process response, in relation to temperature changes [12]. We can circumvent issues with free enzyme activity by immobilizing enzymes on solid supports. It has the ability to improve enzyme activity, selectivity, and stability. It also permits the biocatalyst to be reused as long as the enzyme is active in a continuous process, lowering process costs, particularly in industrial applications [13-17, 13,14,15, 16,17]. The immobilization procedure could improve enzyme activity and solubility in reaction fluids. [16-18, 16,17,18].

2. MATERIALS AND METHODS

2.1 Materials

Water yam was obtained from a local vendor Ketu, Lagos, and identified by a Taxonomist from the University of Lagos, Nigeria. 8-anilino-1-naphthalene sulfonic acid (ANS), Trizma base, sodium chloride and spectroscopic grade ethanol were purchased from Sigma-Aldrich Fine Chemicals, St. Louis, MO, USA. BSA standard and Bradford reagent kit were products of Bio-Rad, Palo Alto, CA, USA. All other chemicals were commercial products of reagent/analytical grade.

2.2 β -Amylase Assay Method

The DNSA method was used to determine β -amylase activity. The reaction mixture contained 100 μ l of 1 % soluble starch in 50 mM sodium acetate buffer (pH 5) and 100 μ l of the enzyme. After 3 min of incubation at 25 °C with 200 μ l of 3,5-DNSA color reagent, the reaction was stopped. The mixture was boiled for 5 min before being cooled in ice and diluted with 1000 μ l of distilled water. A UV-visible spectrophotometer was used to measure absorbance at 540 nm.

The enzyme activity was measured in micromoles of maltose released per minute per milliliter, which is the amount of reducing sugar released by the enzyme during the reaction [19].

2.3 β -amylase immobilization on sodium alginate beads

The beads were made according to Sachin and Sandeep 2012 with slight modification [20]. 4 g of sodium alginate was dissolved in 100 ml acetate buffer to make a 4 % solution. 10 ml of 3.4 mg of the enzyme was mixed with 10 ml of 4 % sodium alginate solution. The beads were formed by dripping the polymer solution from a height of approximately 20 cm into an excess of stirred 0.1 M CaCl_2 solution with a syringe and needle at room temperature in the calcium solution, left to cure for 30 min, filtered, washed thoroughly with distilled water several times, dried with filter paper, and exposed to an open air for 1 hour before use.

Immobilization yield %

$$= \frac{\text{activity of immobilized enzyme}}{A-B} \times 100$$

Where, A= activity of free enzyme added,
B= activity of soluble enzyme

2.4 The effect of substrate concentration on enzyme activity

To determine the kinetic constants K_m , V_{max} , and K_{cat} , 100 μ L of purified β -amylase was

incubated with 100 μL of 50 mM sodium acetate buffer pH 5 at various concentrations of soluble starch solution ranging from 0 to 10mg/ml for 3 min at 25 $^{\circ}\text{C}$. This was also done with amylose, amylopectin, and glycogen. The immobilized enzyme was treated in the same way, with 1 g incubated. A Lineweaver-Burk plot was used to calculate the values of K_m , V_{max} , and K_{cat} .

2.5 Effect of pH on enzyme activity and stability

Glycine-HCl pH 3, sodium acetate (pH 4-5), phosphate (pH 6-7), and Tris-HCl pH 8 were used to measure β -amylase activity at different pH levels. The β -amylase assay was used to carry out the reaction. The maximum activity was set to 100 %, and relative activity was plotted against various pH values.

The immobilized enzyme was treated in the same way. Purified β -amylase's stability was determined by pre-incubating the enzyme at 25 $^{\circ}\text{C}$ with 50 mM of the above-mentioned buffers, followed by a determination of residual activities at different time intervals. The initial value was set to 100 %, and relative activity was plotted against various time intervals.

2.6 Effect of Temperature on Enzyme Activity and Thermodynamic characterization

In a regulated water bath, a mixture of 100 μl of 1 % soluble starch and 100 μl of the appropriately diluted enzyme was incubated for 3 min at different temperatures ranging from 20 $^{\circ}\text{C}$ to 50 $^{\circ}\text{C}$. The assay was performed in accordance with standard procedures.

The effect of temperature on enzyme stability was studied by incubating the enzyme solution at different temperatures without substrate for 1 hour. An aliquot was taken every 10 min and tested for β -amylase residual activity. The $\ln k$ data was plotted

against the reciprocal of absolute temperature. Semi-logarithmic plots were used to calculate and analyze the first-order rate constants for denaturation K_d of the enzyme at various temperatures [21]. Thermodynamic parameters were calculated by rearranging the Eyring absolute rate equation:

$$\text{slope} = -E_a/R \quad 2$$

Where E_a is the Activation energy
 R (gas constant) = 1.987 cal/degree mol^{-1}

$$\Delta H^* = E_a - RT \quad 3$$

$$\Delta G^* = -RT \left(\frac{\ln K_d h}{K_B T} \right) \quad 4$$

$$\Delta S^* = \left(\Delta H^* - \Delta G^* / T \right) \quad 5$$

$$K_d = \left(\frac{K_B T}{h} \right) e^{(-\Delta H^* / RT)} e^{(\Delta S^* / R)} \quad 6$$

where h (Planck's constant) =
 1.584×10^{-34} cal/k,

K_B = Boltzman constant, $[R/N]$ =
 3.298×10^{-24} cal S,

N (Avogadro's no.) = 6.02×10^{23} mol^{-1} .

2.7 Thermal inactivation

Thermal inactivation of purified β -amylase was determined by incubating the enzyme for 60 min in 50 mM sodium acetate buffer pH 5 buffers at 40 $^{\circ}\text{C}$. Aliquots (100 μl) were taken at different times, cooled on ice, and the residual enzyme activity was measured.

The heat stability of the enzyme was investigated by incubating it at 40 $^{\circ}\text{C}$ for 60 min in the presence of 10 mM Fructose, Glucose, Galactose, and Maltose. Each sample's β -amylase level was measured [22]. The following equation was used to calculate the magnitudes of K_{inact} versus thermal condition:

$$\ln A/A_0 = -K_{i\text{anct}} t \quad 7$$

A and A₀ are the enzyme activity at each time and the initial time, respectively

2.8 Effect of Metals on the Enzyme activity and Stability

The enzyme was incubated in 50 mM acetate buffer pH 5 in the presence of each of the following salts: NaCl, CaCl₂, KCl, (NH₄)₂SO₄, ZnSO₄, MgSO₄, FeSO₄, CuSO₄ and HgCl. The enzyme activity was measured according to the standard assay.

The effect of metallic chlorides on stability was determined by incubating the enzyme in the presence of various metals for one hour at 40 °C. Residual activity was measured at 10 min interval. The blank contained the buffer solution with the metal ion under consideration without the enzyme. The control consisted of the enzyme and the substrate without any chloride.

Effect of metallic chlorides on the half-life of the enzyme was determined at a final concentration of 100 mM. The half-life was determined from the relationship:

$$t_{1/2} = \frac{\ln 2}{K_d} \quad 8$$

Where k_d is the deactivation rate constant that can be derived from:

$$V = V_0 e^{-K_d t} \quad 9$$

Where V₀ is the initial enzyme activity while V is the enzyme activity at time t of incubation; and -k_d can be obtained from the plot of:

$$\ln V = \ln V_0 - K_d t \quad 10$$

The D-value (decimal reduction time) is the time (min) to reduce 90 % of the enzyme activity and is given by the following equation

$$D = \frac{\ln 10}{K_d} \quad 11$$

2.9 Application of β-amylase

Wash efficiency of starch stains was studied in the presence of Mama gold and Rana detergents [23]. White cotton cloth pieces stained with a 1 % starch solution were baked at 80 °C for 30 min to ensure a firm binding of stains to the material support.

The cleaning performance was evaluated by varying the cleaner, which included water, water + detergent, water + enzyme, and water + detergent (7 mg/mL) + enzyme.

A stained cotton cloth piece was shaken for 30 min at 50 °C in the presence of the corresponding cleaner mixture. Each mixture obtained solution was collected in order to determine the concentration of reducing sugars released from starch.

The washing process's efficiency in removing starch was expressed as the following equation [24]:

$$\text{Efficiency \%} = \frac{100 * A * 0.95}{B} \quad 12$$

where A is the amount of maltose released (g/mL) during the wash procedure and B is the amount of starch (g/mL) used for staining the cotton cloth piece.

2.10 Reusability of Immobilized β-amylase

Enzyme reusability was checked by carrying out enzyme assay for a couple of times. At the end of each cycle, the derivative was washed with distilled water and activity buffer, a new substrate solution was added to start a new round of reaction. The initial and remaining activities were assayed at pH 5, 25 °C (initial activity at these conditions is regarded as 100 % [25])

2.11 Statistical analysis

All experiments were carried out in triplicates, unless otherwise stated. The results were analyzed using Graph pad prism

(version 5.0) and Kaleida Graph 4.5.4 synergy software. They were expressed as mean \pm standard deviation. $P < 0.05$ was considered as level of significance.

3 RESULTS AND DISCUSSION

β -amylase hydrolyzes starch, amylose, and glycogen but has no effect on maltose, sucrose, fructose, lactose, or galactose (Table 1). The optimal pH after immobilization increased from 5 to 6 (Fig. 1), while the optimal temperature increased from 40 °C to 50 °C (Fig. 2). Figure 3 depicts the Lineweaver-Burk plot for soluble and immobilized enzyme with soluble starch. The K_m and V_{max} values for free enzyme were 2.25 mg/ml and 2.95 $\mu\text{mol}/\text{min}/\text{ml}$, respectively, while the values for immobilized enzyme were 2.13 mg/ml and 1.48 $\mu\text{mol}/\text{min}/\text{ml}$. For soluble starch, amylose, amylopectin, and glycogen, purified β -amylase from water yam exhibited typical Michaelis-Menten kinetics. The purified enzyme with soluble starch had a K_{cat} of 317.5 s^{-1} (Table 2), which is higher than 137.93 min^{-1} obtained from fenugreek β -amylase by Srivastava *et al.*, 2014 [26]. Figure 4 depicts the Lineweaver-Burk plot with and without α -cyclodextrin as an inhibitor. The K_i and $K_{i\text{con}}$ concentrations were 2.70 mg/ml and 3.33 mg/ml, respectively. Figure 5 depicts the Lineweaver-Burk plot of β -amylase from water yam in the presence and absence of maltose as an inhibitor. V_i and V_{ii} values were 2.089 and 2.42 $\mu\text{mol}/\text{min}/\text{ml}$, respectively.

Maltose had a non-competitive inhibition pattern on the enzyme, whereas α -cyclodextrin had a competitive pattern, which is consistent with Srivastava *et al.*, 2014 [26].

The enzyme was most stable at alkaline pH levels of 7 and 8, with approximately 80 % residual activity, while at acidic pH levels of 4, 5, and 6, remaining activities of 65, 70, and 73 percent, respectively, were obtained after incubating for 4 hours (Fig 6). Acidic stable β -amylase from a luminal particle associated microorganism was reported [27], whereas Kolawole *et al.*, (2011) [22], Chiba and

Kuwashima, (2006)[28], and Kolawole *et al.*, (2006)[29] reported alkaline stable β -amylase from Chinese yam tuber, *Digitariaexilis*, and *Eleusine coracana*. Van Damme *et al.*, (2001) [30] proposed that, in addition to the catalytic site, other regulatory parts of the enzyme may be responsible for the stability of β -amylase at alkaline pH.




Fig 1: Effect of pH on both the free and immobilized enzyme




Fig.2: Effect of Temperature on free and immobilized enzyme

After 60 min of incubation, water yam β -amylase retained over 70 % of its initial activity at 50 °C (fig. 7). This could indicate its potential use in the food and beverage industries to convert starch to maltose, where thermostability and thermo-activity of the enzyme are important due to high temperature operating conditions [31].

The K_{inact} value of the inactivation rate constant is the best measure for describing the effect of osmolytes on the enzyme's kinetic stability. The thermo-inactivation data in Table 3 revealed that glucose increased the kinetic stability of β -amylase. It protected the enzyme against heat inactivation by stabilizing it. Sugars are known to protect proteins from losing their function [32,33], chemical [34], and thermal denaturation [35].

Water viscosity and surface tension are also affected by osmolytes [36]. Protein dynamics and structure, particularly protein flexibility, can be affected by environmental parameters such as pH, viscosity, ionic strength, temperature, prosthetic groups, solvent composition, and so on. As a result, the specific nature of interactions that determine osmolyte-mediated protein stability is yet unknown.

Overall, protein stability is determined by a delicate balance of favorable and unfavorable interactions between native and denatured protein states and co-solvent molecules [37].

The stabilizing impact is thus dependent on the type of the protein as well as the co-solvent molecules, and it may not be possible to generalize the effect. Table 4 shows that the activation energy of 6.48 kcal/mol was similar to that of pea epicotyls (6.28 kcal/mol), peanuts (4.5 kcal/mol) [38], and fenugreek [39]. It is related to the reaction rate. At high temperatures, the thermal unfolding of proteins is induced by a large rise in entropy change, which lowers the Gibbs free energy change of the unfolding transition [40]. Ca^{2+} , Mg^{2+} , and Zn^{2+} were added to boost the activity of β -amylase. This is consistent with the findings of Kolawole *et al.* (2011) [22], who found that Ca^{2+} boosted the activity of β -amylase from *E. coracana*, while Cu^{2+} and EDTA decreased activity. Metal binding is linked to variations in metal co-ordination

geometries and influences metal selection [41] (Table 5). The findings show that the thermostability of purified water yam β -amylase is pH dependent, implying that the high stability of water yam β -amylase at alkaline pH at high temperatures and the use of specific metals could imply a high industrial utility potential for the production of various high conversion maltose syrups from raw and liquefied starch (Table 6). Metal specificity appears to be present in β -amylases from several sources for increased thermo-activity and thermostability. The idea of the metal enhancing the chemical potential of the protein and so favoring the folded form over the unfolded state was suggested by Ray *et al.*, (1994) [42].

The main factors that determine an enzyme's stability are its half-life, decimal reduction time (D), and first-order rate constants (K_d) (Table 7). The results show that as the temperature increased, the $t_{1/2}$ and D-value decreased as the K_d increased, indicating that the enzyme is less stable at higher temperatures [43,44].

The addition of Zn^{2+} and Mg^{2+} resulted in a slight increase in both the enzyme half-life and the ΔG^\ddagger of thermal denaturation, as well as a decrease in ΔS^\ddagger . Other metals decreased both the half-life and the ΔG , but increased the ΔS^\ddagger of thermal denaturation (Table 8). The presence of metal ions had no effect on the ΔH^\ddagger of denaturation. The non-covalent bonds were unaffected by thermal denaturation in the presence of metal ions bonding but the opening of the structure by the change in ΔS^\ddagger . It shows that ΔS^\ddagger contributes to the ΔG^\ddagger denaturation but not the ΔH^\ddagger in the presence of all the metals. Thermal unfolding of proteins at high temperatures is caused by a strong increase of the entropy change that lowers Gibbs free energy change of the unfolding transition [41].

Table 1: Effects of various substrate on β -amylase

SUBSTRATE	RELATIVE ACTIVITY (%)
Soluble starch	100
Amylose	91
Amylopectin	87
Glycogen	40
Maltose	0
Sucrose	0
Fructose	0
Lactose	0
Galactose	0

Table 2: Kinetic parameters of β -amylase with various substrates

Substrates	Starch	Amylose	Amylopectin	Glycogen
K_m (mg/ml)	2.24	2.56	2.53	4.27
V_{max} ($\mu\text{mol}/\text{min}/\text{ml}$)	2.94	2.60	3.18	3.74
V_{max}/K_m	1.31	1.02	1.26	0.88
K_{cat} (S^{-1})	317.5	279.6	341.9	402.2
K_{cat}/K_m	141.1	109.2	135.2	94.2




Fig. 3: Effect of concentration on soluble and immobilized enzyme




Fig. 4 Lineweaver-Burk plot of β -amylase from water yam in the presence and absence of α -cyclodextrin as inhibitor. The activity was assayed in 50 mM acetate buffer pH 5 at 25 °C using DNSA method. The substrate concentration range was 0–10mg/ml. $p < 0.05$




Fig. 5: Lineweaver-Burk plot of β -amylase from water yam in the presence and absence of maltose as an inhibitor. The activity was assayed in 50 mM acetate buffer pH 5 at 25 °C using DNSA method. The substrate concentration range was 0–10 mg.

Table 3: The effect of Osmolytes on the kinetic stabilities of the enzyme

Ea of activation is 6.48 kcal/mol




Fig.6: Effect of pH on stability of the β -amylase. The enzyme was prepared by mixing the purified enzyme with two volumes of the buffer solution of pH 4 -8. Incubated at room temperature and the residual activity assayed

Osmolytes (10 mM)	K _{inact} (S ⁻¹)	ΔG 25 °C (Kcal/mol)
Control	6.17 X 10 ⁻⁵	20.76
Glucose	3.80 X 10 ⁻⁵	21.04
Maltose	5.39 X 10 ⁻⁵	20.84
Fructose	5.10 X 10 ⁻⁵	20.87
Galactose	7.10 X 10 ⁻⁵	19.37

hourly.

Table 5: Effects of metal ions on β -amylase activity

Temperature (K)	ΔG^\ddagger kcal/mol	ΔH^\ddagger kcal/mol	ΔS^\ddagger cal/mol/k
293	12.86	5.87	-23.46
298	12.84	5.86	-23.42
303	12.82	5.85	-23.39
308	12.80	5.84	-23.35
313	12.78	5.83	-23.32




Fig 7: Effect of temperature on enzyme stability. The residual activities were expressed relative to the maximum activity at 0 min which was taken as 100 %.

Table 4: Thermodynamic parameters for catalysis of water yam β -amylase

SUBSTRATE	RELATIVE ACTIVITY (%)
None	100
Ca ²⁺	110
Mg ²⁺	122
Zn ²⁺	120
Fe ²⁺	31
Pb ²⁺	22
Na ⁺	100
K ⁺	108
Hg ⁺	14
Iodoacetate	72
Urea	91
Fe ³⁺	89
Cu ²⁺	51
EDTA	81
Sn ²⁺	28
NH ₄ ⁺	78
Hg ⁺	14




Fig. 8: Reusability of β -amylase




Fig. 9: Efficiency of starch stain removal by β -amylase.

Enzyme retained up to 67 % activity after 5 cycles in reusability test after immobilization (Fig. 8), while β -amylase presence made starch removal more efficient as shown in Figure 9.

Table 6: Thermodynamic parameters for thermal inactivation of yam β -amylase at various pH at 40 °C

pH	Ea (Kcal/mol)	Half-life (min)	ΔG^\ddagger kcal/mol	ΔH^\ddagger kcal/mol	ΔS^\ddagger cal/mol/k
5	13.92	41.25	20.89	13.33	-24.25
6	14.41	41.01	20.89	13.79	-22.68
7	16.00	41.46	20.90	15.38	-17.64
8	16.24	42.00	20.90	15.62	-16.87

Table 7: Thermodynamic parameters for thermal inactivation of water yam β -amylase at various temperature

Temperature (K)	Half-life (min)	ΔG^\ddagger kcal/mol	ΔH^\ddagger kcal/mol	ΔS^\ddagger cal/mol/k
293	41.25	20.89	13.33	-24.25
298	31.50	21.07	13.29	-24.47
303	22.35	21.18	13.28	-24.46
308	16.50	21.32	13.27	-24.45
313	10.34	21.35	13.26	-24.29

Ea is 13.92 kcal/mol

Table 8: Thermodynamic parameters for metals on thermal inactivation of water yam β -amylase at 40 °C and pH 5

Metallic Chloride (10mM)	Half-life (min)	ΔG^\ddagger kcal/mol	ΔH^\ddagger kcal/mol	ΔS^\ddagger cal/mol/k
Control	41.25	20.89	13.33	-24.25
Na ⁺	41.25	20.89	13.33	-24.25
Ca ²⁺	40.29	20.88	13.33	-24.19
Zn ²⁺	42.78	20.92	13.33	-24.35
Mg ²⁺	43.31	20.93	13.33	-24.38
NH ₄ ⁺	40.54	20.88	13.33	-24.22

Ea is 13.92 kcal/mol

CONCLUSION

The fact that the β -amylase was active over a wide pH range may imply that the enzyme will be useful in processes that are subjected to wide pH change from the acidic to alkaline range. The enzyme can be employed in food and beverage industries to convert starch into maltose where high value has been placed on the thermostability and thermo-activity of the enzyme.

The high stability of water yam β -amylase at alkaline pH, high temperatures and the use of some specific metals could imply high industrial utility potential for the production of various high conversion maltose syrups from raw and liquefied starch. The low free energy of activation offers low cost in terms of energy requirements.

CONSENT AND ETHICS APPROVAL

Not applicable (the study does not involve human participant and tissues).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

[1] Reddy NS, Nimmangadda A, Sambasiva RKRS. An overview of the microbial α -amylase family. *African J. Biotechnol.* 2003; 2:645–8.

[2] Schramm M, Loyter A. Purification of alpha-amylase by precipitation of amylase-glycogen complex. *Method Enzymol.* 1996; 9:533–7.

[3] Guzmán-Maldonado H, Paredes-López O. Amylolytic enzymes and products derived from starch: A review. *Crit. Rev. Food Nutr.* 1995; 35:373–403.

[4] Coutinho PM, Reilly PJ. Glucoamylase structural, functional and evolutionary relationships. *Proteins.* 1997; 29:334–47.

[5] Pandey A, Nigam P, Soccol CR, Soccol VT, Singh D, Mohan R. Advances in microbial amylases. *Biotechnol Appl. Biochem.* 2000; 31:1052-1058

[6] Van der Maarel MJEC, Van der Veen B, Uitdehaag JCM, Leemhuis H, Dijkhuizen L. Properties and applications of starch converting enzymes of the alpha-amylase family. *J. Biotechnol.* 2002; 94:137–55.

[7] Atichokudomchai N, Jane J, Hazlewood G. Reaction pattern of novel thermostable alpha-amylase. *Carbohydr Pol.* 2006; 64:582–8.

[8] Husain S, Jafri F, Saleemuddin M. *Enzyme Microb. Technol.* 1996; 18:275–80.

[9] Lin KH, Fu H, Chan CH, Lo HFL, Shih MC. Generation and analyses of the transgenic potatoes expressing heterologous thermostable beta amylase, *Plant Sci.* 2008; 174: 649–656.

[10] Moura, G.S., Lanna, E.A.T., Donzele, J.L., Falkoski, D.L., Rezende, S.T., Oliveira, M.G.A. ~~and~~ albino. L.F.T (2016). Stability of enzyme complex solid-state fermentation Subjected to the processing of pelleted diet and storage time at different temperatures. *R. Bras. Zootec.* 45(12):731-736.

[11] Pervez S, Afsheen A, Samina I, Nadir NS. ~~and~~ Shah, A.U Saccharification and liquefaction of cassava starch: an alternative source for the production of bioethanol using amylolytic enzymes by the double fermentation process. *B.M.C. Biotech.* 2014; 14:49–59.

[12] Gutzow I, S. Todorova, N. Jordanov. Kinetics of chemical reactions and phase transitions at changing temperature: general reconsiderations and a new approach, *Bulg. Chem. Comm.* 2010; 42:79–102.

[13] Dos Santos JCS, Bonazza HL, De Matos LJBL, Carneiro EA, Barbosa O, Fernandez-Lafuente R, Gonçalves LRB, De Sant'Ana HB, Santiago-Aguiar RS. Immobilization of CALB on activated chitosan: application to enzymatic synthesis in supercritical and near-critical carbon dioxide, *Biotechnol. Rep.* 2017; 14:16–26.

[14] Rios NS, Pinheiroa MP, Dos Santos JCS, De Fonseca TS, Limac LD, De Mattos MC, Freire DMG, Da Silva Júnior JJ, Rodríguez-Aguadoe E, Goncalves LRB. Strategies of covalent immobilization of a recombinant *Candida antarctica* lipase B on pore-expanded SBA-15 and its application in the kinetic resolution of (R, S)-Phenyl-ethyl acetate, *Mol. Catal. B: Enzym.* 2016; 133:246–258.

[15] Tavanoa OL, Fernandez-Lafuente R, Goulart AJ, Monti R. Optimization of the immobilization of sweet potato amylase using glutaraldehyde-agarose support. Characterization of the immobilized enzyme, *Proc. Biochem.* 2013; 48:1054–1058.

[16] Sheldon R.A, S. van Pelt. Enzyme immobilization in biocatalysis: why, what and how, *Chem. Soc. Rev.* 2013;42: 6223–6235.

[17] DiCosimo R, McAuliffe J, Poulouse AJ, Bohlmann G. Industrial use of immobilized enzymes, *Chem. Soc. Rev.* 2013; 42: 6437–6474.

[18] Rodrigues RC, Hernandez K, Barbosa O, Rueda N, Garcia-Galan C, Dos Santos JCS, Berenguer-Murcia A, Fernandez-Lafuente R. Immobilization of proteins in poly-styrene-divinylbenzene matrices: functional properties and applications, *Curr. Org. Chem.* 2015; 19:1707–1718.

[19] Zhang G, Brox S, and Weiner JH. Extracellular accumulation of recombinant proteins fused to the carrier protein YebF in *Escherichia coli*. *Nature Biotechnology.* 2006; 24(1): 100-104.

[20] Sachin Talekar and Sandeep Chavare. Optimization of immobilization of α -amylase in alginate gel and its comparative biochemical studies with free β -amylase. *Recent Research in Science and Technology*. Recent Res. Sci. Technol. 2012; 4(2): 01-05.

[21] Lonhienne, T, Gerday, C, Feller, G. Psychrophilic enzymes: revisiting the thermodynamic parameters of activation may explain local flexibility *Biochimica et Biophysica Acta-BBA.* 2000; 1543 :1-10.

[22] Kolawole AO, Ajele, JO, Ravi Sirdeshmukh. Purification and Characterization of alkaline-stable β -amylase in malted African finger millet (*Eleusine coracana*) seed. *Plant Biochemistry* 2011; 46:2178-2186

[23] Qureshi AS, Khushk I, Ali CH, Chisti Y, Ahmad A, and Majeed H. “Coproducts of protease and amylase by thermophilic *Bacillus* sp. BBXS-2 using open solid-state fermentation of lignocellulosic biomass,” *Biocatalysis and Agricultural Biotechnology.* 2016; 8: ~~vol. 8, pp.~~ 146–151.

[24] Shukla RJ and Singh SP. “Characteristics and thermodynamics of α -amylase from thermophilic actinobacterium,

Laceyellacchari TSI-2," Process Biochemistry. 2015; 50 (12): ~~vol. 50, no. 12, pp.~~2128–2136.

[25] Tavanoa O.L, R. Fernandez-Lafuente, A.J. Goulart, R. Monti. Optimization of the immobilization of sweet potato amylase using glutaraldehyde-agarose support. Characterization of the immobilized enzyme, Proc. Biochem. 2013; 48:1054–1058.

[26] Srivastava G, Vinay K. Singh, Arvind M. Kayastha. Identification of active site residues of Fenugreek β -amylase: Chemical modification and in silico approach. ~~Plant Physiol. Biochem. Plant physiology and Biochemistry.~~ 2014; 83:217-224.

[27] Martinez TF, Diaz M, Moyano FF. Inhibition of amylases present in luminal particle associated microorganism. ~~J. Sci. Food Agric. Journal of Science, Food and Agriculture.~~ 2002; 82:398–404.

[28] Chiba Y, Kuwashima T. Purification and characterization of alkali stable beta amylase from Chinese Yam (Nagaimo) Tuber. ~~J. of Appl. Glycosci. Journal of Applied Glycoscience.~~ 2006; 53:273–5.

[29] Kolawole AO, Sanni, DM Ajele, JO. Catalytic and thermodynamic properties of *Digitaria exilis* β -amylase. ~~J. of Appl. Glycosci.~~ 2006; 1(4):273-285

[30] Van Damme V.J.M, Hu J, Barre A, Hause B, Baggerman G, Rouge P, *et al.*, Purification, characterization, immunolocalization and structural analysis of the abundant cytoplasmic β -amylase from *Calystegia sepium* (hedge bindweed) rhizomes. ~~Eur. J. Biochem. European Journal Biochemistry.~~ 2001; 268:6263–73.

[31] Swamy MV, Sai Ram M, Seenayya G. β -Amylase from *Clostridium thermocellum* SS 8-a thermophilic, anaerobic, cellulolytic bacterium. ~~Lett. Appl. Microbiol. Letters in Applied Microbiology.~~ 1994;18:301–4.

[32] Carcini, P, Nishiyama Y, Westover A, Itoh M, Nagaoka S, Sasaki N, Okazaki Y, Muramatsu M, and Hayashizaki Y. Thermo-stabilization and thermo-activation of thermolabile enzymes by trehalose and its application for the synthesis of full-length cDNA; ~~PNAS Proceedings of the National Academy of Sciences, USA,~~ 1998; 95:520–524.

[33] Taylor L S, York P, Williams A C, Edwards H G M, Mehta V, Jackson G S, Badcoe I G and Clarke A R. Sucrose reduces the efficiency of protein denaturation by a chaotropic agent. ~~BBA. Biochemical Biophysical Acta~~ 1995;1253: 39–46.

[34] Sola-Pena M, Ferreira-Pereira A, Lemos A P and Meyer-Fernandes J R. Carbohydrate protection of enzyme structure and function against guanidinium chloride treatment depends on the nature of carbohydrate and enzyme. ~~Eur. J. Biochem. European Journal of Biochemistry.~~ 1997; 248:24–29.

[35] Xie G and Timasheff S N. The thermodynamic mechanism of protein stabilization by trehalose; ~~Biophys. Chem. Biophysical Chemistry of Sciences.~~ USA, 1997; 95:520-524.

[36] Schein CH. Solubility as a function of protein structure and solvent components; Biotech. 1990; 8: 308-317.

[37] Timashef S N. Control of protein stability and reactions by weakly interacting cosolvents: the simplicity of the complicated

~~Adv. Protein Chem. Advance Protein Chemistry.~~ 1998; 51 355-43.

[38] Das R, Talat M, Srivastava ON, Kayastha AM. Covalent immobilization of peanut β -amylase for producing industrial nano-biocatalysts: a comparative study of kinetics, stability and reusability of the immobilized enzyme, Food Chem. 2018; 245: 488–499.

[39] Agrawal DC, Yadav A, Kesarwani R, Srivastava ON, Kayastha AM. Immobilization of fenugreek β -amylase onto functionalized graphene quantum dots (GQDs) using Box-Behnken design: Its biochemical, thermodynamic and kinetic studies. ~~Int. J. Biol. Macromol. International Journal of Biological Macromolecules.~~ 2020; 144:170-182.

[40] Pal D, Chakrabarti P. Estimates of the loss of main-chain conformational entropy of different residues on protein folding. Proteins. 1999; 36:332–339.

[41] Vieille C, Epting KL, Kelly RM, Zeikus JG. Bivalent cations and amino-acid composition contribute to the thermostability of *Bacillus licheniformis* xylose isomerase. ~~Eur. J. Biochem. European Journal of Biochemistry.~~ 2001; 268:6291 –301.

[42] Ray R, Jana SC, and Nanda G. Biochemical approaches of increasing thermostability of β -amylase from *Bacillus megaterium* B₆. FEBS Letters. 1994; 356: 30- 33.

[43] Pal A, F. Khanum. Covalent immobilization of xylanase on glutaraldehyde activated alginate beads using response surface methodology: characterization of immobilized enzyme, Proc. Biochem. 2011; 4: 1315-1320

[44] Marin E, Sanchez L, Perez M, Puyol P, Calvo M. Effect of heat treatment on bovine lactoperoxidase activity in skim milk, kinetic and thermodynamic analysis, J. Food Sci. 2003; 68: 89–93.