

Case study

Title: Middle ear adenoma with uncommon presentation: A case report

Abstract

Adenomas of the middle ear are even rarer. Although considered benign tumors, they carry a risk of recurrence and malignant transformation. We present a 37-year-old patient with a hypoacusis of the left ear with purulent otorrhea on a non-marginal tympanic perforation as symptomatology. The audiogram showed transmission deafness, and the scanner found a fleshy attic mass with a preserved ossicular chain. The patient underwent an antro-atticotomy with mass excision, the anatomopathological result of which was in favor of an adenoma of the middle ear. The patient has shown no recurrence for almost 2 years. In conclusion, middle ear adenomas are rare neoplasms and have no specific symptoms or clinical presentation. Total exploration and extirpation is required for treatment, and microscopic and immunohistochemical examinations are mandatory for a definitive diagnosis.

Keywords: case report, adenoma, middle ear, atticotomy

Introduction

Ear tumors form a heterogeneous group whose diagnosis must be evoked in the face of any unilateral damage (hearing, vestibular), any duct polyp, or unfavorable development of a usual treatment must lead to an auditory examination [1]. The adenoma of the middle ear is a benign tumor, but there is a risk of recurrence and malignant transformation [2].

Middle ear adenoma is a rare disease with few published cases in the literature [1-4]. Here we presented a 37-year-old patient with a hypoacusis of the left ear with purulent otorrhea who underwent surgery and was diagnosed histopathologically as an adenoma of the middle ear.

Case Presentation:

A 37-year-old patient consulting for recurrent left ear infections associated with hearing loss. otoscopy find the presence of purulent otorrhea on a non-marginal tympanic perforation and a fleshy polyp in the middle ear. Acoumetry and audiogram find transmission deafness.

A computed tomography (CT) scan of the rocks was performed, which resulted in the presence at the attic level of a fleshy filling with a continuous ossicular chain (Figure 1).

The patient underwent antro-atticotomy with mass removal, the anatomopathological result of which was in favor of an adenoma of the middle ear (Figure 2). The patient has shown no recurrence for almost 2 years.

Discussion

Middle ear adenoma (sometimes incorrectly called "carcinoid tumor" or "ceriminoma" is a rare benign epithelial tumor; only about 100 cases are reported, with a small (10mm) well- limited size [5, 6] . It is derived from pluripotent epithelial cells of the middle ear mucosa, of endodermal origin, which have a double capacity of exocrine differentiation (cells with mucosal secretion granules) and neuroen-docrine differentiation (cells with neurosecretory granules) [6] .

The average age of onset is between 20 and 40 years. The most common symptoms are hearing loss, fullness of the ear, tinnitus and otorrhea. Dizziness and peripheral facial paralysis are more rarely reported [1, 7] .

Clinical examination may reveal non-pulsatile retrotympenic greyish mass. The eardrum is rarely invaded by the tumors [7] .

The main differential diagnoses are usually eliminated by preoperative radiologic imaging [CT and Magnetic resonance imaging (MRI)]: paraganglioma, cholesteatoma, or much more rarely schwannoma of the tympanic segment of the facial nerve, meningioma, etc. [3] . In CT

scan, adenoma occurs as a mass of the eardrum cage that may affect the patient. groin the ossicular chain. Mastoid extension, ossicular lysis and bone erosion – including the tegmen rupture as described in our clinical case – are rare. In MRI, the lesion is elevated after gadolinium injection and is not hypersignal on the diffusion sequences, distinguishing it from the cholesteatoma [6] .

Given the rarity of these lesions and the lack of specific clinical presentation, the diagnosis is rarely mentioned in the preoperative period. The treatment is surgical removal, sometimes with the need for an ossicular sacrifice to allow complete removal [6] . The diagnosis is then confirmed by histological and immunohistochemical examination. Recurrences after complete removal are rare [8].

Conclusion:

Middle ear adenomas are rare neoplasms and have no specific symptoms or clinical presentation. Total exploration and extirpation is required for treatment, and microscopic and immunohistochemical examinations are mandatory for a definitive diagnosis.



Figure 1: CT images of a middle ear adenoma, showing a non specific middle ear mass (arrow).



Figure 2: intraoperative photo showing the adenoma after mastoidectomy (arrow).




Figure 3: photo showing the histological examination of adenoma

References:

1. Isenring D, Pezier TF, Vrugt B, Huber AM. Middle ear adenoma: case report and discussion. *Case Rep Otolaryngol.* 2014;2014:342125. PubMed PMID: 25045567. Pubmed Central PMCID: PMC4090429. Epub 2014/07/22. eng
2. Sterrer E, Windisch F, Frey K, Rettensteiner K, Loader B. Middle ear adenoma with neuroendocrine differentiation : A pediatric case report. *Wien Klin Wochenschr.* 2017 Mar;129(5-6):208-11. PubMed PMID: 27761737. Epub 2016/10/21. eng

3. Cardoso FA, Monteiro EMR, Lopes LB, Avila M, Scarioli BO. Adenomatous Tumors of the Middle Ear: A Literature Review. *Int Arch Otorhinolaryngol*. 2017 Jul;21(3):308-12. PubMed PMID: 28680503. Pubmed Central PMCID: PMC5495583. Epub 2017/07/07. eng
4. Wang L, Ouyang N, Guo HX, Liu QY. Middle ear adenoma with uncommon presentation and literature review. *BMJ Case Rep*. 2017 Jun 13;2017. PubMed PMID: 28611057. Pubmed Central PMCID: PMC5534660. Epub 2017/06/15. eng
5. da Luz Martins M, Boutin P, Bozorg-Grayeli A, Cazals-Hatem D, Bouccara D, Sterkers O. [Middle ear adenoma: report of two cases and review of the literature]. *Ann Otolaryngol Chir Cervicofac*. 2002 Dec;119(6):337-40. PubMed PMID: 12527842. Epub 2003/01/16. Adénome bénin de l'oreille moyenne: a propos de deux cas. fre
6. Torske KR, Thompson LD. Adenoma versus carcinoid tumor of the middle ear: a study of 48 cases and review of the literature. *Mod Pathol*. 2002 May;15(5):543-55. PubMed PMID: 12011260. Epub 2002/05/16. eng
7. Mori E, Kojima H, Wada K, Moriyama H. Middle ear adenoma diagnosed by recurrent facial paralysis. *Auris Nasus Larynx*. 2009 Feb;36(1):75-8. PubMed PMID: 18328653. Epub 2008/03/11. eng
8. Bakhos D, Lescanne E, Fetissof F, Robier A, Morinière S. Neuro-endocrine adenoma of the middle ear: a case study. *Eur Arch Otorhinolaryngol*. 2007 Dec;264(12):1525-8. PubMed PMID: 17639440. Epub 2007/07/20. eng