

Original Research Article
**Growth, Yield and Grain Arsenic Concentration
of Rice Cultivars Under Varying Levels of Soil
Arsenic Contamination**

ABSTRACT

Fourteen cultivars of rice (Summer rice) were studied to evaluate the effects of varying levels of arsenic (As) contamination on growth, yield and grain As contents. Arsenic contamination adversely affected tillering, filled grains, grain yield and straw yield of all rice cultivars, however, the effect varied among the cultivars. In As control treatment, the highest grain yield (75.66 g pot⁻¹) was observed in BR 11 while the lowest grain yield (8.73 g pot⁻¹) was in BRR1 dhan4. Application of 20 ppm As to the soil resulted in complete death of BR 11, BRR1 dhan30, BRR1 dhan33, BRR1 dhan34, BRR1 dhan41, Binadha-4, Biroy and Kalizeera cultivars. Arsenic contamination significantly increased As concentration in grain for all cultivars of rice. Arsenic concentration on rice grains grown in soils without As contamination ranged from 0.06 ppm in BR 11 to 0.13 ppm in BRR1 dhan32 and BRR1 dhan41, while such values were ranged from 0.231 ppm in Kalizeera with 10 ppm As to 0.743 ppm in BRR1 dhan40 with applied 20 ppm As to soils. Among the 14 rice cultivars, the BRR1 dhan32 BR 11 and Najirshail appeared to be more tolerant to As contamination.

Key words: Rice, Yield, Arsenic Contamination, Grain Arsenic

1. INTRODUCTION

Arsenic (As) is an important geogenic contaminant found ubiquitously in the earth that threatening the health of 150million people worldwide [1-2]. The major sources of As exposure to humans includes drinking water and different food items,

especially rice. Exposure of As is associated with various chronic and acute health problems to humans that include skin lesions, cardiovascular diseases, diabetes, cancer, and so on [2]. The As bound to Fe hydroxides, oxyhydroxides, and oxides released into groundwater through biogeochemical processes and resulted in widespread As contamination in Bengal delta [3-4]. Geogenic As contaminated ground water are being used for rice cultivation in West Bengal and Bangladesh. This common practice has resulted in As build-up in soil through the years [5-6]. Background As concentration in Bangladesh soils is in the range of 4-8 mg As kg⁻¹. Research studies indicate that, long-term use of As contaminated water for irrigation purposes may result in elevated As concentration in soils and plants, and may lead to crop yield loss and elevated As concentration in cereals, vegetables and other agricultural products [7]. Irrigation using As contaminated groundwater results in significant loading of As in soil and as a consequence subsequent bioaccumulation in rice grains and elevated levels of As have also been shown to lead to substantial yield losses [8-9]. Rice can accumulate several-fold higher As than other cereal crops such as wheat and maize [2, 6, 10]. Every year approximately, 36.60 million tons of rice is produced from 11.42 million hectares of land of Bangladesh [11]. The As accumulation in rice grains varies significantly in different rice varieties [12]. The concentrations of As in different parts of all rice varieties increased significantly with the increase of its concentrations in soil. However, the identification of a suitable low As accumulating rice variety and its use in breeding programs for the development of a suitable rice variety is now time demanding. Under these circumstances, a successful solution can be to identify rice varieties that accumulate low As in grains.

2. MATERIAL AND METHODS

The experiment was conducted at the net-house of the Department of Soil Science of Bangladesh Agricultural University, Mymensingh, during the Aman season (Summer rice). The rice cultivars tested were BR 11, BRRI dhan30, BRRI dhan31, BRRI dhan32, BRRI dhan33, BRRI dhan34, BRRI dhan39, BRRI dhan40, BRRI dhan41, Binadhan-4, Najirshail, Biroi, Kalizeera, and Pajam were collected from Genetic Resources and Seed Division, BRRI, Gazipur. The soil for the experiment was silt loam in texture, pH 6.7, organic matter 2.0%, total N 0.12%, available P 9.8

ppm, exchangeable K 0.11 me%, available S 10.5 ppm and total As 2.1 ppm. Three levels of As contamination was induced in soil by adding viz. 0, 10 and 20 ppm and tested on 14 rice cultivars (10 modern and 4 local cultivars). The trial was set in a Completely Block Design with three replications. A total of 126 plastic pots were used in the experiment. Each plastic pot had the diameter of 43 cm and height 40 cm receiving 10 of kg soil. Thirty-five days old seedlings were transplanted in the pots and three hills were placed in an equal distance and each hill consisted of 3 seedlings. Urea, TSP, MoP, gypsum and zinc oxide were applied @ 100 mg kg⁻¹ N, 25 mg kg⁻¹ P, 50 mg kg⁻¹ K, 25 mg kg⁻¹ S and 5 mg kg⁻¹ Zn, respectively. The entire TSP, MoP, Gypsum and zinc were applied during the final pot preparation. Urea was applied in three equal splits; at 7, 30 and 60 days after transplanting. Intercultural operations were employed as and when necessary. Data were recorded on plant height, effective tillers pot⁻¹, panicle length, filled grains panicle⁻¹, 1000-grain weight, grain yield pot⁻¹ and straw yield pot⁻¹. The grain As content was determined after digesting the samples with HNO₃ and H₂O₂ 120°C followed by flow injection hydride generation atomic absorption spectrophotometer with UNICAM model No. 969 with hydride generator assembly using matrix-matched standards. The analysis of variance for various characters of the crop was done following the MSTATC computer program. The mean comparison of the treatments was made by the Duncan's Multiple Range Test (DMRT).

3. RESULTS AND DISCUSSION

3.1 Effective tillers

The effective tillers pot⁻¹ decreased significantly due to As contamination to soil. This reduction was significant for all of the varieties. The number of effective tillers pot⁻¹ in As control condition was 19.31 and that in 10 and 20 ppm As was 8.10 and 1.57, respectively (Table 1). The high yielding and local varieties selected for the experiment showed significant variation in terms of number of effective tillers pot⁻¹ (Table 1). The highest number of effective tillers pot⁻¹ was obtained in Najirshail with the value of 19.11 which was statistically similar to BRR1 dhan32. The interaction of As levels and different high yielding and local varieties selected for the experiment showed remarkable variation in terms of number of effective tillers pot⁻¹ (Table 2). Application of 10 and 20 ppm As significantly reduced the number of effective tillers

pot⁻¹. The highest number of effective tillers pot⁻¹ (30.37) was found in BRRi dhan32 with 0 ppm As application. The cultivar BRRi dhan41 failed to produce any effective tillers due to the application of 10 and 20 ppm As and the other seven rice varieties BR 11, BRRi dhan30, BRRi dhan33, BRRi dhan34, Binadhan-4, Biroi and Kalizeera failed to produce plants in 20 ppm As treatment. The corresponding reductions in tiller numbers also reported in Aman rice ranged from 0–22 % [13]. Similarly, a significant effect on plant height and effective tiller numbers has been observed when a different variety of rice is cultivated in soil artificially enriched with arsenic acid [14].

Table 1. Effects of added As and varieties on the yield contributing characters, grain and straw yield and their As concentration of different rice cultivars

Factors	Number of effective tillers pot ⁻¹	Number of filled grains pot ⁻¹	Grain yield pot ⁻¹ (g)	Straw yield pot ⁻¹ (g)	Grain As (ppm)
As added (ppm)					
0	19.31a	89.54a	42.27a	62.54a	0.10b
10	8.10b	50.12b	11.30b	19.47b	0.45a
20	1.33c	8.75c	2.12b	3.80b	0.64a
S.E. (±)	0.41	3.52	2.73	3.04	0.05
Cultivars					
BR 11	14.22b	48.31b	33.57a	39.74ab	0.31cd
BRRi dhan30	7.44dc	57.82ab	17.58cd	31.39b	0.26c
BRRi dhan31	10.45bc	67.33a	24.90b	29.09bcd	0.40bc
BRRi dhan32	15.01ab	58.71ab	12.00d	34.39b	0.34bcd
BRRi dhan33	9.00cd	35.95c	19.27cd	23.43d	0.31d
BRRi dhan34	5.22d	49.40b	8.28d	14.42d	0.22e
BRRi dhan39	10.00bc	36.00c	12.33d	16.69d	0.48ab
BRRi dhan40	5.34d	54.86b	20.02b	28.81b	0.49a
BRRi dhan41	1.44e	24.41d	2.91e	16.64d	0.13e
Binadhan-4	10.00c	48.86b	23.93b	27.21bcd	0.35bcd
Najirshail	19.11a	68.38a	32.55a	45.43a	0.35bcd
Biroi	10.00bc	37.05c	18.55c	31.92b	0.23d
Kalizeera	10.56bc	54.84b	19.56c	28.75b	0.15e
Pajam	7.45dc	50.59b	14.40d	32.55b	0.38bcd
S.E. (±)	1.21	3.78	1.52	2.07	0.05

In a column figures having same letters do not differ significantly at 5% level by DMRT

3.2 Filled grains

The filled grains pot^{-1} was decreased significantly by addition of As to soil. The number of filled grains pot^{-1} recorded in 0 ppm As was 89.54 and that in 10 and 20 ppm As was 50.12 and 8.75, respectively (Table 1). Higher As concentration causes grain sterility and grains panicle^{-1} were also significantly affected by the As treatments [13, 15]. All the rice cultivars showed significant variation in terms of number of filled grains pot^{-1} (Table 1). The highest number of filled grains pot^{-1} was obtained in Najirshail with the value of 68.38 which was statistically similar to BRR1 dhan31 and BRR1 dhan30. The interaction of As levels and different high yielding and local varieties selected for the experiment showed remarkable variation in terms of number of filled grains pot^{-1} (Table 2). Application of 10 and 20 ppm As significantly reduced the number of filled grains pot^{-1} . The highest number of filled grains pot^{-1} (114.31) was found in BRR1 dhan40 with 0 ppm As application.

Table 2. Interaction effects of added As and varieties on the yield contributing characters of different rice cultivars

Cultivars	Number of effective tillers pot^{-1} (g)			Number of filled grains pot^{-1} (g)		
	As0	As10	As20	As0	As10	As20
BR 11	28.33ab	14.33d-g	0.00m	93.03b-d	51.91j-n	0.00r
BRR1 dhan30	13.66d-h	8.67g-m	0.00m	103.57ab	69.88g-j	0.00r
BRR1 dhan31	20.67b-d	7.00g-m	3.67i-m	94.07b-d	81.85c-h	26.08o-q
BRR1 dhan32	30.37a	12.00e-i	2.67j-m	102.29ab	61.12i-l	12.73qr
BRR1 dhan33	20.67b-d	6.33g-m	0.00m	70.66f-j	37.20m-o	0.00r
BRR1 dhan34	10.33e-j	5.33h-m	0.00m	91.81b-e	56.40i-m	0.00r
BRR1 dhan39	18.00c-e	8.00g-m	4.00i-m	58.48i-l	33.56n-p	15.97p-r
BRR1 dhan40	12.00e-i	3.33i-m	0.68lm	114.31a	37.39m-o	12.89qr
BRR1 dhan41	4.33i-m	0.00m	0.00m	73.24d-i	0.00r	0.00r
Binadhan-4	20.68b-d	9.33f-l	0.00m	84.46b-g	62.13h-k	0.00r
Najirshail	30.33a	17.33c	9.67e-k	91.11b-f	70.52f-g	43.51k-o
Biroi	23.33a-c	6.68i-m	0.00m	84.56b-g	26.58o-q	0.00r
Kalizeera	24.00a-c	7.68g-m	0.00m	93.03a-c	71.50e-j	0.00r
Pajam	13.68b-h	7.33g-m	1.33k-m	98.88a-c	41.57l-o	11.33qr
S.E. (\pm)	1.501			3.658		

In a column figures having same letters do not differ significantly at 5% level by DMRT

3.3 Grain yield

Grain yield decreased significantly with As addition and levels of As application. Grain yield was considered as one of the most important parameters in variety selection in terms of As tolerance or sensitivity. The grain yield of rice cultivars was statistically affected by As contamination (Table 1). The application of 10 and 20 ppm As in soil produced 11.30 g and 2.12 g grain pot⁻¹, respectively, whereas in control (0 ppm As) the grain was 42.27 g pot⁻¹ i.e. yield reduction was almost 73.31% and 94.98 %, respectively over control. Previously, progressively decreased of rough rice yield from about 7 to 2 t/ha in 2006 and 9 to 3 t/ha in 2007 across the soil As gradient [9]. A number of authors [16-17] also reported yield reduction as a result of higher levels of As concentrations. Generally higher levels of As addition of As resulted in greater reduction in yield and the application of As to soil resulted in 80% yield reductions in T. Aman rice (BRRI dhan33), indicating a strong residual effect of added As on the next crop [13]. In winter and summer rice, above ground biomass of rice genotypes progressively decreased because of soil As gradient from 0 to 60 mg/kg [18]. The grain yield of rice varied significantly in different cultivars of Aman rice (Table 3). The highest grain yield (33.57 g pot⁻¹) was found in BR 11 which was statistically similar to Najirshail. On the other hand, the lowest grain yield (2.91 g pot⁻¹) was found in BRRI dhan41. Almost similar result represents by another study in Bangladesh [17]. The yield markedly declined at 20 mg kg⁻¹ As treatment and above. The grain yield was reduced to 93.1% for 50 mg kg⁻¹ As treatment, whereas its reduction was only 1.8% in 1.0 mg kg⁻¹ As treatment compared to the control. The interaction effect of added As and different rice varieties selected for the experiment significantly affected the grain yield (Table 3). The application of 10 and 20 ppm As reduced the grain yield of all cultivars of rice selected for this experiment. In As control treatment, the highest grain yield (75.66 g pot⁻¹) observed in BR 11. The lowest grain yield (8.73 g pot⁻¹) in As control treatment was found in BRRI dhan41. The addition of 10 ppm As significantly reduced the grain yield of rice compared to that obtained in As control treatment. The application of 20 ppm As significantly decreased the yield and most of the varieties failed to produce grain yield except BRRI dhan31, BRRI dhan32, BRRI dhan39, BRRI

dhan40 and Najirshail. Thus, it can be said that in respect to grain yield Najirshail is the most tolerant variety to As contamination.

3.4 Straw yield

Straw yield of rice cultivars was statistically affected by the As contamination (Table 1). The yield due to As contamination varied from 62.54 g pot⁻¹ in control (0 ppm As) to 19.47 g and 3.80 g pot⁻¹ in 10 and 20 ppm As treatment, respectively. Application of 10 and 20 ppm As to soil caused 64.87 % and 93.93 % straw yield reduction respectively of rice. Similarly straw yield was reduced by 7–64 % in the rice [13]. The straw yield of rice ranged from 14.42 g pot⁻¹ in BRR I dhan34 to 45.43 g pot⁻¹ in Najirshail (Table 1). The interaction effect of different rates of As and cultivars of rice was significant in producing the straw yield (Table 3). The application of 10 ppm As decreased the straw of all rice varieties but the degree of decrease in straw yield was quite variable. The application of 20 ppm As significantly decreased the yield and some of the varieties failed to produce straw yield viz. BRR I dhan33, BRR I dhan34, BRR I dhan41 and Binadhan-4. Considerable reduction in straw biomass in the higher As treatment recorded in this experiment primarily as a result of lower tillers pot⁻¹ and stunted rice plant. The same type of declining As uptake pattern for straw is found for one variety of rice grown in a glasshouse [19]. Exposure to As, causing a reduction in shoot biomass and/ or growth, was also reported by many research findings [20-21].

3.5 Grain arsenic

Arsenic concentration is an important consideration for the selection of As tolerant rice genotypes. Arsenic concentration in grains of rice cultivars was significantly increased due to application of different levels of As in soil (Table 1). The rate of increase was significant in all of the cultivars. Arsenic concentration in grain of control pot (0 ppm As) was 0.10 ppm and that in pot receiving 10 and 20 ppm As was 0.45 and 0.64 ppm, respectively. The As concentrations in different cultivars ranged from 0.13 ppm in BRR I dhan41 to 0.49 ppm in BRR I dhan40. There was a

significant interaction effect of different rates of As and cultivars of rice on As concentration of rice grains (Table 3). The application of 10 and 20 ppm As increased the grain As concentration of rice. All the varieties selected for the experiment showed higher As concentration in the pots receiving 10 and 20 ppm As in soil compared to As control pot. Among the treatment combinations $As_{20} \times$ BRRI dhan40 had the highest As concentration (0.743 ppm) in grain and followed by $As_{20} \times$ BRRI dhan39 and $As_{20} \times$ BRRI dhan31. The lowest As concentration (0.063 ppm) was found in $As_0 \times$ BR 11 treatment combination. Cultivation of rice in soils with elevated concentration has been reported a number of authors [22-23]. The As concentrations in rice grain vary widely depending on the cultivars, As status of soil and irrigation water [8, 16]. The grain As concentration increased from values of 0.22–0.32 $\mu\text{g g}^{-1}$ with the As_0 treatment to 0.59–0.81 $\mu\text{g g}^{-1}$ with the As_{20} treatment, but varied inconsistently between the varieties.

Table 3. Interaction effects of added As and cultivars on grain yield, straw yield and grain As concentration of different rice cultivars

Cultivars	Grain yield pot ⁻¹ (g)			Straw yield pot ⁻¹ (g)			Grain As (ppm)		
	As0	As10	As20	As0	As10	As20	As0	As10	As20
BR 11	75.66a	25.05f-h	0.00k	83.57a	32.79gh	2.87m-o	0.063e	0.559ab	-
BRR1 dhan30	39.40de	13.35g-k	0.00k	66.62b-e	25.99hi	1.57no	0.095e	0.417a-d	-
BRR1 dhan31	59.14bc	10.60g-k	4.95i-k	66.64b-e	14.06i-o	6.57k-o	0.105de	0.458a-c	0.625ab
BRR1 dhan32	17.25ab	17.26g-k	1.48i-k	80.08a-c	19.50h-m	3.58m-o	0.133d-e	0.333b-e	0.550ab
BRR1 dhan33	55.21bc	2.61i-k	0.00k	63.97c-e	6.31k-o	0.00o	0.097e	0.525a-c	-
BRR1 dhan34	17.54g-i	7.29i-k	0.00k	27.52g-i	15.75i-o	0.00o	0.108de	0.333b-e	-
BRR1 dhan39	26.19fg	9.40h-k	1.39i-k	32.76gh	13.49i-o	3.81l-o	0.108de	0.642ab	0.683ab
BRR1 dhan40	47.06de	10.01g-k	2.98i-k	61.54de	21.30h-k	3.59m-o	0.104de	0.625ab	0.743a
BRR1 dhan41	8.73h-k	0.00k	0.00k	42.40fg	7.53j-o	0.00o	0.128de	-	-
Binadhan-4	56.90bc	14.90g-k	0.00k	63.52de	18.12h-n	0.00o	0.116de	0.583ab	-
Najirshail	52.78cd	26.49fg	18.39g-i	81.46ab	34.18gh	20.65h-l	0.082e	0.333b-e	0.620ab
Biroi	49.24c-e	6.42i-k	0.00k	68.29a-e	23.78h-j	3.68m-o	0.075e	0.375a-e	-
Kalizeera	52.06cd	6.63i-k	0.00k	64.94c-e	21.31h-k	0.00o	0.071e	0.231c-e	-
Pajam	34.58ef	8.18i-k	0.43jk	72.25a-d	18.46h-n	6.93k-o	0.078e	0.450a-c	0.600ab
S.E. (±)	2.845			2.864			0.053		

In a column figures having same letters do not differ significantly at 5% level by DMRT

4. CONCLUSION

From this experiment, it was found that high level of As contamination to soil significantly affected the yield contributing characters viz. number of effective tillers number of filled grains of rice cultivars. The grain and straw yields of these rice cultivars were also markedly reduced by the artificial introduction of As into soil. A high As uptake in rice grain and rice straw was observed in this experiment. Rice cultivation in the naturally As-contaminated soil of Bangladesh may also reduce the yield of rice grain and straw. Among the fourteen rice cultivars, BRRI dhan32, BR 11 and Najirshail appeared to be tolerant to As contamination. Therefore, at high levels of As contamination, the varieties viz. BRRI dhan32, BR 11 and Najirshail may be considered for cultivation. However, further studies with varying levels of As contamination with different varieties in the field are needed to reach a final conclusion. Therefore, choosing less As-responsive rice genotypes to grow in As affected soil could effectively minimize As bioaccumulation in rice grains, leading to a reduction in health risks caused by As exposure.

REFERENCES

1. Majumder, S., and Banik, P. Geographical variation of arsenic distribution in paddy soil, rice and rice-based products: a meta-analytic approach and implications to human health. *Journal of Environmental Management*. 2019; 233:184–199.
2. Shikawa, S., Arai, T., and Makino, T. Agronomic strategies for reducing arsenic risk in rice, in *Arsenic Contamination in Asia*, eds H. Yamauchi and G. Sun (Singapore: Springer). 2019: 181–198.
3. Majumdar, A., and Bose, S. Toxicogenesis and metabolism of arsenic in rice and wheat plants with probable mitigation strategies, in *Arsenic: Risks of Exposure, Behavior in the Environment and Toxicology*, ed R. Knežević (Hauptpage, NY: Nova Science Publisher). 2017: 149–166.
4. Raessler, M. The arsenic contamination of drinking and groundwaters in Bangladesh: featuring biogeochemical aspects and implications on public health. *Arch. Environmental Contamination and Toxicology*. 2018; 75: 1–7.

5. Panaullah, G.M., Alam, T., Duxbury, J.M., Loeppert, R.H., Meisner, C. A., Lauren, J.G. and Waddington, S.R. Arsenic in Soil and Its Effect on the Growth of Rice at a High-Arsenic Site in Bangladesh. The 18th World Congress of Soil Science (July 9-15, 2006). Philadelphia, Pennsylvania, USA.
6. Upadhyay, M. K., Shukla, A., Yadav, P., and Srivastava, S. A review of arsenic in crops, vegetables, animals and food products. *Food Chemistry*. 2019; 276: 608–618.
7. Hossain, M.B., Jahiruddin, M., Loeppert, L.H., Panaullah, G.M., Islam, M. R. and Duxbury, J.M. The effects of iron plaque and phosphorus on yield and arsenic accumulation in rice. *Plant and Soil*. 2008; 317: 167-176.
8. Islam, S., Rahman, M.M., Islam, M.R. and Naidu, R. Effect of irrigation and genotypes towards reduction in arsenic load in rice. *Science of the Total Environment*. 2017; 609: 311-318.
9. Panaullah GM, Alam T, Hossain MB, Loeppert RH, Lauren JG, Meisner CA, Ahmed ZU, Duxbury JM. Arsenic toxicity to rice (*Oryza sativa* L.) in Bangladesh. *Plant and Soil*. 2009; 317:31-39.
10. Williams, P.N., Raab, A., Feldmann, J. and Meharg, A.A. High levels of arsenic in South Central US rice grain: consequences for human dietary exposure. *Environmental Science and Technology*. 2007; 41: 2178-2183.
11. BBS (Bangladesh Bureau of Statistics). Yearbook of Agricultural Statistics of Bangladesh. Bangladesh Bureau of Statistics. Government of Bangladesh. 2020.
12. Norton, G. J., Duan, G., Dasgupta, T., Islam, M. R., Lei, M., Zhu, Y..... Meharg, A. A. Environmental and genetic control of arsenic accumulation and speciation in rice grain: comparing a range of common cultivars grown in contaminated sites across Bangladesh, China, and India. *Environmental Science and Technology*. 2009; 43(21), 8381-8386.
13. Khan, M., Islam, M. R., Panaullah, G. M., Duxbury, J. M., Jahiruddin, M., & Loeppert, R. H. Accumulation of arsenic in soil and rice under wetland condition in Bangladesh. *Plant and Soil*. 2010; 333(1): 263-274.
14. Wang F-M, Chen Z-L, Zhang L, Gao Y-L, Sun Y-X. Arsenic uptake and accumulation in rice (*Oryza sativa* L.) at different growth stages following soil incorporation of roxarsone and arsanilic acid. *Plant Soil*. 2006; 285: 359–367.

15. Islam, S., Islam, M.R., Jahiruddin, M. and Islam, M.M. Genotypic variation for yield, yield attributes and grain arsenic content of boro rice grown in arsenic contaminated soils. *Bangladesh Journal of Crop Science*. 2008; 19(20):183-190.
16. Rahman, M.A., Hasegawa H., Mahfuzur Rahman M., Arifur Rahman M. and Miah M.A.M. Accumulation of arsenic in tissues of rice plant (*Oryza sativa* L.) and its distribution in fractions of rice grain. *Chemosphere*. 2007; 69(6): 942-948.
17. Azad, M., Kalam, A., Islam, M., Alam, A., Mahmud, H., Islam, M.A. and Rahman, M. Arsenic uptake and phytotoxicity of T-aman rice (*Oryza sativa* L.) grown in the As-amended soil of Bangladesh. *The Environmentalist*. 2009; 29(4): 436-440.
18. Iqbal, M., Rahman, G.K.M.M., Panaulah, G.M., Kabir, H. and Biswas, J.C. Influence of soil arsenic levels on biomass production and relationship the concentration of arsenic between Rice straw and grain. *Asian Journal of Soil Science and Plant Nutrition*. 2019; 5(2): 1-10.
19. Rahman M. A, Hasegawa H, Mahfuzur Rahman M, Mazid Miah MA, Tasmin A. Arsenic accumulation in rice (*Oryza sativa* L.): human exposure through food chain. *Ecotoxicology Environmental Safety*. 2008; 69: 317–324.
20. Imamul Huq, S.M., Shila, U.K. and Joarder, J.C. Arsenic mitigation strategy for rice using water regime management. *Land Contamination and Reclamation*. 2006; 14 (4): 805-812.
21. Islam, M.R., Islam, S., Jahiruddin, M. and Islam, M.A. Effects of irrigation water arsenic in the rice-rice cropping system. *Journal of Biological Science*. 2004; 4: 542-546.
22. Williams, P. N., Islam, M.R., Adomako, E.E., Raab, A., Hossain, S.A., Zhu, Y.G., Feldmann, J. and Meharg, A.A. Increase in rice grain As for regions of Bangladesh irrigating paddies with elevated arsenic in ground waters. *Environmental Science and Technology*. 2006; 40: 4903-4908.
23. Habiba, K.K., Jahiruddin, M., Islam, M.R. and Islam, S. Phosphorus increases arsenic toxicity in rice. *Bangladesh Journal of Seed Science and Technology*. 2007; 11(1&2): 31-36.