

## Original Research Article

### A comparative study between two different fractionation of High Dose Rate Brachytherapy in locally advanced carcinoma of Uterine Cervix after pelvic concurrent Chemoradiotherapy

#### ABSTRACT:

**Aims** **Background:** Uterine cervical cancer is the commonest form of gynecologic malignancy in Bangladesh. In locally advanced cases radiotherapy with a combination of external beam radiotherapy (EBRT) with concurrent chemotherapy and intracavitary brachytherapy (ICRT) is the mainstay of treatment. The **aim** of this study is to compare the treatment outcome and acute complications following treatment with 9 Gy (gray) in two fractions of ICRT with EBRT.

**Methodology:** A prospective study was carried out in the Department of Oncology, Bangabandhu Sheikh Mujib Medical University, Department of Radiation Oncology, National Institute of Cancer Research and Hospital, Dhaka during the period of 2017 to 2018. ~~Biopsy proven a total sixty patients of uterine cervical cancer were included and received EBRT and inj. Cisplatin 40 mg/m<sup>2</sup> weekly followed by HDR (High dose rate) ICRT. EBRT treatment was delivered by Co-60 teletherapy with 50 Gy in 25 fractions of 2Gy per fraction over a period of 5 weeks. Then two fractions of ICRT with 9 Gy in two fractions weekly for 2 weeks was given to the patients of arm A and 7 Gy in three fractions weekly for 3 weeks to arm B. Treatment was completed in 7 weeks in arm A and 8 weeks in arm B. Every patient was evaluated weekly during treatment. Then follow up was done at week 6, 12 and at 6 months after completion of treatment to see the response and toxicities.~~

**Results:** The mean age of patient at diagnosis was 50 years. During follow up at 6 month after completion of treatment, complete remission was 90% and 86% respectively for arm A and arm B. The overall complete response was 88%. The common toxicities associated with treatment were bladder and rectal toxicities, skin reaction and hematologic complications which were managed conservatively. During follow up after 6 months, rectal and bladder toxicities were similar in both arms.

Formatted: Font: Bold, Complex Script  
Font: Not Bold

Formatted: Justified

Formatted: Font: Bold, Complex Script  
Font: Not Bold

Formatted: Complex Script  
Font: Not Bold

Formatted: Font: Bold, Complex Script  
Font: Not Bold

Formatted: Font: Bold, Complex Script  
Font: Not Bold

Formatted: Complex Script  
Font: Not Bold

Formatted: Font: Bold, Complex Script  
Font: Not Bold

Formatted: Font: Bold, Complex Script  
Font: Not Bold

**Conclusion:** This study showed that a total dose of 18 Gy ICRT in two fractions of 9 Gy over 2 weeks is equally effective in short term local control with acceptable toxicities in comparison with a total dose of 21 Gy in three fractions of 7 Gy ICRT .

*Key words: Brachytherapy, Gray, Carcinoma cervix, Treatment response, Toxicity*

Formatted: Justified

## INTRODUCTION

Formatted: Justified

Uterine cervical cancer is the fourth most common cancer in women worldwide, and the seventh overall.<sup>1-6</sup> ~~with there is~~ an estimated 604,127 new cases in 2020.<sup>7</sup> A large majority (around 85%) of the global burden occurs in the less developed regions, where it accounts for almost 12% of all female cancers.<sup>7-11</sup> There were an estimated 341,831 deaths from cervical cancer worldwide in 2020, accounting for 7.5% of all female cancer deaths. Almost nine out of ten (87%) cervical cancer deaths occur in the less developed regions.<sup>4,7</sup>

Formatted: Superscript

In Bangladesh and many other developing countries, unfortunately the incidence and mortality rates of carcinoma uterine cervix remains high, predominantly due to late detection.

Formatted: Superscript

<sup>12-16</sup> There is no nationwide data of cancer in Bangladesh, but according to the cancer registry report of National Institute of Cancer Research and Hospital (2014) carcinoma cervix is the second leading cancer among women.<sup>2,17</sup> More than 80% cases usually present at an advanced stage with a high mortality rate. According to WHO report, age standardized incidence of cervical cancer in Bangladesh is 29.4 per lac women. Mortality from this disease is 17.9 per lac women. One third cervical cancer cases of world are found in South-Asian region, especially in India, Bangladesh and Pakistan.<sup>3</sup> In Bangladesh this is the commonest gynecological cancer, about 70% of the hospital cases of gynecological cancer are cervical cancer. About 11,956 new cervical cancer cases are diagnosed annually in Bangladesh. Cervical cancer is the 2<sup>nd</sup> most common female cancer in women aged 15 to 44 years in Bangladesh.<sup>3,18</sup>

Formatted: Superscript

A number of factors may influence the choice of treatment for carcinoma cervix, including tumor size, histology, stage, evidence of lymph node metastasis, risk factors for complications of surgery or radiotherapy and preference of patient.<sup>4,19-23</sup> Depending on stage, primary treatment consists of surgery, radiotherapy (RT) or a combination of radiotherapy and chemotherapy.<sup>24-27</sup>

Comment [D1]: Statements should not be started by abbreviations

Formatted: Superscript

Formatted: Highlight

Comment [D2]: Statements should not be started by abbreviations

Formatted: Highlight

<sup>24-27</sup> **CCRT** is the cornerstone of choice for FIGO stage IIB, IIIA, IIIB and IVA carcinoma of the cervix and is an excellent alternative to surgery to selected patients with stage IA, IB, or IIA diseases. **RT** for primary cervical cancer consists of a combination of external beam radiotherapy (EBRT) and intracavitary radiotherapy (ICRT), except stage IA disease where ICRT alone may be used.<sup>5, 28</sup> Primary radiotherapy is the treatment of choice

for locoregionally advanced disease with a careful balance of pelvic EBRT with high energy photons and ICRT, and must be administered at high doses (>80 -90Gy) and in a short time (<55 days) with the best technological resources available.<sup>629</sup>

Intracavitary brachytherapy delivers a high radiation dose directly to the tumor while sparing the surrounding normal tissues. HDR brachytherapy for carcinoma of the cervix is widely used because of its advantages of a short treatment time, rigid immobilization, patient convenience, and out-patient treatment.<sup>730</sup>

Treatment duration of cervical carcinoma with radiotherapy (EBRT and ICRT) should be as short as possible (within 8 weeks), and any planned or unplanned interruptions or delays should be avoided. Overall treatment time should not exceed 56 days including brachytherapy and should ideally be 49 days or less. Lower pelvic tumor control and survival rates are observed in invasive carcinoma of the uterine cervix when the overall treatment time in a course of irradiation is prolonged.<sup>831</sup>

Bangladesh is a developing country with a dense population. In Bangladesh 80% of carcinoma of uterine cervix presents at a fairly advanced stage with a high mortality rate. We have a variety of problems in managing cervical cancer including patient load, which is very high due to inadequate number of radiotherapy centre. So decreasing the insertion of HDR brachytherapy from 3 to 2 fractions gives similar result, then it will help patients by reducing treatment cost and decreasing repeated attendance in hospital, as well as reduction in patient load in radiotherapy centres to some extent and more patients will get chance of treatment.

So, the objective of this study was to observe and compare the local control of disease and complications following treatment of locally advanced carcinoma cervix with two fractions of HDR brachytherapy after standard concurrent chemoradiation.

## **MATERIALS AND METHODS**

It was a prospective analytical study and conducted in Department of Oncology, Bangabandhu Sheikh Mujib Medical University (BSMMU), Dhaka and Department of Radiation Oncology, National Institute of Cancer Research and Hospital (NICRH), Dhaka. The duration was one year from 2017 to 2018.

A total 60 patients with clinically diagnosed and histologically proven locally advanced squamous cell carcinoma of the uterine cervix (Stage IIB- Stage IVA) was selected as sample and random sampling was done by computer generated method to divide them in both arms. The aim of this study was to see the effects of different fractionation of brachytherapy as the primary outcome and secondary outcome was to compare the toxicities.

Ethical approval was taken from the institutional review board (IRB) of BSMMU, informed consent was taken from each patient before enrolling in the study.

For radiotherapy the target volume was whole pelvis encompassing the extent of primary tumor and the pelvic lymph nodes. During EBRT whole pelvis was treated with 2 Gy per fraction, 5 days in a week with a total dose of 50 Gy for 5 weeks in a Cobalt 60 teletherapy machine with SSD of 100 cm, Inj. cisplatin 40 mg/m<sup>2</sup> was given weekly to the patients on days 1, 8, 15, 22 and 29. After EBRT, all the patients of both arms were treated with HDR ICRT. A dose of 9 Gy per fraction, 2 fractions in 2 weeks for arm A and 7 Gy per fraction, a total of 3 fractions over 3 weeks for arm B to the point-A were given. A total ICRT dose of 18 Gy and 21 Gy were delivered for arm A and B respectively

**Response Evaluation Criteria for Solid Tumors (RECIST-~~response evaluation criteria for solid tumors~~)** criteria was followed to assess the treatment response. For toxicity assessment, 'Toxicity criteria of the Radiation Therapy Oncology Group (RTOG) and the European Organization for Research and Treatment of Cancer (EORTC) 1995' criteria was used. SPSS software (version 20) was used for data analysis. Chi square test was used for comparison of demographic variables and qualitative data.

## OBSERVATIONS AND RESULTS

Patients with locally advanced carcinoma of uterine cervix were enrolled in this study and demographic and baseline characteristics were comparable in both groups (table I),

**Table I: Patient characteristic (n=60)**

Characteristics	Arm A (n=30)	Arm B (n=30)
Age at diagnosis		
31 to 60	26	25
Above 60	4	5
Early age of marriage (≤16 year)	25	24
Lower Economic condition	18	19
Grand multiparous	17	15
<b>Sign and symptom</b>		
Post Coital Bleeding	25	24
Inter-menstrual Bleeding/Postmenopausal bleeding	20	18
Excessive per vaginal Discharge	27	28
Pelvic Pain	10	12
Dysuria	6	7
Rectal Pain	0	0
Loss Of Appetite	19	18
Anemia	22	23

Formatted: Justified

Formatted: Line spacing: 1.5 lines

Formatted: Centered, Line spacing: 1.5 lines

Formatted Table

Formatted: Line spacing: 1.5 lines

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: Bold, Complex Script  
Font: Bold

Formatted: Left, Line spacing: 1.5 lines


Formatted: Complex Script Font: Bold

Formatted: Justified

Most of the patients was diagnosed as stage IIB disease in both arms, 18(60%) and 16(53.33.0%) patients in Arm A and B respectively.12 patients(40.0%) in Arm-A & 13 patients(43.33%) in Arm-B were in Stage IIIB & there was only 1 patient from stage IVA in Arm-A (Figure 1).

Formatted: Centered

Figure 1: Graphical presentation of clinical stage of disease in both arms



Formatted: Justified

At 1<sup>st</sup> follow up, there were 18(60%) patients of Arm-A and 16 (53.33%) of Arm-B with stage IIB disease and complete response observed in 16 (53%) of Arm-A and 13 (43.3%) of Arm-B patients. There was no statistically significant difference in complete and partial response between two arms on the basis of staging (p>0.05).

Formatted: Centered, Indent: Before: -0.27 cm, After: -0.28 cm, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Space Before: 0 pt, After: 0 pt, Line spacing: 1.5 lines

Formatted: Space Before: 0 pt, After: 0 pt, Line spacing: 1.5 lines

Formatted Table

Formatted: Font: Bold, Complex Script  
Font: Bold

Formatted: Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Formatted: Font: Bold, Complex Script  
Font: Bold

Formatted: Indent: Before: 0.11 cm, Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Formatted: Font: Bold, Complex Script  
Font: Bold

Formatted: Indent: Before: 0.11 cm, Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Table II: Distribution of the patients on the basis of staging and response at 6 week after completion of treatment

Total Patients n= 30 ( each arm)	No. of patients N_(%)	Complete response N_(%)	Partial Response N_(%)	Progressive Disease N_(%)	p value
<b>Stage IIB</b>					
▶ Arm-A	18(60%)	16(53.33%)	02(6.7%)	00(0.0%)	0.530
▶ Arm-B	16(53.33%)	13(43.33%)	03(10%)	00(0.0%)	
<b>Stage IIIB</b>					
▶ Arm-A	12(40%)	11(36.6%)	01(3.33%)	00(0.0%)	0.587
▶ Arm-B	13(43.3%)	11(36.7%)	02(6.7%)	00(0.0%)	
<b>Stage IVA</b>					

Formatted: Line spacing: 1.5 lines	▶ Arm-A	0(0%)	00(0.0%)	00(0.0%)	00(0.0%)
Formatted: Line spacing: 1.5 lines	▶ Arm-B	01(3.33%)	00(0.0%)	01(0.0%)	00(0.0%)

Formatted: Justified

▶ Then response evaluation was done at 2<sup>nd</sup> follow up which was at 12<sup>th</sup> week after completion of treatment. Response was similar like 1<sup>st</sup> follow up and there was no progression of disease

Final follow up was done at 6 month (24 weeks) after completion of treatment and it was observed that 90% of patients had complete response in Arm A. In Arm B 86% had complete response. The overall complete remission was 88%. Statistical analysis revealed there was no significant difference but arithmetically this is proven that Arm-A patients had better response than Arm-B.

**Table III: Distribution of the patients according to treatment response at 6 month**

Formatted: Font: Bold, Complex Script  
Font: Bold

Formatted: Line spacing: 1.5 lines

Formatted Table

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: Bold, Complex Script  
Font: Bold

Formatted: Line spacing: 1.5 lines

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Line spacing: 1.5 lines

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Justified

Response	Arm A (total=30) N (%)	Arm B (total =30) N (%)	Total
Complete response	27(90)	26(86)	53(88)
Partial response	3(10)	4(13)	7(12)
Progressive disease	0	0	0

Formatted: Space After: 0 pt

Formatted Table

Formatted: Centered, Space After: 0 pt

Formatted: Centered, Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Centered, Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: Not Bold, Complex Script  
Font: Not Bold

The frequencies of common toxicities related to treatment shown on table IV which are not statistically significant. Table shows that overall radiotherapy related toxicities were more in Arm A than that of Arm B but statistically it was not significant and all the toxicities were managed by conservative treatment. Treatment discontinuation or hospitalization for toxicity management was not needed during treatment and follow-up period. Most of the patients suffered from rectal and bladder toxicities in both arms.

**Table IV: Distribution of patients according to acute toxicity**

Variables	Group (Total=30)		p value
	Arm-A [n (%)]	Arm-B [n (%)]	
<b>▶ Skin Reaction</b>			
▶ Grade 1	10(33)	9(30)	0.967
▶ Grade 2	8(27)	7(23)	
<b>▶ Bladder toxicity</b>			
▶ Grade-1	12 (40)	10 (33)	0.878
▶ Grade 2	8(26.66)	6(20)	

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Centered, Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: Not Bold, Complex Script Font: Not Bold

<b>Haematologic toxicities</b>			
Grade 1	10(33)	11(37)	0.473
Grade 2	5(17)	3(10)	
<b>Rectal toxicity/ Rectal discomfort</b>			
Grade-1	9 (30)	10 (33)	0.370
Grade 2	10(33.33)	6(20)	

Formatted: Justified

During follow up at 6 months after completion of treatment 3 patients in arm A and 2 patients in arm B developed grade II bladder toxicities and only 2 patients in arm A developed rectal grade II toxicities, there was no rectal toxicity in arm B. the late bladder and rectal toxicity was higher in arm A than arm B but it was not statistically significant (p value 0.2899).

**Table V: Distribution of patients according to toxicity at 3<sup>rd</sup> follow-up**

Formatted: Space After: 0 pt

Formatted Table

Formatted: Centered, Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Centered, Space After: 0 pt

Formatted: Space After: 0 pt

Toxicity	Arm A(total=30) N_(%)	Arm B (total=30) N_(%)	p value
Bladder	3(10)	2(7)	0.850
Rectal	2(7)	0	

**DISCUSSION**

Formatted: Justified

Diagnosed patients of locally advanced carcinoma cervix (stage IIB to IVA) of squamous cell variety were enrolled in this study. Etiological factors of carcinoma cervix are early age of marriage, low socioeconomic condition, multiparity, illiteracy, long term oral contraceptive pill use and this study correlates with previous findings.

For locally advanced carcinoma cervix radiotherapy is the main modality of treatment and intracavitary brachytherapy is an essential part of it. In recent years, owing to the obvious physical advantages of shortened treatment time and better geometric placement HDR

brachytherapy has gained popularity. No clear consensus of the appropriate number of fractions or appropriate dose per fraction has been reached. Various fractionation schemes have been used experimentally in search of the optimal technique. The number of fractions has varied from as low as 1 to as many as 16. The dose per fraction to point A has varied from 3 to 17 Gy/fraction.<sup>932</sup>

Our present study was done using the HDR Microselectron with an Iridium 192 source and the HDR MicroselectronNucleotron applicator. We used a dose of 9 Gy/fraction of 2 fractions in Arm A and 3 fraction of 7 Gy/fraction in Arm B. The effects and toxicities were observed during and upto six months after completion of treatment. Both clinical examination and pap's smear test was done to see the presence of any microscopic residual disease. Complete remission was observed in 90% of patients in arm A and it was 86% in arm B, the overall complete response was 88%. Statistical analysis revealed there was no significant difference but arithmetically this is proven that Arm-A patients had better response than Arm-B.

The most prevalent acute toxicities in both the arms were bladder and rectum related toxicities, skin reaction, vaginal mucositis. No patient in both arms developed grade III or grade IV toxicity and there was no interruption of treatment due to toxicity. Although treatment related toxicities were slightly more in arm A and was managed well but it was not statistically significant (p value <0.05).

During follow up after 6 months, only 10% of in arm A and 6% in arm B developed grade II bladder toxicities and only 6% in arm A developed rectal grade II toxicities, which is similar to findings of other studies.<sup>4033</sup>

Different studies have effectively shown that in choosing the number HDR fraction and consequently the dose per fraction, it is the prescribed dose received by the critical organs that is important. If the critical ~~organ receive~~organs receive a smaller percentage of the point A dose, we can use larger dose per fraction without increasing morbidity.<sup>4434</sup> The advantage of using fewer fractions is patient convenience and improved patient compliance. Two fractions of 9 Gy brachytherapy can be completed in 2 weeks whereas three fractions of 7 Gy needs 3 weeks to complete the treatment. Reducing the risk of multiple exposures to anesthetic agents and minimizing the number of hospital attendance makes this schedule cost effective. This schedule also reduces the patient load in radiotherapy centers to some extent which is a major consideration in a developing country like ours where radiotherapy canters are overburdened with cancer patients.

## CONCLUSION

Formatted: Justified

HDR brachytherapy- 2 fractions of 9 Gy after concurrent chemoradiotherapy is equally effective with the brachytherapy of 3 fractions of 7 Gy after concurrent chemoradiotherapy for the control of locally advanced carcinoma cervix but more convenient regarding time and cost.

Formatted: Justified, Space After: 12 pt, Line spacing: Double

## **RECOMMENDATIONS**

Formatted: Font: 11 pt, Bold, Complex Script Font: 11 pt

Formatted: Justified

## **CONSENT**

Written informed consent was taken from all patients

Formatted: Line spacing: 1.5 lines

## **REFERENCE**

- [1. Qalawa, Sh., Eldeeb, A., & Hassan, H. Young Adult Women's intention regarding breast and cervical cancer screening in Beni-Suef. Scientific Research Journal, 2015; 3\(3\): 11-24.](#)
- [2. Hassan H. Early Stage Cervical Cancer: Survivorship and Fertility preservation. American Research Journal of Oncology, 2020; 2\(1\): 1-3.](#)
- [3. Said S., Hassan H., Sarhan A. Effect of an Educational Intervention on Women's Knowledge and Attitude Regarding Cervical Cancer. American Journal of Nursing Research, 2018; 6\(2\): 59-66. doi: 10.12691/ajnr-6-2-4.](#)
- [4. Nady F., Said M., Youness E., Hassan H. Effect of Nursing Intervention Program on Quality of Life Improvement for Women Undergoing Gynecological and Breast Cancer Treatment. Assuit Scientific Nursing Journal, 2018; 6\(15\): 62-77.](#)
- [5. Nady F., El-Sherbiny M., Youness E., Hassan H. Effectiveness of Quality of Life Planned Teaching Program on Women Undergoing Gynecologic Cancer Treatment. American Research Journal of Oncology, 2018; 1\(1\): 1-17.](#)
- [6. Hassan H., Ali R., Abd El Salam S., Kamal H. Impact of an Educational Program on Sexual Dysfunction Associated With Cervical Cancer. Journal of Cancer Research and Treatment, 2021; 9\(2\): 22-31. DOI:10.12691/jcrt-9-2-1](#)
- [7. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA: a cancer journal for clinicians. 2021 May; 71\(3\): 209-49.](#)

Formatted: Justified

**Formatted:** Line spacing: 1.5 lines

8. Hassan H., Mohammed R., Ramadan S., Masaud H. Call for Alleviating Sexual Issues among Cervical Cancer Survivors' Women in Northern Upper Egypt. *Journal of Obstetrics Gynecology and Reproductive Sciences*, 2021; 5(3): 1-11. DOI: 10.31579/2578-8965/066

9. Atwa A., Hassan H., Ahmed S. The impact of a hospital-based awareness program on the knowledge of patients about breast cancer and cancer cervix. *International Journal of Studies in Nursing*, 2019; 4(1): 20-29. doi:10.20849/ijsn.v4i1.537.

10. Hassan H., Masaud H., Mohammed R., Ramadan S. Self-Knowledge and Body Image among Cervical Cancer Survivors' Women in Northern Upper Egypt. *Journal of Applied Health Sciences and Medicine*, 2021; 1(1): 1-12

11. Mohammed F., Shahin M., Youness E., Hassan H. Survivorship in Women Undergoing Gynecological and Breast Cancer Treatment in Upper Egypt: The Impact of Quality of Life Improvement Educational Program". *American Research Journal of Gynaecology*. 2018; 2(1): 1-28. doi:10.21694/2577-5928.18001

12. Ali R., Abd El Salam S., Kamal H., Hassan H. Women with Cervical Cancer: Impact of an Educational Program their Knowledge. *Journal of Obstetrics Gynecology and Reproductive Sciences*, 2021; 5(2): 1-8. DOI: 10.31579/ 2578-8965/063

13. Abd El Salam S., Hassan H., Kamal K., Ali R. Sexual Dysfunction of Women's Associated with Cervical Cancer. *Journal of Applied Health Sciences and Medicine*, 2021; 1(2): 12-27.

14. Hassan H., Ramadan S., Ali R., Kamal H. Sexual Issues among Cervical Cancer Survivors' Women in Northern Upper Egypt. *Journal of Advanced Trends in Basic and Applied Science*, 2021; 1(1): 1-11.

15. Zagloul M., Naser E., Hassan H. Cervical Cancer Knowledge, Attitude, and Practices: Educational Program Management for Female Workers at Port Said University. *International Journal of Studies in Nursing*, 2020; 5(3): 1-16. doi:10.20849/ijsn.v5i3.776.

16. Masaud H., Hassan H., Mohammed R., Ramadan S. Women's Sexual Distress Associated with Cervical Cancer. *Sumerianz Journal of Medical and Healthcare*, 2021; 4(1): 28-34. doi.org/10.47752/simh.41.28.34

17. Cancer Registry Report, National Institute of Cancer Research and Hospital, Mohakhali, Dhaka, 2012-2014, pp 08.

18. Bruni L, Barrionuevo-Rosas L, Albero G, Serrano B, Mena M, Gómez D, Muñoz J, Bosch FX, de Sanjosé S. ICO/IARC Information Centre on HPV and Cancer (HPV Information Centre). Human Papillomavirus and Related Diseases in Bangladesh. Summary Report 2017 July 27-: 6

**Formatted:** Complex Script Font: Not Bold

**Formatted:** Justified, Line spacing: 1.5 lines

**Formatted:** Font color: Auto, Complex Script Font: Not Bold

**Formatted:** Font color: Auto, Complex Script Font: Not Bold

**Formatted:** Line spacing: 1.5 lines

19. Masaud H., Abd Rabo R., Ramadan S. Hassan H. [Impact of Protocol of Nursing Intervention on Sexual Dysfunction among Women with Cervical Cancer. Journal of Nursing Science Benha University, 2021; 2\(2\): 203-224](#)

20. Hassan H., Mohammed R., Ramadan S., Masaud H. [Impact of an Educational Program on Sexual Issues among Cervical Cancer Survivors' Women in Northern Upper Egypt. Journal of Obstetrics Gynecology and Reproductive Sciences, 2021; 5\(1\): 1-16. DOI: 10.31579/2578-8965/061](#)

21. Ali R., Kamal H., Hassan H., Abd El Salam S. [Impact of an Educational Program on Sexual Distress Associated With Cervical Cancer. Journal of Applied Health Sciences and Medicine, 2021; 1\(1\): 30-42](#)

22. Mohamed A., Hassan H., Gamel W., Arafa A. [Awareness about breast and cervical cancers among nursing students in Beni-Suef University. Journal of Nursing Education and Practice, 2019; 9\(5\): 44-51. doi.org/10.5430/jnep.v9n5p44](#)

**Formatted:** Font color: Auto, Complex  
Script Font: Not Bold

423. Klopp AH, Eifel PJ. Chemoradiotherapy for cervical cancer in 2010. *Current oncology reports*; 2011 Feb 1; 13(1): 77-85.

**Formatted:** Justified, Line spacing:  
1.5 lines

24. Ramadan S., Hassan H., Masaud H., Mohammed R. [Women's Body Image Distress Associated with Cervical Cancer. Journal of Obstetrics Gynecology and Reproductive Sciences, 2021; 5\(3\): 1-6. DOI: 10.31579/2578-8965/062](#)

**Formatted:** Line spacing: 1.5 lines

25. Kamal H., Ali R., Abd El Salam S., Hassan H. [Self-Knowledge among Women with Cervical Cancer. Journal of Cancer Research and Treatment, 2021; 9\(1\): 12-21. DOI: 10.12691/jcrt-9-1-2](#)

26. Abd El Salam., Ali R., Hassan H., Kamal H. [Outcome of an Educational Program on Body Image Distress Associated with Cervical Cancer. Journal of Advanced Trends in Basic and Applied Science, 2021; 1\(1\): 12-20](#)

27. Hassan, H., Bayoumi, M., & Atwa, A. [Emotional Distress Associated with Gynecologic and Breast Cancer in Beni-Suef City. International Journal of Science and Research, 2016; 5\(2\): 1118-1129.](#)

**Formatted:** Font color: Auto, Complex  
Script Font: Not Bold

528. Dueñas-Gonzalez A, Cetina L, Mariscal I, de la Garza J. Modern management of locally advanced cervical carcinoma. *Cancer treatment reviews*; 2003 Oct 31.29(5):389-99.

**Formatted:** Justified, Line spacing:  
1.5 lines

629. Colombo N, Carinelli S, Colombo A, Marini C, Rollo D, Sessa C, ESMO Guidelines Working Group. Cervical cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology*; 2012 Oct 1.23(suppl\_7):vii27-32.

730. Pazdur R., Wagman L.D., Camhausen K.A., Hoskins W.J. (eds), *Cancer Management : A Multidisciplinary Approach*, 13th edition; 2012. UBM Medica LLC, New York

831. Barrett A, Dobbs J, and Roques T, Practical Radiotherapy Planning 4th Edition, CRC Press; 2009. 370-380.

932. Patel FD, Sharma SC, Negi PS, Ghoshal S, Gupta BD. Low dose rate vs. high dose rate brachytherapy in the treatment of carcinoma of the uterine cervix: a clinical trial. International Journal of Radiation Oncology\* Biology\* Physics; 1994 Jan 15. 28(2): 335-41.

4033. Patel FD, Rai B, Mallick I and Sharma SC, High-dose-rate brachytherapy in uterine cervical carcinoma. International Journal of Radiation Oncology\* Biology\* Physics; 2005.62(1), pp.125-130.

4434. Fowler JF. The radiobiology of brachytherapy. in: A.A. Martinez, C.G. Orton, R.F. Mould (Eds.) Brachytherapy HDR and LDR. Nucletron International, B.J, Leersum, Netherlands; 1990.121-137.



UNDER PEER REVIEW

Formatted: Font:

Formatted: Font color: Auto, Complex Script Font: Not Bold

Formatted: Font: Not Italic, Complex Script Font: Not Italic

Formatted: Font: Not Italic, Complex Script Font: Not Italic

Formatted: Complex Script Font: Not Bold

Formatted: Font: 10 pt, Complex Script Font: 10 pt

Formatted: Justified