

Original Research Article

Effect of Industrial Coconut Oil Effluent Irrigation on Germination, Growth and Yield of *Talinum fruticosum* (Talinaceae)

Abstract

The increase in food production has been a major concern to developing countries. Due to the high demand for certain crops especially vegetables, farmers resort to the use of waste water to cultivate these crops during the dry season. This research work aimed to investigate the effects of industrial coconut oil effluent (ICE) on the germination, growth and yield of *Talinum fruticosum* (water leaf). The physicochemical composition of the effluent was determined using standard analytical methods. Germination and vegetative growth (the plant height, number of leaves, leaf area and fresh weight) were studied on *T. fruticosum*. The plants were subjected to irrigation with 20%, 40%, 60%, 80%, 100% and 0% (control) concentrations of ICE. The experiment on the vegetative growth studies was carried out in a Completely Randomized Design (CRD). The data were collected every two weeks, subjected to Analysis of Variance (ANOVA) and means were compared by using Duncan's new multiple range test (DNMRT) at $P < 0.05$. The results of physicochemical analysis of ICE showed that most parameters were above FEPA standard limits except lead. Results from germination and vegetative studies revealed that mean values from treatment 0% (control) were significantly higher at $P < 0.05$ and that the germination of seeds and seedling growth gradually declined as the concentration of effluent increased. Therefore, adequate dilution of the effluent is required before disposal and reuse of ICE for irrigation purposes.

Keywords: Seedlings, *Talinum fruticosum*, Vegetable production, Waste water reuse, Water leaf.

INTRODUCTION

The increase in food production has been a major concern to Nigeria and many other tropical countries. Leafy vegetables are one of the foods obtained from plant sources and stand out as the most popular, being used in the preparation of most dishes eaten by man (Schippers, 2000). Nutritionally, *Talinum fruticosum* (L.) Juss. (water leaf) is high in crude protein (22.1%), ash (33.95%) and crude fiber (11-12%); rich source of vitamins A and C, lipids, proteins, β -carotene and minerals such as calcium, potassium and magnesium (Ezekwe *et al.*,

Comment [u1]: Some Journal guidelines for authors recommend cite $P < .05$

Formatted: Font: Italic

Comment [u2]: Scientific name and authority according International Plant Names Index (IPNI.org) and data base from Missouri Botanical Garden, (Tropicos.org)

2001), which are essential for the maintenance of human health (Bolaji *et al.*, 2007; Nya and Eka 2008; Osuagwu, 2008; Nya and Eka 2008). The consumption of vegetables in diet has been reported to protect the human body from degenerative diseases and their main protective action had been attributed to their antioxidants content (Swarna *et al.*, 2013). The continuous demand for water leaf has increased the need to cultivate it all year round. The use of effluent or waste water is an important alternative source of water for irrigation, especially during the dry season and in regions where water is scarce to meet up with the rising demand for fresh vegetables (Uaboi-Egbenni *et al.*, 2009; Ogunkunle *et al.*, 2013). The use of industrial effluents irrigation in the production of leafy vegetables is not a new concept and has been practiced in many developed and developing countries (Nweze and Eji, 2005; Nzekwe *et al.*, 2009). Effluents may be considered beneficial and harmful to agriculture and the environment. Waste water may contain nutrients that can be extensively used for irrigation, when treated or diluted properly (Ubochi *et al.*, 2019). Ivy *et al.* (2015) reported that proper treatment of waste water can deliver positive benefits to the farming community and society at large. Conversely, Kwaya *et al.* (2021) reported that most of the *Corchorus olitorius* L. (Malvaceae), irrigated with coconut oil (*Cocos nucifera* L., Arecaceae) effluent did not favour the plants' growth and development at higher concentrations (80 and 100%).

Formatted: Font: Italic

Coconut fruit (*Cocos nucifera* L.) is of domestic and industrial importance because of the oil obtained from the solid endosperm (meaty part of coconut fruit). The industrial process of producing coconut oil generates huge quantities of waste. During the extraction process, about 80% of the water results in coconut oil waste water. Most times, the waste water is discharged into the environment without proper treatment, while some urban farmers divert effluents to farm lands to irrigate their vegetable farms for adequate output (Fatoba *et al.*, 2011). Many investigators reported that effluents from distilleries, sugar industry, pulp and paper mills and textile industries cause a gradual decline in seed germination and seedling growth with increasing concentration of the effluent. However, the rate of inhibition was different for different seeds. Studies suggested that industrial effluents could be used safely for agricultural purposes with proper dilution, monitoring and use management practices to minimize any potential adverse effects on soil and plants (Medhi *et al.*, 2011; Narain *et al.*, 2012). Therefore, the study aims to determine the effect of industrial coconut oil effluent irrigation on germination, growth and yield of *Talinum fruticosum* L.

Comment [u3]: What about treatment of these waste water?

Materials and methods

Comment [u4]: In capital letters

Study location

The planting was carried out in the Botanic Garden at the University of Nigeria, Nsukka (UNN) of 7° 35' 00" E longitude and 6° 15' 00" N latitude geographic coordinates. Nsukka is both a commercial and agrarian town producing varieties of vegetable crops including *T. fruticosum*. The soil and plant samples were analyzed in the Department of Soil Science, and the Department of Crop Science respectively, of the University of Nigeria, Nsukka.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Collection of materials

Viable seeds of *Talinum fruticosum* were obtained from the Botanic Garden of the Department of Plant Science and Biotechnology, University of Nigeria Nsukka (UNN). Coconut oil effluent was collected from a coconut oil-producing company in Nsukka and analyzed at the Department of Biochemistry, UNN. The topsoil and sandy soil were collected from the Botanic Garden of the Department of Plant Science and Biotechnology, UNN and poultry manure was collected from the poultry farm at the Faculty of Agriculture, UNN.

Preparation of sSoil sSamples: The top garden soil, poultry manure and sandy soil were mixed in a ratio of 3:2:1 to formulate the amended soil. The sandy soil was added to facilitate drainage and easy uprooting of the seedlings for analysis. Two-kilogram aliquots of amended garden soil were weighed using an electronic digital weighing balance (Furi, model: FEJ-600) and put into 180 polythene bags.

Preparation of eEffluent: The various concentrations of coconut oil effluent used for the irrigation experiments were prepared as follows: 0% control (100 ml tap water), 20% concentration (20 ml effluent + 80 ml tap water), 40% concentration (40 ml effluent + 60 ml tap water), 60% concentration (60 ml effluent + 40 ml tap water), 80% concentration (80 ml effluent + 20 ml tap water), and 100% (100 ml effluent).

Nursery bed preparation

Two nursery baskets filled with amended soil were used. The seeds of *Talinum fruticosum* were nursed in the nursery baskets filled with amended soil samples and watered daily till the plants were ready for transplanting.



Plate 1: *Talinum fruticosum* growing in a nursery (four weeks after planting).

Comment [u5]: sowing?

Transplanting

The seedlings were transplanted four weeks after germination into 180 polythene bags each containing 2 kg of amended soil. Transplanting was done in the evening to avoid the effect of transpiration on the seedlings. The plants were watered with tap water at an interval of two days for two weeks for the seedlings to stabilize before the application of effluent. Thereafter, 100 ml of the effluent concentrations of 0%, 20%, 40%, 60%, 80% and 100% respectively were applied to the seedlings. Irrigation was done at an interval of two days for 12 weeks.

Comment [u6]: I suppose they are the same of polyethylene bags

Germination Studies

The germination studies were carried out in the Postgraduate Laboratory of the Department of Plant Science and Biotechnology, University of Nigeria Nsukka. Twenty seeds of *T. fruticosum* were sown in each of 18 sterilized Petri dishes lined with filter papers. Then, 2 ml of tap water (0%, control) and 2 ml each of effluent solution at 20%, 40%, 60%, 80% and 100% were respectively added in different labelled Petri dishes. One milliliter of tap water was added to the control and 1 ml effluent diluents were added to the various dilutions respectively, at two-day intervals to keep the plates moist. The experiment was replicated three times. They were left to stand for seven days and the treatments were monitored after sowing to know the treatment that can retard or enhance germination. Records of the following were taken:

- (i) Days of germination: The number of days it took the seeds to germinate was recorded.
- (ii) Germination percentage: This was calculated for the various treatments respectively following the equation presented by Omosun *et al.* (2008)

$$\text{Germination (\%)} = \frac{\text{total number of normal germinated seeds}}{\text{total number of sown seeds}} \times 100.$$

Comment [u7]: You can note as follows, it is easier

Germination (%) = 100 x (total number of normal germinated seeds / total number of sown seeds)

Chemical **a**Analysis of **I**ndustrial **c**oconut **o**il **e**ffluent

Some selected chemical properties such as nitrogen, phosphorus, copper, lead, chromium, and exchangeable cations (K, Ca and Mg) were determined by the leaching ~~method~~ method of official methods of analysis... (AOAC, 2002). The biological oxygen demand (BOD) and **c**Chemical oxygen demand (COD) were determined following Winkler's method as described by Horvitz and Latimer (2006). The pH of the samples was measured with a glass electrode in 1:2 sample-water ratio suspensions using a digital pH meter (Metrohm, Model: 872 Labomaga symbol).

Comment [u8]: It is not cited in References

Comment [u9]: It is not cited in References

Experimental **d**esign

The experiment was carried out in a Completely Randomized Design (CRD) comprising six treatments. Each treatment was replicated eighteen times given a total of 180 experimental units.

Statistical **a**nalysis

The data from each treatment were subjected to analysis of variance (ANOVA) and means were separated using Duncan's Multiple Range Test (DMRT) at $p < 0.05$ level of significance, using Statistical Package for Social Sciences (IBM SPSS) version 20.

Comment [u10]: See the Journal guide lines for authors, but correct expression would be $P < .05$

Formatted: Font: (Default) Times New Roman, 12 pt

Results

Comment [u11]: In capital letters

Chemical Composition of Industrial Coconut Oil Effluent (ICE)

The values of some chemical components of industrial coconut oil effluent (ICE) shown in Table 1 revealed that the data obtained from ICE (pH, BOD, COD, N, P, K, Cd, Cr) were above Federal Environmental Protection Agency (FEPA) accepted standards for irrigation water, except lead that was within acceptable limits. Fats and oils content was quite high and above and beyond the recommended standard for discharge on land. pH was acidic as well.

Table 1: Chemical analysis of industrial coconut oil effluent (ICE).

Properties	ICE	FEPA(1991) for Irrigation wWater
pH	4.0	6.5-9
DO_(mg/L)	8.50	NA
BOD (mg/L)	8,200	4.00
COD_(mg/L)	12,667	120
Fats & oil	20.34	10
Nitrogen (N) (mg/L)	9.5	0.10
Phosphate (PO ₄) (mg/L)	5.35	0.02
Potassium (K) (mg/L)	4.63	0.2
Magnesium (Mg) (mg/L)	33.40	0.50
Calcium (Ca) (mg/L)	26.50	0.20
Lead (Pb) (mg/L)	4.36	5.00
Cadmium (Cd) (mg/L)	1.14	0.01
Chromium (Cr) (mg/L)	0.82	0.10

Key: BOD= Biochemical oxygen demand; COD= Chemical oxygen demand; DO= Dissolved oxygen; ICE = Industrial Coconut oil Effluent, FEPA= Federal Environmental Protection Agency, NA = Not Available.

Seed Germination

The seed percentage germination gradually declined as the treatment increased (Table 2). The highest percentage germination (74%) was recorded for 0% (control) treatment, followed by treatment with 20% ICE (58%). Treatment with 100% ICE gave the least seed percentage germination of 8%.

Table 2: Percentage of seed germination with different Industrial Coconut Effluent (ICE) concentrations.

ICE Concentration	No. of Germinated Seeds	Percentage Germination
0% (control)	35	74%
20%	28	58%
40%	23	46%
60%	20	40%

Comment [u12]: I believe your analysis considered in first place the possibility of toxic and harmful effluent aspect, because I saw that you not included in your analysis Ni, Cu, Fe, Hg, etc. However, as you say the values are enough above those given by FEPA. It has enough Pb.

Comment [u13]: Values resalted are correct?

Formatted: Highlight

Formatted: Highlight

Comment [u14]: OR 0.02?

80%	9	18%
100%	4	8%

Effect of Effluent on Plant Growth and Yield of *T. fruticosum*

The impact of coconut oil effluent irrigation on plant height, number of leaves, leaf area and fresh and dry weight are shown in Tables 3, 4, 5, 6 and 7 respectively. From the 2nd to the 12th week after treatment (WAT) in all the treatments, 0% (control) treatment gave the highest ($P < 0.05$) mean plant height followed by 20% while 100% treatment had the lowest mean plant height (Table 3).

The control gave the highest number of leaves ($P < 0.05$) from the 2nd to 12th-WAT, while the 100% treatment had the lowest number (Table 4). At 4, 8, and 10 WAT, 20%, 40%, 60% and 80% treatments were the same for each week. However, at 6 WAT, 60%, 80%, and 100% treatments produced the same number of leaves which were significantly ($P < 0.05$) lower than the number of leaves produced by 20% and 40% treatments.

The highest mean leaf area was produced by the 0% (control) treatment from the 2nd to 12th WAT and this was significant at $P < 0.05$ while 80% and 100% treatments gave low leaf area (Table 5). The highest leaf fresh and dry weight were obtained in 0% treatment (control) followed by 20% treatment while 100% ICE treatment had the least (Tables 6, 7). The plant height, number of leaves, leaf area, stem girth, root length and fresh and dry weight of *Talinum fruticosum* grown in the control (0%) treatment were significantly higher ($P < 0.05$) than all the growth and yield parameters measured in ICE treated plants.

Table 3: Mean plant height (cm) of *Talinum fruticosum* irrigated with different concentrations of industrial coconut oil effluent (ICE).

ICE Treatments	Week after Treatment (WAT)					
	2	4	6	8	10	12
0% (control)	28.0 ± 1.16 ^a	30.7 ± 1.45 ^a	32.3 ± 1.45 ^a	35.7 ± 2.33 ^a	38.3 ± 2.40 ^a	42.0 ± 1.53 ^a
20%	26.0 ± 0.58 ^{ab}	28.0 ± 1.53 ^{ab}	29.3 ± 1.76 ^{ab}	30.7 ± 1.20 ^b	31.3 ± 1.20 ^b	34.3 ± 1.20 ^b
40%	25.0 ± 1.00 ^{bc}	26.3 ± 1.86 ^{bc}	26.7 ± 1.45 ^{bc}	27.7 ± 1.45 ^{bc}	28.7 ± 1.45 ^{bc}	31.7 ± 0.88 ^{bc}

60%	24.3 ± 0.88 ^{bc}	25.3 ± 0.88 ^{bc}	26.0 ± 1.53 ^{bc}	26.7 ± 1.67 ^{bc}	27.7 ± 1.67 ^{bc}	30.0 ± 2.52 ^{bc}
80%	23.7 ± 0.33 ^{bc}	24.7 ± 0.33 ^{bc}	25.3 ± 0.88 ^{bc}	26.0 ± 1.00 ^{bc}	27.0 ± 1.53 ^{bc}	27.7 ± 1.45 ^c
100%	23.0 ± 0.58 ^c	23.3 ± 1.20 ^c	24.3 ± 0.33 ^c	24.7 ± 0.33 ^c	25.0 ± 1.00 ^c	26.3 ± 1.86 ^c

Data are presented in means ± standard error. Means with different alphabets on the same column are significantly different using Duncan New Multiple Range Test (DNMRT) at P < 0.05.

Table 4: Mean number of leaves of *T. fruticosum* irrigated with different concentrations of ICE.

ICE Treatments	Week after Treatment(WAT)					
	2	4	6	8	10	12
0% (control)	15.0 ± 0.58 ^a	24.7 ± 1.20 ^a	28.0 ± 2.52 ^a	29.0 ± 2.08 ^a	30.0 ± 3.06 ^a	32.0 ± 3.06 ^a
20%	14.7 ± 0.88 ^a	21.3 ± 1.86 ^{ab}	22.7 ± 1.20 ^b	25.7 ± 0.88 ^{ab}	26.7 ± 0.88 ^{ab}	27.0 ± 0.58 ^b
40%	14.3 ± 0.88 ^a	19.7 ± 3.28 ^{abc}	21.0 ± 1.16 ^b	24.0 ± 2.08 ^{ab}	24.3 ± 1.76 ^{ab}	25.3 ± 0.33 ^{bc}
60%	14.3 ± 0.88 ^a	17.0 ± 2.65 ^{bc}	19.0 ± 0.58 ^{bc}	23.0 ± 0.58 ^b	24.0 ± 0.58 ^b	20.7 ± 1.76 ^{cd}
80%	14.0 ± 0.58 ^a	15.3 ± 1.45 ^{bc}	16.0 ± 1.16 ^c	21.7 ± 1.76 ^b	22.0 ± 2.08 ^{bc}	18.3 ± 0.88 ^d
100%	13.7 ± 1.20 ^a	14.0 ± 1.53 ^c	14.7 ± 1.76 ^c	15.3 ± 1.45 ^c	18.0 ± 1.16 ^c	17.7 ± 1.45 ^d

Data are presented in means ± standard error. Means with different alphabets on the same column are significantly different using Duncan New Multiple Range Test (DNMRT) at P < 0.05.

Table 5: Mean leaf area (cm²) of *T. fruticosum* irrigated with different concentrations of ICE.

Treatments	Week after Treatment (WAT)					
	2	4	6	8	10	12
0% (control)	39.07 ± 0.93 ^a	58.93 ± 3.85 ^a	62.07 ± 5.62 ^a	69.30 ± 8.04 ^a	83.67 ± 3.76 ^a	102.00 ± 6.43 ^a
20%	22.67 ± 2.29 ^b	41.67 ± 1.67 ^b	53.00 ± 4.73 ^a	53.93 ± 2.99 ^{ab}	54.60 ± 5.80 ^b	59.87 ± 5.39 ^b
40%	20.13 ± 6.06 ^{bc}	31.20 ± 7.02 ^{bc}	34.40 ± 2.81 ^b	40.37 ± 12.37 ^{bc}	43.33 ± 5.46 ^{bc}	52.33 ± 7.84 ^b
60%	17.27 ± 1.46 ^{bc}	21.30 ± 3.53 ^c	23.03 ± 1.35 ^{bc}	34.33 ± 6.69 ^{bc}	37.16 ± 6.02 ^{cd}	40.35 ± 7.71 ^{bc}
80%	15.83 ± 2.20 ^{bc}	16.47 ± 3.15 ^c	17.77 ± 3.12 ^c	22.65 ± 1.98 ^c	24.43 ± 2.72 ^{de}	29.18 ± 6.14 ^d
100%	12.37 ± 1.45 ^c	24.40 ± 6.84 ^c	31.03 ± 8.76 ^{bc}	19.14 ± 2.32 ^c	14.07 ± 2.07 ^c	10.30 ± 1.30 ^d

Data are presented in means ± standard error. Means with different alphabets on the same column are significantly different using Duncan New Multiple Range Test (DNMRT) at P < 0.05.

Table 6: Mean fresh weight (g) of *T. fruticosum* irrigated with different concentrations of ICE.

ICE Treatments	Week after Treatment(WAT)					
	2	4	6	8	10	12
0% (control)	26.82 ± 3.45 ^a	30.97 ± 3.16 ^a	40.68 ± 3.72 ^a	52.02 ± 3.79 ^a	60.85 ± 3.77 ^a	70.99 ± 1.06 ^a
20%	22.43 ± 0.36 ^{ab}	25.68 ± 3.73 ^{ab}	30.25 ± 3.08 ^{ab}	40.29 ± 2.70 ^b	52.71 ± 1.64 ^{ab}	66.29 ± 1.95 ^{ab}
40%	18.92 ± 3.90 ^{ab}	21.52 ± 0.74 ^{ab}	28.86 ± 3.47 ^{ab}	37.99 ± 3.78 ^{bc}	47.83 ± 2.83 ^{ab}	61.50 ± 1.76 ^{bc}
60%	17.33 ± 2.89 ^{ab}	18.31 ± 2.86 ^b	26.61 ± 3.98 ^b	33.69 ± 4.47 ^{bc}	43.62 ± 3.00 ^b	57.97 ± 1.93 ^{cd}
80%	15.69 ± 2.87 ^b	18.00 ± 3.00 ^b	23.36 ± 4.40 ^b	31.51 ± 1.84 ^{bc}	38.81 ± 1.42 ^b	53.65 ± 2.65 ^d
100%	14.79 ± 3.58 ^b	16.57 ± 3.23 ^b	20.97 ± 3.69 ^b	28.94 ± 0.82 ^c	20.47 ± 10.02 ^c	43.19 ± 3.65 ^e

Data are presented in means ± standard error. Means with different alphabets on the same column are significantly different using Duncan New Multiple Range Test (DNMRT) at P < 0.05.

Table 7: Mean leaf dry weight (g) of *T. fruticosum* irrigated with different concentrations of ICE.

ICE Treatment	Week after Treatment (WAT)					
	2	4	6	8	10	12
0%(control)	2.70 ± 0.34 ^a	3.27 ± 0.14 ^a	4.07 ± 0.37 ^a	5.20 ± 0.38 ^a	6.08 ± 0.38 ^a	7.10 ± 0.11 ^a
20%	2.25 ± 0.04 ^{ab}	2.84 ± 0.30 ^{ab}	3.03 ± 0.30 ^{ab}	4.03 ± 0.28 ^b	5.27 ± 0.16 ^{ab}	6.56 ± 0.22 ^{ab}
40%	1.89 ± 0.39 ^{ab}	2.15 ± 0.10 ^{bc}	2.89 ± 0.35 ^b	3.80 ± 0.38 ^{bc}	4.78 ± 0.28 ^{bc}	5.85 ± 0.13 ^{bc}
60%	1.73 ± 0.29 ^{ab}	1.83 ± 0.29 ^c	2.66 ± 0.40 ^b	3.37 ± 0.45 ^{bc}	4.36 ± 0.30 ^{cd}	5.66 ± 0.33 ^c
80%	1.71 ± 0.21 ^{ab}	1.80 ± 0.30 ^c	2.33 ± 0.44 ^b	3.15 ± 0.18 ^{bc}	3.88 ± 0.14 ^{de}	4.62 ± 0.31 ^d
100%	1.47 ± 0.35 ^b	1.65 ± 0.32 ^c	2.07 ± 0.12 ^b	2.90 ± 0.08 ^c	3.32 ± 0.27 ^e	3.76 ± 0.18 ^e

Data are presented in means ± standard error. Means with different alphabets on the same column are significantly different using Duncan New Multiple Range Test (DNMRT) at P < 0.05.

DISCUSSION

The chemical components of industrial coconut effluent had higher values than FEPA (1991) values for use as irrigation water for crops. The increase in BOD and COD was a result of a high level of pollutants in wastewater. This makes the effluent unsuitable for irrigation purposes. The pH value was acidic. The acidic nature of ICE may have originated from the corrosion of the machine used in processing the coconut fruits as well as organic acids found in fresh fruit (Mbagwu, 2003). Fats and oils content were above the recommended standard

Comment [u15]: Cite in References

for discharge on land. From the results, there was a high presence of heavy metals (Pd, Cd, Cr) in the effluents.

In this study, the results obtained from seed germination indicate that the lower concentrations of the effluent could promote germination of *T. fruticosum* seeds while higher concentrations of ICE had an inhibitory effect. The reduction in the germination of seeds irrigated with the ICE is due to the high amount of oil present in the effluent. This result is in line with the findings of Osubor and Oikeh (2013) that attributed the reduction of seed germination to the oily nature of palm oil industrial effluent on maize.

In this present study, there was a decline in plant height, number of leaves, leaf area and fresh and dry weight. The reduction in height of *T. fruticosum* was due to the high level of oil in the effluent which cause a deficiency in the availability of nutrients needed to maintain physiological processes involved in plant growth. These findings agree with the work of Omosun *et al.* (2008) who reported that the oily nature of the effluent may create an anaerobic condition in the soil which led to the reduction in uptake of nutrients needed for plant growth.

The decrease in the number of leaves may be attributed to the presence of heavy metals in the effluent. This is in line with the findings of Jadoon *et al.* (2013) who noted that heavy metal accumulation in vegetables had negative impacts on the growth and yield due to impaired uptake of water and nutrients.

Meanwhile, the mean leaf area declined with an increase in the application of effluent. This may be due to the toxic nature of the ICE as reported by Uaboi-Egbenni *et al.* (2009) who investigated the impact of industrial effluents (toiletries and plastic effluent) from different industries on *Abelmoschus esculentus* (L.) Moench (Malvaceae). They observed that effluents induced detrimental effects on leaf width and length. The fresh weight of *T. fruticosum* in the control was significantly higher than in other treatments due to the high COD, BOD, oil content of the effluent and heavy metals. This is in line with the reports of Narain *et al.* (2012) that different concentrations of distillery effluent influence the fresh and dry weight of *Cicer arietinum*.

CONCLUSION

In water-scarce countries, the reuse of effluents for irrigation of various vegetables is a very effective method to meet the demand, for adequate water for food supply. From this study, it was revealed that industrial coconut oil effluent inhibited plant germination and growth. Higher concentrations of industrial coconut effluent caused delayed germination, low fresh and dry weight, retarded growth and reduced leaf area. However, low concentrations (20% to 60%), favoured plant growth and development. Hence, proper management of ICE may result in healthy growth and yield of vegetables and improves soil fertility.

Comment [u16]: Really, do you think so? There is a saying: Take care, lest the cure be worse than the disease.

REFERENCES

America Public Health Association (APHA) (1998). *Standard Methods for the Examination of Water and Wastewater*. American Public Health Association, Washington, D.C. 1287pp.

Begum, A. S., Alam, J.M., Rahman, S. S. and Rahman, M. (2010). Effect of industrial effluents on germination and seedling growth of three leafy vegetables. *Bangladesh Journal of Science Industrial Research*, 45(2): 101 - 104.

Bernard, A. (2008). Cadmium and its adverse effects on human health. *Indian Journal of Medical Research*. 128(4): 557 – 564.

Bolaji, P. T., Komolafe, G. O. and Alli E. (2007). Drying characteristics of selected local vegetables. *Nigerian Food Journal*, 26: 138 - 142.

Dohare, D., and Rahul M. (2014). Biological treatment of edible oil refinery waste water using activated sludge process and sequencing batch reactors. A review. *International Journal of Engineering Science and Research Technology*, 3(12) 7 – 13.

Ekpo, I. A., Osuagwu, A. N. Okpako, E.C. Agbor, R. B., Ekanem, B. E. (2012). Physico-chemical and the morphological features of *Talinum triangulare* (water leaf), exposed to hydrocarbon polluted soil. *World Journal of Environmental Biosciences*, 2(1): 1-6.

Comment [u17]: All this References I didn't find cited in the text

Ezekwe, M.O., Besong, S. A., Igbokwe, P. E. (2001). Beneficial influence of purslane and waterleaf supplement to humans, *Federation of American Societies for Experimental Biology (FASEB) Journal*, 16: 639 - 643.

Fatoba, P. O., Olorunmaiye, K. S and Adepoju, A. O (2011). Effects of soaps and detergents wastes on seed germination, flowering and fruiting of tomato (*Lycopersicon esculentus*) and okra (*Abelmoschus esculentus*) plants. *Ecology, Environment and Conservation*, 17(1): 7- 11.

Fontem D. A. and Schippers R. R. (2004). *Talinum triangulare* (Jacq.) Wild. Innovation. <http://www.prota4u.org/search.asp>. Accessed 12th August, 2016.

Comment [u18]: I didn't find cited in the text of Ms.

Formatted: Font: Not Italic

Ivy, N., Hossain, M.K. and Hossain, M.L. (2015). Effects of industrial effluents on germination and early growth of selected agricultural crops. *Journal of Agronomy*, 14(11) 43-48.

Jadoon, K. B. I., Ali S., Jadoon, K. B. Q., Shakoob, B. M., Bharwana, A. S., and Farooq, A. M. (2013). Effects of irrigation with wastewater from different industries on vegetables grown in the vicinity of Faisalabad, Pakistan; A review. *International Research Journal of Plant Science*, 4 (6): 144 - 148.

Kwaya, V. B., Nweze, N. O., Wante, S. P., Barka, P. M. and Tahir A. A. (2021). Effect of coconut oil effluent on the morphological characteristics of *Corchorus olitorius*. *International Journal of Plant Research*, 11(1):7-13.

Mbagwu, F. O. (2003). Concentration of toxic substances in edible crops following use of wastewater in self-help farming practices by rural families in Obukpa, Nsukka. *International Journal of Agriculture and Biological Science*, 2(1): 31- 38.

Medhi, V. J., Jalukdar, A. K. and Deka, S. (2011). Impact of paper mill effluent on growth and development of certain agricultural crops. *Journal of Environmental Biology*. 32: 185 -188.

Narain, K., Bhat, M. M., Abhilash P. C. and Yunus, M. (2012). Impact of distillery effluent on seedling growth and pigment concentration of *Cicer arietinum* L., *Journal of Environmental Research and Development*.6: 3 – 9.

Nweze, N. O and Eji, M. O. (2005). Studies on the sewage effluent, river and tap water in the irrigation of *Amaranthus viridis* L. (Amaranthaceae). *Nigerian Journal of Plant Protection*, 22: 53 – 64.

Nya, E. J. and Eka, M. J. (2008). Genetic variability and heritability studies of desirable metric character in *Talinum triangulare* land races in South Eastern Nigeria. *Journal of Agronomy*, 6: 459 - 463.

Nzekwe, U., Nweze, N. O., Ugwoke, C.E.C., Nwamara, J. U and Omeh, S. C. (2009). Production of edible vegetable using sewage water. How safe for human health? Lettuce (*Lactuca sativa*) S. Khus. (*Ceompositae*). *International Journal of Biochemistry*, 1 (1): 82 – 84.

[Official methods of analysis of AOAC International.....Cite](#)

Ogunkunle, C.O., Abdulrahman, A.A., Aluko, T.A., Kolawole, O.S., Fatoba, P.O. and Oladele F.A. (2013). Anatomical response of *Amaranthus hybridus* Linn. as influenced by pharmaceutical effluents. *Notulae Scientia Biologicae*, 5 (4): 31-437.

Omosun, G., Markson, A. A., and Mbanasor, O. (2008). Growth and anatomy of *Amaranthus hybridus* as affected by different crude oil concentrations. *Journal of Scientific Research*, 3(1): 70 – 74.

- Osuagwu G. G. (2008). Proximate and vitamin content of Nigerian *Pterocarpus* species, *Nigerian Food Journal*, 26 (1): 21 – 24.
- Osubor, C. O. and Oikeh, E. I. (2013). Impact of palm oil mill effluent (POME) on soil physico-chemical properties and some vegetative parameters of maize (*Zea mays* L.) plant, *Nigerian Society for Experimental Biology*, 13(1-2): 19 – 22.
- Schippers, R.R. (2000). *African Indigenous Vegetables- An Overview of the Cultivated Spices*, NRI/ACP, EU Chatten, UK.
- Swarna, J., Lokeswari, T. S., Smita, M. and Ravindharan, R. (2013). Characterization and determination of in vitro-antioxidant potential of betalains from *Talinum triangulare* Lacquod Wild. *Journal of Food Chemistry*. 141: 4382-4390.
- Uaboi-Egbenni, P.O., Okolie P.N., Adejuyitan O.E., Sobande A. O. and O Akinyemi. O. (2009). Effect of industrial effluents on the growth and anatomical structures of *Abelmoschus esculentus* (L.) (okra). *Africa Journal of Biotechnology*, 8 (14): 3251 - 3260.
- Ubochi, K.C., Nweze, N.O. and Ojua E.O. (2019). Effects of coconut oil effluent (ICE) irrigation practices on some soil chemical properties and nutrients composition of *Talinum fruticosum* L. *International Journal of Ecology and Environmental Sciences* 45 (3): 303 – 310.