

### **A Review on Genetic Analysis of Rice (*Oryza Sativa* L.) Crop of yield contributing Traits**

#### **Abstract**

Rice is the primary source of energy for the international population. In any crop breeding programme, the presence of genetic variability is vital for broadening the gene pool. An excessive magnitude of variability in the population is required for selecting a genotype with excessively applicable traits. The heritability estimates along with the amount of genetic variability provide us with an idea of genetic gain due to choose. Specific direct developments (number of panicles/plants, filled grains/plant, and test weight) and indirect tendencies (tiller number/plant, plant peak) determine the yield component of rice. In the present overview, we draw a clear picture of how variability affects the selection method and also how yield is related to distinct direct and indirect developments. Genotypes show high variability, and suitable characters can be used in a similar breeding programme. A glimpse of molecular advances in rice breeding has also been integrated into the gift overview.

**Keywords:** Additive, gene action, selection; heritability; Rice Marker Assisted selection

#### **Introduction:**

[1] The rice (*Oryza sativa* L.) crop presents meals for the majority of the arena's populace, offering nearly 21% of its strength. [2] Rice is one of the most important food grains and an economically viable crop, ranking 2nd (2nd) next to wheat with 770.3 million tonnes produced. For the expression of a particular person, distinct gene movements are worrying. In the rice crop, di allele evaluation provides valuable information on gene action by utilising distinct yield and its contributing traits. In some specific scenarios, traditional breeding is likewise helpful to apprehend the type of gene movement involved in the expression of a trait. In rice crop besides breeding techniques, addition of herbal components peptides, nano particles. [3] With biological activities may also boom gene expression. Grain yield is enormously affected by diverse biotic (bacteria, viruses, fungi) and abiotic (water logging, salinity) stresses. Under the complicated environmental situation in any crop breeding

programme, direct selection for yield is blunder-prone and creates confusion. As a result, the appropriate selection strategy is dependent on the genetic variability present in the population as well as the relationship between various morpho-agronomic developments and yield. Estimation of the correlation coefficient offers a concept for approximately dating the extraordinary developments in grain yield. affiliation analysis helps plant breeders to find superior genotypes by specialising in perfect tendencies. So, the prevailing paper gives an overview of the concept of genetic diversity and additionally describes how the yield issue is decided through distinct direct and indirect results. [4] Amino acids natural extracts and vital oils.

### **GENETIC VARIABILITY IN RICE**

Any breeding program, especially, depends on the quantity of genetic variability present in the population. A man's or woman's phenotypic observable variant is the end result of genetic variability and environmental variability. For any crop development programme, the estimation of genetic variability is an important requirement. As a way to discover the genetic and non-genetic components of a populace, partitioning of overall phenotypic variability is critical for any successful breeding programme. A study was conducted to evaluate 35 rice genotypes for genetic variability, genetic development, and heritability for yield and its associated traits. For developments like days to panicle initiation, plant peak, days to 50% flowering, variety of panicle or plant, panicle length (cm), and range of spikelets or panicles, an extensive range of variants is changed into "found. [5] A significant distinction was discovered among numerous genotypes in a look at with rice crop by taking into consideration 13 (13) characters to estimate the nature and significance of variability, genetic strength, and heritability. In a study with rice, comparable large genetic variability was additionally discovered for all of the characters. Slight genotypic and phenotypic coefficients of variance become observed in trends like range of grains/panicle, productive tillers/plant, and seed yield/plant. [6] In a latest examine variety analysis became carried out with forty (40) wild rice germplasm traces for twelve (12) Quantitative characters the usage of D2 analysis. Seven (7) clusters were made based on the overall performance of genotypes, wherein Cluster I own the largest (32) quantity of genotypes, followed by Cluster IV and different clusters. [7] To examine the character association and genetic variability, observations have been recorded for 49 rice cultivars. The focus of the investigation shifted to individual association in order to determine the relationship between yield and its associated traits. Significant variations among genotypes were discovered after thinking

about all the developments. For traits like a wide variety of grains or panicle types, grain yield, and check weight, PCV and GCV became determined to be slight to excessive [8]. Estimation of correlation changed into additionally executed alongside with genetic variability and heritability for one hundred rice germplasms by the usage of alpha lattice design having three replications. ANOVA revealed that tremendous variations have been observed for all of the tendencies studied.

### **CORRELATION AND PATH COEFFICIENT ANALYSIS IN RICE**

[9] It is critical to understand the feasibility of combining multiple trends rather than selecting secondary trends for the genetic advantage of essential (primary) traits [10]. The goal of breeding programmes for increasing rice production is to use a more genetically diverse population as well as appropriate selection techniques to boost yield by its elemental characters. Correlation evaluation helps in figuring out the relationship between yield and its component trends [11], Whereas route evaluation is used to divide the influence of characters into direct and indirect effects. In course analysis (a statistical device), a direction chart is used for estimation of relationships between prediction variables and reaction variables. [12] Yield is a complicated man or woman who is linked to its element tendencies (quantity of panicle or plant, range of grains or spikes, weight, and so on). [13,14]. The overall performance of the rice crop depends on the relationship among yield and its related important tendencies. In such circumstances, choice is ineffective if we focus on the yield trait. Consequently, course analysis and the correlation coefficient facilitate segmenting the yield-contributing developments into direct effect and oblique impact components [15]. Course evaluation is also use as a spread criterion for improvement of rice crop [16]. An effective correlation has a look at became found in rice crop for grain yield and its associated developments. According to one study, increasing the panicle duration and panicle range per plant can improve trait 1000 grain weight [17]. Another look at the excellent correlation of grain yield with related trends; trait one thousand grain weight can be advanced by increasing the panicle period and panicle variety according to plant. Kumar et al. [18], Khush [19], Bhutta et al. [20], and Chhange and Devi [21] All found a comparable correlation. Facts about trait relationships enable plant breeders to locate suitable genotypes in various environmental conditions [22]

### **MODERN APPROACHES IN RICE BREEDING**

Molecular breeding of rice for excessive yield, stepped-forward grain nice, and excessive environmental adaptability is essential for feeding the sector's swiftly growing population [8]. To meet the protection of meals delivered by 2030, crop yield per unit region must be increased by 50% [23]. Excessive yield is the number one goal pursued by rice breeders. For the future rice development program, tremendous rice breeding using molecular design in the model of ideal plant architecture is critical [24]. A few current studies with aromatic rice indicate variations in aroma when they're grown in areas other than their normal, traditional developing regions [25]. In the Biriya Cheera variety, aroma is manufactured because of the presence of the BADH2 gene, which is also present within the famously scented Basmati rice [26]. For increasing the primary breeding goal, which is grain yield, distinct related tendencies like grain quantity in step with panicle, grain form, a wide variety of tillers, leaf perspective, and leaf length need to be enhanced. Gene OsbHLH079 is related to the management of leaf attitude and grain shape in rice via brassinosteroid biosynthesis and signalling pathways [27]. A study changed into one conducted on grain form and grain weight to discover the effect of 7 genes: GS2, GS3, GS5, GW5, GS7, SLG7, and GW8. For the development of traits such as grain duration, grain thickness, grain width, and thousand grain weights, the genetic effect differed from what was expected for each allelic mixture [28]. OsCYP96B4 is a semi-dwarfism gene found in rice plants that influences the content of amino acids, aminobutyrate (GABA), saccharides, and other secondary metabolites. Molecular observe on heterosis for unique characters like plant height, days to heading and grain yield have been additionally performed [29-32]. Parental material became two (2) subspecies of rice (*O. sativa* spp. Indica and *O. sativa* spp. Japonica), and eight QTLs were detected inside the loci [33]. The CRISPR/Cas9 technique was used to perceive the OsEWL4 gene, which is a regulator of individual plant biomass and tiller variety. The technique employs a centred mutation of rice FLW genes [34]. Overexpression and knockdown transformants of the OsNAR2.1 gene changed rice production as anticipated. With the exchange of expression level of OsNAR2.1 gene adjustments the worldwide methylation at whole genome level which in the long run modifications plant peak and grain yield [35]. Any other have a look at located out that a big set of genes are expressed differentially at the shoot apical meristem and regulate traits like number of tillers in keeping with plant and panicle numbers in rice plant [36]. Most of the above novel techniques are used to increase the yield ability of rice crops through expertise in molecular mechanisms.

## **CONCLUSION**

Enhancement of yield is the top objective in any rice development programme. Genetic variability with excessive slender feel heritability should be present in a populace for powerful choice of economically important traits. Proper dissection of nature and the significance of genetic variability are critical for the selection of traits desired by farmers, such as large seeds and high yield ability, as well as traits preferred by plant breeders, such as resistance to extreme biotic and abiotic pressure. A have a look at on relationship among yield and its extraordinary component trends assist plant breeders to go for direct and oblique selection. Specific molecular strategies are used to discover the genes responsible for the expression of traits. So extra emphasise should receive to characterise diverse germplasm with the aid of using different molecular tactics so they can be used in destiny breeding programme correctly.

#### References:

- [1] Saleh MM, Salem KF, Elabd AB. Definition of selection criterion using correlation and path coefficient analysis in rice (*Oryza sativa* L.) genotypes. Bulletin of the National Research Centre. 2020 Dec;44(1):1-6.
- [2] Baroudy AAE, Ali A, Mohamed ES, Moghanm FS, Shokr MS, Savin I, Lasaponara R. Modeling land suitability for rice crop using remote sensing and soil quality indicators: the case study of the Nile delta. Sustainability. 2020;12(22):9653.
- [3] Saad AM, El-Saadony MT, El-Tahan AM, Sayed S, Moustafa MA, Taha AE, Ramadan MM. Polyphenolic extracts from pomegranate and watermelon wastes as substrate to fabricate sustainable silver nanoparticles with larvicidal effect against *Spodoptera littoralis*. Saudi Journal of Biological Sciences. 2021;28(10):5674-5683.
- [4] El-Sobki AE, Saad AM, El-Saadony MT, El-Tahan AM, Taha AE, Aljuaid BS, Salem RE. Fluctuation in amino acids content in *Triticum aestivum* L. cultivars as an indicator on the impact of post-emergence herbicides in controlling weeds. Saudi journal of biological sciences. 2021;28(11): 6332-6338.
- [5] Chhaya R, Nilanjaya LP, Sethi S, Kumari P, Kushwaha N, Zala RG, Jaiswal A. Genetic divergence studies for agro-morphological and yield attributing traits in locally adopted rice varieties of Bihar;2021.
- [6] Divya J, Raju CD, Rao LS, Aparna K. Genetic Variability Studies in genetically diverse rice genotypes. Int. J. Curr. Microbiol. App. Sci. 2018;7(4):1169-1172.

- [7] Koli GK, Upadhyay S, Rathi SR, Gupta S, Kumar K, Singh PK, Sinha B. Genetic diversity analysis for yield and yield attributing traits of wild germplasm in rice. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(1):949-951.
- [8] Sreedhar S, Reddy UR. Association studies for yield and its traits in rice (*Oryza sativa* L.) genotypes. *International J. of Current Microbiology and Applied Sci*. 2019;8(1):2337-42.
- [9] Adjah, Kossi Lorimpo, et al. "Genetic variability, heritability and correlations for milling and grain appearance qualities in some accessions of rice (*Oryza sativa*L.)." *Physiology and molecular biology of plants*. 2020;26.6:1309-1317.
- [10] Akatwijuka R, Rubaihayo PR, Odong TL. Correlations and path analysis of yield traits in sorghum grown in southwestern highlands of Uganda. *African Crop Science Journal*. 2019 Aug 23;27(3):437-44.
- [11] Saha SR, Hassan L, Haque MA, Islam MM, Rasel M. Genetic variability, heritability, correlation and path analyses of yield components in traditional rice (*Oryza sativa* L.) landraces: Variability and traits association in rice. *Journal of the Bangladesh Agricultural University*. 2019 Mar 28;17(1):26-32.
- [12] Mongiano G, Titone P, Pagnoncelli S, Sacco D, Tamborini L, Pilu R, Bregaglio S. Phenotypic variability in Italian rice germplasm. *European Journal of Agronomy*. 2020 Oct 1; 120:126131.
- [13] Zhou J, Xin X, He Y, Chen H, Li Q, Tang X, Zhong Z, Deng K, Zheng X, Akher SA, Cai G. Multiplex QTL editing of grain-related genes improves yield in elite rice varieties. *Plant cell reports*. 2019 Apr;38(4):475-85.
- [14] Zhou Y, Tao Y, Yuan Y, Zhang Y, Miao J, Zhang R, Liang G. Characterisation of a novel quantitative trait locus, GN4-1, for grain number and yield in rice (*Oryza sativa* L.). *Theoretical and applied genetics*. 2018;131(3):637-648.
- [15] Jayasudha S, Majhi PK, Singh DK, Korada M, Khaire AR, Singh P, Singh SK. Genetic Variability, Character Association and Path Analysis for Yield and its Related Traits in Rice (*Oryza sativa* L.) Genotypes. *International Journal of Plant & Soil Science*. 2021 Dec 22:437-46.
- [16] Xu F, Wang Z, Lu G, Zeng R, Que Y. Sugarcane ratooning ability: research status, shortcomings, and prospects. *Biology*. 2021 Oct 15;10(10):1052.

- [17] Sarker U, Oba S, Ercisli S, Assouguem A, Alotaibi A, Ullah R. Bioactive phytochemicals and quenching activity of radicals in selected drought-resistant *Amaranthus tricolor* vegetable Amaranth. *Antioxidants*. 2022 Mar 17;11(3):578.
- [18] Kumar S, Bhuvanewari S, Devi EL, Sharma SK, Ansari MA, Singh IM, et al. Estimation of genetic variability, correlation and path analysis in short duration rice genotypes of Manipur. *Journal of Agri Search*. 2017;4(2):112-118.
- [19] Asibi AE, Chai Q, Coulter JA. Rice blast: A disease with implications for global food security. *Agronomy*. 2019 Aug 15;9(8):451.
- [20] Bhutta MA, Munir SANA, Qureshi, MK, Shahzad AN., Aslam KASHIF, Manzoor H, Shabir G. Correlation and path analysis of morphological parameters contributing to yield in rice (*Oryza sativa*) under drought stress. *Pak J Bot*. 2019;51(1):73-80.
- [21] Chhangte L, Devi TR. Correlation and path analysis studies in aromatic rice germplasm of North-East region of India. *The Pharma Innovation Journal*. 2019;8(10):01-04.
- [22] Gonçalves DDL, Barelli MAA, Oliveira TCD, Santos PRJD, Silva CRD, Poletine J. P, Neves LG. Genetic correlation and path analysis of common bean collected from Caceres Mato Grosso State, Brazil. *Ciência Rural*. 2017;47(8)
- [23] Tang R, Cui D, Zhou J, Li W, Ma X, Han B, Guo X, Zhao Z, Han L. Comparative analysis of genetic diversity of rice (*Oryza sativa* L.) varieties cultivated in different periods in China. *Genetic Resources and Crop Evolution*. 2021 Apr;68(4):1439-51.
- [24] Tang R, Cui D, Zhou J, Li W, Ma X, Han B, Guo X, Zhao Z, Han L. Comparative analysis of genetic diversity of rice (*Oryza sativa* L.) varieties cultivated in different periods in China. *Genetic Resources and Crop Evolution*. 2021 Apr;68(4):1439-51.
- [25] Sansenya S, Hua Y, Chumanee S. The correlation between 2-acetyl-1-pyrroline content, biological compounds and molecular characterization to the aroma intensities of Thai local rice. *Journal of Oleo Science*. 2018;67(7):893-904.
- [26] Veerabhadraswamy M, Sindhumole P, Mathew D, Shylaja MR, Joseph J, Augustine R, James, D. Molecular analysis of aroma gene (*BADH2*) in Biriyanicheera: A tropical aromatic rice genotype from Kerala, India. *Molecular Biology Reports*. 2022;1-7.
- [27] Seo H, Kim SH, Lee, BD, Lim JH, Lee SJ, An G, Paek NC. The rice basic Helix–

Loop–Helix 79 (OsBHLH079) determines leaf angle and grain shape. *International journal of molecular sciences*.2020;21(6):2090.

[28] Gull S, Haider Z, Gu H, Raza Khan RA, Miao J, Wenchen T, Liang G. InDel marker based estimation of multi-gene allele contribution and genetic variations for grain size and weight in rice (*Oryza sativa* L.). *International journal of molecular sciences*. 2019;20(19):4824.

[29] Abd El-Aty MS, Abo-Youssef MI, Galal AA, Salama AM, Salama, AA, El-Shehawi, A. M, El-Tahan AM. Genetic behavior of earliness and yield traits of some rice (*Oryza sativa* L.) genotypes. *Saudi Journal of Biological Sciences*; 2021.

[30] Hassanin AA, Saad AM, Bardisi EA, Salama A, Sitohy MZ. Transfer of anthocyanin accumulating delila and roseal genes from the transgenic tomato micro-tom cultivar to moneymaker cultivar by conventional breeding. *Journal of Agricultural and Food Chemistry*. 2020;68(39):10741-10749.

[31] Tang R, Cui D, Zhou J, Li W, Ma X, Han B, Guo X, Zhao Z, Han L. Comparative analysis of genetic diversity of rice (*Oryza sativa* L.) varieties cultivated in different periods in China. *Genetic Resources and Crop Evolution*. 2021 Apr;68(4):1439-51.

[32] Saleh MM, Salem KF, Elabd AB. Definition of selection criterion using correlation and path coefficient analysis in rice (*Oryza sativa* L.) genotypes. *Bulletin of the National Research Centre*. 2020;44(1):1-6.

[33] Yu Y, Zhu M, Cui Y, Liu Y, Li Z, Jiang N, Sui G. Genome sequence and QTL analyses using backcross recombinant inbred lines (BILs) and BILF1 lines uncover multiple heterosis-related loci. *International Journal of Molecular Sciences*. 2020;21(3):780.

[34] Gao Q, Li G, Sun H, Xu M, Wang H, Ji J, Zhao X. Targeted mutagenesis of the rice FW 2.2-like gene family using the CRISPR/Cas9 system reveals OsFWL4 as a regulator of tiller number and plant yield in rice. *International journal of molecular sciences*. 2020;21(3):809.

[35] Fan X, Chen J, Wu Y, Teo C, Xu G, Fan X. Genetic and global epigenetic modification, which determines the phenotype of transgenic rice?. *International Journal of Molecular Sciences*. 2020;21(5):1819.

[36] Zhang Q, Shen L, Wang Z, Hu G, Ren D, Hu J, Qian Q. OsCAF1, a CRM domain containing protein, influences chloroplast development. *International journal of molecular sciences*. 2019;20(18):4386.

UNDER PEER REVIEW