

## **Case study**

**Non surgical management of immature permanent teeth with open apex and large periapical lesion- 8 months follow up.**

### **Abstract**

**Aim:** The aim of this case report is to present a non surgical management of immature permanent maxillary central incisor with open apex and large periapical lesion within 8 months follow up.

**Methods & Materials:** The traumatic case was presented where the calcium hydroxide and iodoform paste (Metapex®) was placed in the root canals of immature permanent teeth with open apex and large periapical lesion. The teeth involved were evaluated radiographically at regular intervals for the 8 months after placement of the paste. At the end of 8 month the case showed continued root growth and the apical closure(apexification) with no evidence of periapical radiolucency. Conventional endodontic treatment was then performed.

**Conclusion:** In this clinical case, the calcium hydroxide and iodoform paste (Metapex®) was found to induce apical closure when assessed radiographically and clinically . After a period of 8 months the case showed continued root growth.

**Keywords:** Metapex, Calcium hydroxide, Apexification etc.

### **INTRODUCTION**

Trauma to the anterior teeth is a relatively common occurrence during childhood resulting in inflammation or necrosis of pulp and subsequent incomplete development of root apices.<sup>[1]</sup> Such teeth present with wide root canals and open apices causes the endodontic treatment technically more difficult and time consuming.<sup>[2]</sup> Due to the lack of apical barrier to contain the root filling material, obturation of the root canal with an open apex is difficult. The treatment of choice in such type of case is apexification, i.e. establishing an apical barrier.<sup>[3]</sup>

Apexification is any method responsible to induce a calcific apical barrier in a root

with an open apex or continued development of apex of an incomplete

root in teeth with non vital pulp". Traditionally, calcium hydroxide  $\text{Ca}(\text{OH})_2$  has been widely used for the apexification procedure due to its biological and healing performances. Calcium hydroxide combination with other materials has been proved to be the most commonly used material for apex closure. Apexification with calcium hydroxide involves repeated stimulations , for a period of atleast 6- 24 months, until apical closure is achieved.<sup>[4]</sup> Metapex, a silicone oil-based calcium hydroxide paste containing 38% iodoform is very popular (28) and has

been used as a root canal filling material in primary teeth. Use of it in apexification has also been reported. Iodoform and to the viscous and oily vehicle, which may prolong the action of the medicament. The present case report describes the successful non surgical management of immature permanent maxillary central incisor with open apex and large periapical lesion with 8 months follow up.

### **CASE REPORT**

A 13 year old female patient, reported to the Department of Pediatric and Preventive dentistry, K.D Dental college, Mathura, U.P, India with a chief complaint of swelling in association with her permanent maxillary right central incisor. On further examination it was found that there was no signs of caries or fracture but was sensitive to percussion and palpation. A radiographic image revealed an open apex and an associated large periapical lesion in relation to the right maxillary central incisor. The dental history disclosed that the patient has suffered dental trauma 2 years back with a recurrent swelling.

After administration of local anesthesia, a rubber dam was placed and the access cavity was prepared. Working length was determined with a no. 15 k file and root canal preparation was done till no. 80 k file (DENSPLY) using circumferential filing motion. The canal was irrigated copiously with 1.25% sodium hypochlorite (Vishal Dentocare Pvt Ltd, Hyderabad) and saline. The canal was dried with paper point, and triple antibiotic paste (metronidazole, minocycline, and ciprofloxacin in the ratio of 1:1:1) mixed with propylene glycol was placed as an intracanal

The superior antimicrobial effects of calcium hydroxide may be due to the combination with medicament and the access cavity was sealed with a temporary restoration.

At 14 days recall appointment, the intracanal medicament was replaced with metapex and patient was recalled at 2, 3, 5, 6 and 8 month intervals. The patient was clinically asymptomatic and radiographic evaluation showed reduction in periapical lesion and disclosed significant apical development of the tooth at 3 months interval. Follow up evaluation at 6 months from the time, revealed similar clinical findings and more healing of periapical lesion but the apex was not closed. After 8 months from the time of the induction apex was found closed radiographically.

### **DISCUSSION**

Apexification is the treatment of choice after confirmation of pulpal necrosis in immature permanent teeth. The procedure involves canal cleaning and shaping or the removal of all necrotic tissue, and hermetic sealing of the tooth by placement of suitable material to the apex to avoid bacterial infiltration. The prime importance of apexification procedure is to attain an apical stop for compaction of obturating material.<sup>[5]</sup> Various materials has been used for induction of apical barrier formation in nonvital permanent teeth such as calcium hydroxide paste, lime powder mixed with different vehicles, tricalcium phosphate, collagen phosphate, osteogenic protein-1, bone protein, oxidized cellulose, Mineral trioxide aggregate etc.<sup>[6]</sup>

Calcium hydroxide has been the first choice material for apexification because

of its antimicrobial action on bacterial cellular components mainly on lipopolysaccharide (LPS) (Jiang et al., 2003) and its high pH causes induction of hard tissue formation (Javelet et al., 1985). lime in contact with vital tissue in the apical area, seems to cause tissue reactions similar to those in the coronal pulp. Apexification procedure usually takes 6-24 months and refilling of root every 3-6 months is favoured.<sup>[7]</sup>

Metapex paste has been found to be an effective material for achieving apexification for younger permanent teeth. Metapex, releases lime slowly and tends to remain in the canal and exert its effects over a considerable period of time. In the present study, glass ionomer cement was used to restore the access cavity after placement of metapex in the root canal. This was done to supply an adequate coronal seal, and prevent any micro leakage.

A study by Ghosh et al., showed a 100% success with metapex as compared with calcium hydroxide mixed with water and non-setting calcium hydroxide with water.<sup>[8]</sup>

Similar finding was reported for apexification by Gu et al., where there was complete root development and apical closure.<sup>[9]</sup>

The present case indicate good results with metapex as an apexification agent taking into consideration the cost factors, simple placement and radiographic interpretation. This report shows both clinical and radiographic success in achieving apexification. Unlike with barrier formation, continued root growth was observed for the treated tooth.

## CONCLUSION

Metapex shows good clinical and

radiographic success in promoting continued root growth and inducing root end closure and also healing of periapical lesion in immature necrotic young permanent teeth.



Fig1.Pre-operative radiograph



Fig.2 2 months follow



Fig.3 3 months follow




Fig.4 5 months follow




Fig.5 6 month follow up




Fig.6 8 month

## REFERENCES

1. Anantharaj A, Praveen P, Venkataraghavan K, Prathibha RS, Sudhir R, Murali Krishnan B. Challenges in pulpal treatment of young permanent teeth a review. *J Dent Scien Res* 2011;2:142
2. Moore A, Howley MF, O'Connell AC. Treatment of open apex teeth using two types of white mineral trioxide aggregate after initial dressing with calcium hydroxide in children. *Dent Traumatol* 2011;27(3):166-173
3. Purra AR, Ahangar FA, Chadgal S, Farooq R. Mineral trioxide aggregate apexification: A novel approach. *J Conserv Dent* 2016;19:377-380
4. Sheehy EC, Roberts GJ. Use of calcium hydroxide for apical barrier formation and healing in non-vital immature permanent teeth: A review. *Br Dent J* 1997;183:241-246
5. Mohammadi Z, Dummer PMH. Properties and applications of calcium hydroxide in endodontics and dental traumatology. *International Endodontic Journal* 2011; 44:697- 730.
6. Lu YM, Qin JN. A comparison of the effect between vitapex paste and antibiotic paste in apexification. *Shanghai Kou Qiang Yi Xue* 2004; 13:449-51.
7. Jiang J, Zuo J, Chen SH, Holliday LS. Calcium hydroxide reduces lipopolysaccharidestimulated osteoclast formation. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* 2003; 95:348-54.
8. Weng QF. Clinical study on the effect of Vitapex paste in apexification. *Hua Xi Kou Qiang Yi Xue Za Zhi*. 2004;22(3):214-215.
9. Gu HX, Xu Q, Liu LM, OuyangY. Treatment of chronic apical periodontitis teeth complicated by open apices with Vitapex in the adults. *Shanghai Kou Qiang Yi Xue*. 2007;16(2):140-143.