

Original Research Article

Analysis of the hive yield of three species of bees for honey production in Eastern Honduras.

ABSTRACT

Objectives: The analysis of the performance of honey production hives of three species of honeybees was carried out.

Study design: three harvests per year of honey per hive of each species, with a sampling of ten hives per species, data were evaluated by ANOVA analysis, and Duncan mean separation test developed by the IBM SPSS 26.0 program.

Place and duration of the study: This research was developed in the department of Paraíso, municipality of Danlí, Honduras, Central America, in Cerro El Águila, latitude 13° 56' 59" N, and longitude 86° 13' 59" W, the sampling was carried out in 2020.

Methodology: the qualitative variables were evaluated for each species by *In Situ* observation that characterized the honeycombs of the three species, and the data were subjected to an analysis of variance and Duncan's mean separation, where the dependent variable is the average yield of Kg of honey per hive.

Results : according to the ANOVA we have that there are significant differences in the harvest of the three species of bees ($P = .05$), and in the analysis of separation of means the three averages of honey yield of the three species observed are not homogeneous ($P = .05$), the *Apis mellifera* produces 63.1 kg / hive-year, the *Tetragonisca angustula* obtained an average of 0.679 kg / hive-year, and *Melipona beecheii* obtained 13.19 kg/hive-year.

Conclusion: In the three species we observe that they have differences in the structures of the construction of their hives, their entrance, the construction of their cells, production and storage of honey and pollen, birth of queens, and method of defense; and when evaluating the average annual yield per hive of the three species we find that the three species of bees are different averages of honey production in kg / hive-year.

Keywords: Bees, *Apis mellifera*, hives, *Melipona beecheii*, honey, *Tetragonisca angustula*.

1. INTRODUCTION

Beekeeping could provide more diversified productions such as natural foods rich in vegetable protein, vitamins and minerals. Today they are lost or ceased to be produced due to ignorance or technological insufficiencies, especially in the fields of Latin America, and Africa, pollen, honey and royal jelly, products that could or reverse the hunger of these peoples (1). In Honduras the production of honey has experienced a gradual development in recent years, unfortunately it is mostly only for the local market. There are some 3 800 beekeepers, many of them constituted in companies dedicated to the production, processing, and marketing of honey, are grouped in 122 associations of producers, and all these in turn in the ANAPIH (2) . Contreras describes that it is important to analyze meliponiculture, not only as an economic activity that can provide resources to local populations, but as a biocultural heritage of relevance for the maintenance of the ecosystem (3). In addition, he mentions that the native human-bee relationship and its upbringing implies a network of knowledge and knowledge that are part of the worldview of the peoples.

The study aims to give an essential contribution to the country to characterize honey production in yield per hive-year, observing three varieties of bees, two native to Mesoamerica, as well as making their contributions regarding food security, the rural economy and human health, since information is limited, knowing that there are many producers of melipona bees in the country today, and that it has been an ancient activity recorded since the times of the heyday of the Mayan civilization; López cites that archaeological evidence suggests that it is in the region of

the Yucatan Peninsula where meliponiculture originated, approximately 1400-1900 years ago, and where its products, such as wax and honey, came to represent one of the pillars of the Mayan civilization. (4). By estimating the productions of these three species, and characterizing their hives, data that will be available to produce honey of different native species of bees of Honduras, technifying them at home, to improve efficiency in productivity, and to be able to give added value to beekeeping products, this item has special importance for the economy of the country, also helping the conservation of biodiversity and the general habitat.

1.1 Characteristics of Honey

Honey is a sweet, unfermented substance produced by bees from the nectar of flowers or secretions on or from living plants; that they collect, transform and combine with specific substances and that they finally store and mature in honeycombs. It is composed of water, fructose, and glucose, as well as acids, minerals, amino acids, proteins, enzymes, aromas, proteins, and other substances (5). Likewise, the content of hydroxymethylfurfural (HMF) should be values ≤ 40 mg/Kg and the diastatic activity not less than 8 units of diastase (DN) (6). Ureña cites that the most common components found in honey are water (17.1%), sugars (82.4%), proteins (0.1%) and other components that include vitamins, minerals, aromatic substances, and organic acids, among others (0.4%). In addition, five biologically active enzymes are found in it: invertase, diastase, glucooxidase, catalase and an acid phosphatase. (7)

In what represents the productivity of honey in our region, propitiatory dates and diversity of offerings in the Madrid Codex (Figure 1), which reflects that it is an ancient activity in the Mesoamerican region, the management of stingless bees, which were sacred to the Mayan civilization. This has been one of the most developed cultural areas in the breeding of native stingless bees (Hymenoptera: tribe Meliponini), also known as melipona bees; In the Yucatan Peninsula through the Mayan culture, he stood out in this activity called meliponiculture, whose antiquity can be traced back to the pre-Columbian period. (8)




Figure 1. Madrid Codex. Mayan Polymeniculture.
(9)

1.2 Description of the species studied

The best known and studied bee in our environment are the bees of the species: *Apis Mellifera* Africanized bee introduced in America to increase honey production. But not everyone knows about native bees such as the jimerito (*Tetragonisca angustula*) and the royal bee jicote (*Melipona beecheii*) (Figure 2), which produce an excellent quality of honey, produced entirely for medicine and supplementary food. The biological characteristics of the three varieties of bees studied are found: the size of the bee in the *Aphis mellifera* the size is usually 20 to 30 mm in length and robust, the *Tetragonisca angustula* is smaller measures 2 to 8 mm in length and thin and the *M. beecheii* is slightly larger can reach 8 to 25 mm in length and robust, as shown in Figure 2 (10)



Figure 2 *Tetragonisca angustula*(a), *Aphis mellifera*(b), *Melipona beecheii*(c)

1.2.1 Description of native bees or meliponas: jimerito (*Tetragonisca angustula*)

To systematically unify the name of this species we will give the scientific name that is *T. angustula*, its abdomen is yellowish brown, a thorax and dark head bright. This species of bee, We can find it throughout our country in heights ranging from 600 to 1 800 meters above sea level in different formations of wooded environments, and to the surprise of many can also be found in urban areas where they are established in buildings, tubes, cemeteries, and any space that you find wide for their nests (Figure 2). They form quite numerous populations with a large number of workers that are reported more than 5000 individuals depending on the space of the hive, they are very docile bees that can be easily managed by the whole family, since they do not have a stinger (11). Bees do not store honey in honeycombs, instead store the honey in amphorae (pots), which must be broken or crushed to extract the honey. Breeding honeycombs composed of wax have overlapping disc shapes separated from each other by earwax pillars, forming spaces that allow the circulation of bees (Figure 3). As for the physicochemical characteristics of the honey of this species we can highlight, which has a humidity of 30%; acidity 70 %; sugars 50%, pH of 3-3,5, degrees °Brix 79%, a clearly crystalline color and acidic flavors. . (12)

1.2.2 Native bee description: Royal Jicote or morito (*Melipona beecheii*)

It is another type of *Melipona* bee or native, but larger in the Mesoamerican region, known as real honey jicote bees, they are dark in color with light stripes on the abdomen almost the size of honey *Apis*. Looking inside these hives we can see a lot of similarity with the involucre of the breeding honeycombs and the honey pots of the jimeritos, the only difference is in the size. Its entrance to the nest is built with a mixture of mud and resin called batumen or geopropolio of circular shape where only one bee fits at a time and that will have a permanent guardian bee. Breeding honeycombs built with earwax are disc-shaped placed on several floors with space between the levels for bee circulation. Their honeys have different uses in home medicine, for bronchitis and stomach infections (13). The characteristics of the honey of this species are, humidity 30%, sugars 60%, pH 3.6, acidity 75%, soluble solids 74.6%, Degrees °Brix 81%, amber color, acid flavor, light cloudy appearance, and texture of moderate viscosity. (14)

1.2.3 Description of European or Africanized bees (*Aphis mellifera*)

This species as well as the native or *melipona* bees, they are considered as the main pollinators of some wild and cultivated plants (flowering plants or angiosperms). In this way, they help the conservation of ecosystems and improve the quality and quantity of agricultural products. Most bees are hairy-bodied and have a feathery appearance; they carry an electrostatic charge with which they attract small pollen particles. All of this helps the pollen stick to your body. With their legs they transfer pollen to pollen baskets that can be of two types: cups or corbiculas. The physicochemical characteristics of the honey produced by *A. mellifera* in terms of humidity has 25%, Degrees °Brix 77.9, total solids 83.45%, sugars 69.1%, acidity 40% with pH 3.8, light color and viscosity and flavors different sweet types. (15)



Figure 3 Types of bee hives used. *Melipona beecheii* in the wooden box (a), *Aphis mellifera* (b), *Tetragonisca angustula* hive in pumpkin (c).

2. MATERIAL AND METHODS

This research was developed in the department of Paraíso, municipality of Danlí, Honduras, Central America, in Cerro El Águila, latitude 13° 56' 59" N, and longitude 86° 13' 59" W, with an average height of 1 400 m.a.s.l., which is a mountainous area with a rich diversity of fauna and flora, the sampling was carried out in 2020, evaluating the productivity of three species of bees which are: *Apis mellifera* bees European or Africanized, bees native to the region *Tetragonisca angustula* better known as jimerito, and *Melipona beecheii* known as royal jicote or morito, from which three harvests were obtained for each species of bee, the first harvest and honey harvest was made in the months of March and April taking advantage of the flowering of coffee, the second in June and July with the flowering of citrus, oaks, oaks, and medicinal flowers; the third was held between November and December as native plants and wildflowers bloom.

2.1 Achievement of melipona bees in the field

To make the achievement of colonies of meliponas or native bees, they can be extracted directly from the trees of the forest, without forgetting that most of these species have the need to house trunks, coconuts, cavities that have left other animals, caverns, bricks of houses, different places that offer it a safe cavity and where they can build their nest, in our case, logs were used to house the hives of royal jicote, where we used two types of dry trees, the guamo (*Inga feuilleei*), and the oak *Quercus Robur*(Fig. 4-a). The colonies of Melipona bees will always form their colonies of access to the nest what we call entrance, this will be protruding from the trunk and there we will see the flow of bees at the entrance and exit, this entrance can be up, down or in the middle of the nest, here the person who is going to make the capture must be malicious with some experience and who knows very well what direction it may be towards the nest so as not to deteriorate it, the way in which we can have access to the nest, handling very carefully so as not to damage the nest when extracting it. Once we have the nest extracted, the largest number of bees and the queen in our trunk in the case of the royal jicote, we proceed to take out the reserves of honey and pollen, which at no time should go to the trunk, since this can bring pests to the hive, before obtaining the colonies you must have the site in the place where it will be placed or housed, It should be a quiet place, shaded and away from strong winds and bases protecting bees from pest attacks.

2.2 Method of obtaining native bees using traps

To obtain native bees we also use traps, the first thing we must do is clean the containers with water, then we will use an attractive solution that is composed of 70% alcohol, where wax and propolis of the same bees are dissolved so that it adheres inside the trap. With this substance we must impregnate the container well, drain the residues of this attractant well and put the containers to dry in the sun for 15 days to ensure that the aroma of alcohol escapes completely: this type of traps must be placed in very strategic places where the swarms of meliponas or native bees can possibly arrive, these were placed in trees, preferably where we know that there were nests of bees, in the case of *T. Angustula* bees. Pumpkin (*Lagenaria siceraria*) better known as jicaros were used in our country as shelters for all the hives used (Fig. 4-c), since it is a type of natural container, and this species works very well.

2.3 Installation of Apis mellifera hives

Ten hives were installed with two raises or frame boxes each, ten frames per box, these were located at two meters between hives and in the same area, and each hive had 20 frames, of which 60% were with children and pollen and 40% of honey, this in relation to the harvest time. From the hives observed, average records of the honey production of each hive were made, in three different harvests that were made in the year, which were extracted from the honeycombs and centrifuges, to then weigh the honey produced.

2.4 Fieldwork

Once the hives were installed through direct observation of the three species studied, 10 hives were evaluated for each species of bees placed randomly at different points, and the productive cycle of one year of each hive was observed, making three harvests per species, without altering the natural conditions of the area where all the hives were installed.

2.4.1 Characterization of hives by species:

The following qualitative variables are evaluated for each species:

Nature of the entrance of the nest, real cells, location of the real cell, honeycomb shape, method of harvesting pollen, rate of production of queens, and method of defense, which were determined by observation of the hives of each species, and the reality is analyzed in its natural context and as it happens, taking out and interpreting phenomena according to what was observed by the researchers who were in the field.

2.4.2 Honey production yield:

Quantitative analysis: the yield in kg of honey of annual production per hive of each species was determined, as well as the yield in weight in Kg of honey per harvest considering three annual harvests for each of the species.

To determine the harvest time, it depends a lot on the flowering season, considering also that if there is a lot of rain the bees do not work at their usual pace, so they were harvested in non-rainy weather, and in high flowering time, in the case of *Apis mellifera*, when the first 4 rises are full and the honey is ripe in the hive, it is time to make the collection of the honey, which was carried out in the evening, then the honey was transferred to the processing plant to centrifuge and strain it, to then be weighed; in the case of stingless bees, when the honey is ready, the bees usually shake outside the hive, and with the flowering period, we realize that the hive is ready to make the harvest, which was checked by direct review of the honeycombs, then we open the hive to extract the honey, by means of a syringe the honey was removed from each hive, with great care not to damage the wax of the honeycombs, always in the afternoon, and later this honey was weighed by each hive, in the case of the hive, these are washed with clean water in case honey is spilled and resealed with clay to avoid attacks of other insects.



Figure 4 Extraction of Honey from hives. *Melipona beecheii* in trunk, EPL (a), Honey harvest (b), *Tetragonisca Angustola* hive in pumpkin(c).

2.5 Statistical analysis.

Table I. Summary of characteristics of the hive

The samples were subjected to an analysis of variance, where the dependent variable is the average yield of Kg of honey per hive, evaluating as the main factor the three species of bees, and as a subfactor three harvests per year, evaluating in blocks ten that represent the hives by species. In addition, analysis was made of separation of Duncan's stockings by bee species, by annual harvest (first, second and third), and by each hive evaluated (block). The statistical program SPSS 26.0 was used to analyze the data by means of a univariate analysis.

3. RESULTS AND DISCUSSION

3.1 Characterization of hives by honeybee species.

Feature	<i>Aphis mellifera</i> European-Africanized	<i>Tetragonisca angustula</i> Jimerito	<i>Melipona beecheii</i> Royal Jicote
Nest entry nature	They do not perform any specialty at the entrance of the nest, it is directly.	Usually made of earwax.	Usually made of mud. Nest entrances frequently Radial.
Real cells	Large specialized real cells.	Real specialized cells larger than workers and males	No specialized real cells
Location of the actual cell.	The cells are located near the brood chamber and pollen.	Near the periphery of the honeycomb. It does not include those that make cells in clustered form.	Interspersed in the honeycomb with cells of workers and males.
Honeycomb deposit form.	Hexagonal	Spherical, honey pots.	Spherical, honey pots.
Harvest pollen.	Trap is needed for collection	This accumulates in the spherical pots.	This accumulates in the spherical pots.
Queen production rate	Frequent	Relatively rare	Frequent
Method of defense	It has a stinger with poison that harms humans and animals.	It does not harm any harm to a person or has a specific method of defense. No sting.	It has sticky substances, but it does not harm people. No sting.

From the observation of the hives of the three species we denote that *A. mellifera* does not make any entry into the nest, in exchange the *T. angustola* and the *M. beecheii* elaborates an entrance of wax to the nest, with respect to the real cells, the *A. mellifera* and the *T. angustola* are large, and the *M. beecheii* does not have specialized cells; as far as the location of the real cell is concerned, in *Apis M.* they place them near the brood chamber and pollen, in the *T. angustola* it is near the periphery of the honeycomb, and in the *M. beecheii* they are intermingled in the honeycomb with the cells of workers and males; in the case of honey deposits to *Apis M.* the shape of the honeycomb is hexagonal, in the case of the other two species they store honey in honeycomb in a spherical shape, called pots; propolis in the case of *Apis M.* needs traps to collect it, in the case of stingless bees, they store it in spherical pots; In the reproduction of queens it is denoted that in *A. mellifera* and *M. beecheii* are frequent, contrary to *T. angustola* are very rare new queens; and the method of defense only *A. mellifera* has a sting that has a poison that can harm humans and animals, while the other two species jimerito and jicote morito have no defense system, which makes them more vulnerable to attack by other insects, that is why they are better known as melipona bees (Table I); this characterization of the hives is in accordance with what is described in the species (16) *T. angustola* and *M. beecheii*, which have been domesticated and adapted for a long time in Mesoamerica.

3.2 Statistical analysis on yields per hive and per harvest.

Table II. Factorial arrangement for ANOVA of honeybee yield. This table presents the honey yields in Kg. per hive of each of the three species, and the three annual harvests that were made, where data were obtained from ten hives per species for analysis.

FACTOR	HARVEST	Blocks									
		I	II	III	IV	V	VI	VII	VIII	IX	X
<i>Tetragonisca angustola</i>	FIRST	0.40	0.52	0.30	0.45	0.46	0.48	0.49	1.00	1.20	1.00
	SECOND	0.35	0.45	0.22	0.25	0.26	0.35	0.46	0.59	1.00	0.90
	THIRD	1.00	1.00	0.80	0.85	0.70	0.40	0.89	0.90	1.50	1.20
<i>Melipona beecheii</i>	FIRST	2.20	2.90	2.30	3.00	4.00	4.60	4.90	3.00	4.00	5.00
	SECOND	3.00	4.20	4.00	3.60	2.00	2.50	3.00	4.00	5.00	10.00
	THIRD	5.00	3.00	3.60	4.60	6.00	5.20	5.80	6.00	4.50	11.00
<i>Apis Mellifera</i>	FIRST	19.00	21.00	21.00	19.00	17.00	25.00	19.00	21.00	23.00	21.00
	SECOND	25.0	23.00	21.00	25.00	32.00	32.00	21.00	32.00	25.00	17.00
	THIRD	19.00	21.00	17.00	15.00	13.00	21.00	15.00	17.00	19.00	15.00

Table III. Analysis of Variance. Applying the univariate ANOVA analysis in SPSS 26.0, we have that there are significant differences in the average honey harvest in kilograms of the three bee species studied ($P = .05$), and there are no significant differences between the average yield of the hives in kilograms of honey ($P = .05$).

Dependent variable: Yield Kg.					
	Tipo III Suma de cuadrado s	Df	Cuadrado medio	F	Sig.
Fountain					

Corrected model	7175.56 ^a	13	551.97	61.29	0.00
Intercept	6816.80	1	6816.80	756.98	0.00
Bee species	7048.99	2	3524.49	391.38	0.00
Harvest	74.88	2	37.43	4.16	0.02
Block(hives)	51.71	9	5.75	0.64	0.76
Error	684.40	76	9.01		
Total	14676.75	90			
	5.0				
Total, corrected	7859.96	89			

a. R Square = 0.913 (R squared = 0.898)

Table IV. Test of Separation of stockings by species. According to Duncan's mean separation analysis, we denote that the three average honey yields in kg/hive of the three observed bee species are not homogeneous at a significance level of (P =.05)

Bee Species	N	Subset		
		1	2	3
<i>Tetragunista angustula</i>	30	0.68		
<i>Melipona beecheii</i>	30		4.40	
<i>Apis mellifera</i>	30			21.03
Sig.		1.00	1.00	1.00

Table V. Analysis of separation of means per harvest of each species of bee.

Duncan ^{a,b}						
	Segunda <i>Tetragunista angustula</i>	10	0.48			
	Primera <i>Tetragunista angustula</i>	10	0.63			
	Tercera <i>Tetragunista angustula</i>	10	0.92			
	Primera <i>Melipona beecheii</i>	10		3.59		
	Segunda <i>Melipona beecheii</i>	10		4.13		
	Tercera <i>Melipona beecheii</i>	10		5.47		
	Tercera <i>Apis mellifera</i>	10			17.20	
	Primera <i>Apis mellifera</i>	10				20.60
	Segunda <i>Apis mellifera</i>	10				25.30
	Sig.		0.70	0.10	1.00	1.00

When evaluating the harvest by species of bee by Duncan mean separation test we have that the average honey production averages in kilograms of the three species are different, for *T. angustula* in the analysis of its three annual harvests by weight they are homogeneous (P =.05), that is, the honey yield of the average harvest in the three seasons in a year are homogeneous, it also happens with the species *M. beecheii*, its average yield is homogeneous in the three annual harvests, only in the *A. mellifera* the yield in kg. of honey of the three harvests by weight are not homogeneous (P =.05), that the three averages in kilograms of honey obtained in the hives are different.




Figure 5. Average harvest per species.

In this graph we denote the averages of honey in Kg. for each of the three species of bees and the three annual harvests per species, where the native bees evaluated tend to raise the average gradually in the three harvests, not so with the *A. mellifera*, which showed that in the second harvest the highest average was obtained, and the last harvest had the lowest yield.

In the case of the bees *T. angustula* Nates-Parra describes that it can be obtained from 500ml to 1 liter / year (0.704 – 1.408 Kg / year.) , in our samples the average values obtained are 2,037 kg honey / hive-year, slightly higher than the average of Nates; as for (17)the *M. beecheii*, he comments in his study carried out in Cuba that the annual production per hive is 10.5 kg / year, being able to reach 13.5 kg/year; when compared with our averages, we denote the closeness of the averages, our average is 13.19 kg honey/ hive-year, which remains in this close range of productive yield. (18)

The bee *A. mellifera* as (19) found in the Altiplano Ecuador a yield of 25.08 Kg / harvest-hive, and in our case we obtained averages of 20.9 kg honey / hive / harvest, and 63.1 Kg of honey / hive-year; in the same way in the state of Veracruz, Mexico production data is presented where it specifies that: the technical level produces between 60 to 70 kg, semi-technified between 30 to 45 kg, and non-technified between 25 to 30 kg of honey per hive per year (20), performance that is similar with our averages of Kg. honey/hive are within this range of a hive of technical level of this work, which states that in both research works cited in different places they are within the similar range of our yields per annual hive in this species.

In what corresponds to the medicinal value of honey, this food is of utmost importance, for its infinity of properties that it has, it already offers new opportunities for rural income for the added value it can generate, García suggests that for the use of honey as a nutraceutical the content of phenolic acids is considered, flavonoids, ascorbic acid, proteins, carotenoids, and certain enzymes such as glucose oxidase and catalase. Its antimicrobial action against *Helicobacter pylori* is well documented. In addition to the formation of peroxide, glucose oxidase acts on fibroblasts and activators of epithelial cells. It has bactericidal properties against pathogenic and enteropathogenic bacteria (21). In relation to cosmetic applications, it has great nourishing and antiseptic effects on the skin, which is why it is used in a variety of beauty products that are found especially in beauty centers. (22) comments that for all this honey represents a range of opportunities for the development of new products, mainly native species. Also the entry of cheap honeys into the country in the coming years will be

an enormous challenge for beekeepers, there is food insecurity, since only a small number of families have the opportunity to fully cover all the links of what food security entails; it is necessary to focus on those areas, where families are in food uncertainty, (rural areas) working families, who struggle day by day, but that the socioeconomic conditions do not favor them, since although their income improves, the prices of basic basket, public services, and transport; they increase, they do not have access to quality education, that makes illiteracy levels continue to increase, therefore there is no improvement in technologies applied to the countryside, to their coffee farms, to their small business (23), for the above the production of honey becomes an alternative to improve family income, improving access to and availability of food, generating income through the production and processing of beekeeping products; Bees are part of the biodiversity on which we all depend to survive, they provide high-quality foods.

The growing agriculture is a threat to native species, given that in many cases it contributes to the destruction of the natural habitat of these species, Martínez describes that the transformation of the natural environment to move to the agricultural exploitation of the soil has caused a very negative effect on the bee wildlife: on the one hand, the most favorable places for nesting have been destroyed, and on the other, the plant species that provided the original food source have been eliminated. (24)

According (25), to Climate Change affects beekeeping mainly indirectly, however, a direct impact lies in the alteration of the behavior and physiology of bees, they adapt to cope with environmental conditions; so it is important to take mitigation measures to reduce the impact of Climate Change on Beekeeping, however, they must be aimed at a common good, since the survival of man affects the different ecological systems, in such a way that the actions to be taken to face Climate Change.

4. CONCLUSION

Regarding the hives of the three species was observed that they have differences in the structures of the construction of their hives, their entrance, the construction of their cells, production, and storage of honey and pollen, birth of queens, and method of defense. When evaluating the average annual yield per hive of the three species we find that *Apis mellifera*, *Tetragonisca angustula* and *Melipona beecheii* are different averages of honey yields in kg/hive-year. Honey production in our country offers an income opportunity for rural areas, which contributes to the food security of rural populations, in addition to improving native ecosystems, and hives offer high quality food with nutraceutical properties.

REFERENCES

1. *Beekeeping and food security*. **Green, Mayda M.** [ed.] Instituto de Ciencia Animal. 1, 2014, Revista Cubana de Ciencia Agrícola, Vol. 48, pp. 25-31. 0034-7485.
2. *Physicochemical determinants of honey quality: a literature review*. **Pineda Ballesteros, E., Castellanos Riveros, A., and Téllez Acuña, F. R.** Colombia : s.n., 02 02 2019, Cuadernos de Desarrollo Rural.
3. **Contreras Cortés, Leonardo Ernesto Ulises, Vázquez García, Amparo, Aldasoro Maya, Elda Miriam, & Mérida Rivas, Jorge.** *Knowledge of native stingless bees and generational change among the Lacandon Maya of Nahá*. Chiapas : s.n., 2020. pp. 205-225. Vol. 56.

4. *Decoloniality as an alternative for the conservation of biodiversity. The case of meliponiculture in the Yucatan Peninsula.* **López Barreto, Mauricio Feliciano.** 1, 2021, *Peninsula*, Vol. 16, pp. 29-53. 1870-5766.
5. **Venancio Martínez,** . usc.gal. *University of Santiago de Compostela.* [Online] 2010. [Cited: 20 04 2022.]
https://www.usc.gal/export9/sites/webinstitucional/gl/investigacion/grupos/malaterra/publicaciones/IV_Ciclo/Tema_13_Benancio_Trabajo_de_la_miel.pdf.
6. *Chemical composition of honey and its relationship with health benefits.* **Chaviano, María Elena García, et al.** 1, Cuba : s.n., 01 01 2022, *Revista Médica Electrónica*, Vol. 44.
7. *Evaluation of the possible adulteration of commercial bee honeys of Costa Rican origin when compared with artisanal honeys from specific apiaries.* **Ureña Varela, Maurico, Arrieta Bolaños, Esteban, Umaña, Eduardo, Zamora, Luis Gabriel, & Arias Echandi, María Laura.** 57, 2007, *Latin American Archives of Nutrition*, Vol. 1, pp. 63-69.
8. *Bees of the Mayab, from the MAYAN cosmogony TO AN ETHNOBIOLOGICAL COLLECTION OF THE ROGER ORELLANA REGIONAL BOTANICAL GARDEN.* **Canto, , et al.** [ed.] A.C Centro de Investigación Científica de Yucatán. s.l. : Centro de Investigación Científica de Yucatán, A.C, 2021. 2395-8790.
9. **Foundation for the advances of Mesoamerican Studies.** FAMSÍ. [Online] 05 26 2022. http://www.famsi.org/mayawriting/codices/pdf/4_madrid_rosny_bb_pp79-112.pdf.
10. *Colonization, impact and control of Africanized honey bees in Mexico.* **Guzman, , et al.** 2, Mexico : s.n., 06 2011, *SciELO Analytics*, Vol. 42.
11. **Montenegro, Lanza, Johel David.** *repositorio.una.edu.n.* Faculty of Rural Development, National Agrarian University. Managua : s.n., 2013. Thesis.
12. *Quality standards for medicinal uses of Meliponinaemiel in Guatemala, Mexico and Venezuela.* **Vit, , Medina, and Enríquez, María Eunice.** 2, Mexico : s.n., 01 04 2004, *Bee World*, Vol. 85, p. 4. 5.
13. *Breeding Stingless Bees.* **Sandker, and Totaro,** . 03 2019, *Promabos*, p. 1.
14. *Physico-chemical and organoleptic characterization of Melipona beecheii honey obtained in agroforestry systems.* **Fonte, , et al.** 03, Habana Cuba : s.n., 09 2013, *Pastos y Forrajes*, Vol. 36, p. 3.
15. *IDENTIFICATION OF FLORA AND NUTRITIONAL ANALYSIS OF HONEY FOR BEEKEEPING PRODUCTION.* **Insuasty, , Martínez, and Jurado,** . 01, 06 2016, *Biotechnology in the Agricultural and Agroindustrial Sector*, Vol. 14, pp. 37-42.
16. **Arnold, Noemi: Zepeda, Raquel: Vásquez Dávila, Marcos: Aldasoro Maya, Mirian.** *The stingless bees and their cultivation in Oaxaca, Mexico, with catalog of species.* First Edition. San Cristobal de las Casas. : El Colegio de la Frontera Sur., 2018. pp. 33-47. 978-607-8429-53-0.
17. *Meliponiculture: an activity that generates income and environmental services.* **Nates Parra, Juan Manuel and Rosso Londoño,** . [ed.] Bee Research Laboratory. Bogota : s.n., 2005, *LEISA Revista de agroecología*.
18. *Melipona beecheii Bennett (Hymenoptera: Apidae): origin, studies and meliponiculture in Cuba.* **Genaro, Julio A: Loringa, Walberto.** [ed.] University of Nebraska. 1-18, Gainesville, FL : s.n., 2018, *Insecta Mundi*, Vol. 0643. 0749-6737.
19. *Relationships between honey production, defensiveness and diameter of breeding cells of Apis mellifera L; in the Ecuadorian highlands.* **Masaquiza-Moposita, Diego Armando, Curbelo Rodriguez, Lino Miguel and Diaz Monroy, Byron Y Arenal Cruz, Amílcar.** 3, Camagüey : s.n., 2019, *Revista de Producción Animal*, Vol. 31. 2224-7920.
20. *Beekeeping situation in Mexico and prospect of honey production in the State of Veracruz.* **Muciño, Luis Enrique Soto, Elizarraras, and Muciño, Ivonne Soto.** 07, Veracruz Mexico : s.n., 03 2017, *Revista de Estrategias del Desarrollo Empresarial*, Vol. 03, pp. 50-51.
21. *Chemical composition of honey and its relationship with health benefits.* **García-Chaviano, María Elena, Armenteros-Rodríguez, Esther, Escobar-Álvarez, María del**

- Carmen, García-Chaviano, Jorge Andrés, Méndez-Martínez, Jesús, & Ramos-Castro, Guillermo.** 1, Matanzas : s.n., 2022, Revista Médica Electrónica, Vol. 44. 1684-1824.
22. *History and current affairs of skin products, cosmetics and fragrances.* **González Minero, Francisco José, & Bravo Díaz, Luis.** 5-12, 2017, Especially Plant Derivatives, Vol. 58. 23409894.
23. **Torres Mejía, Juan & Torres Mejía, Fredy & Lopez Toro, Claudia.** *Analysis of food and nutrition security, The covid 19, western region of Honduras.* Santa Rosa de Copan : NEXOS, 2022. Vols. Vol. 35, No. 01, pp. 41-59/March 2022.
24. *Importance of bee diversity (Hymenoptera: Apoidea) and threats it faces in the tropical ecosystem of Yucatan, Mexico.* **Martínez-Puc, Jesús Froylán, & Merlo-Maydana, Flavio Eudaldo.** 2, 2014, Journal of the Selva Andina Animal Science, Vol. 1, pp. 28-34.
25. *Impact of Climate Change on Beekeeping.* **Medellin.** 3, 2012, Beekeeping National Atlas of bees, pp. 4, 5, 6.

UNDER PEER REVIEW