

A REVIEW ON THE EFFECT OF MICRONUTRIENTS IN HEAT STRESS ALLEVIATION IN DAIRY ANIMALS

ABSTRACT

Heat stress from global warming and climate change, which result in an ongoing rise in Earth's temperature, has a detrimental effect on the growth and health of dairy animals. The animal experiences heat stress when it cannot release enough heat to maintain homeothermy. The degree of heat stress dairy animals experience depends on the ambient temperature and humidity. In dairy animals, heat stress reduces feed intake, milk production, reproductive efficiency, and immune function. This article will concentrate on the micronutrients that lessen the damaging effects of heat stress on dairy animals. Micronutrients are vital substances that life requires in minute amounts. Major minerals, trace minerals, and vitamins are all included. Micronutrients support the maintenance of animal production, enhance nutrient utilization, effectively combat oxidative stress, and strengthen the weak immune system. Minerals are crucial for maintaining an animal's normal physiological processes. However, it's believed that animals' reactions to heat stress increase mineral loss through excretion. Therefore, adding minerals to the diet (such as DCAD, Zinc, Chromium, Selenium, etc.) may help to reduce the harmful effects of heat stress. Vitamins serve as cofactors for enzymes, act as catalysts in a number of metabolic pathways, and are crucial for an animal's normal growth and development. The addition of vitamins (such as Vitamin E, Niacin, etc.) to dairy animals' diets may also help to mitigate the harmful effects of heat stress.

INTRODUCTION

In the present scenario, heat stress is a major concern in rearing dairy cattle. Heat stress occurs when an animal's heat load is greater than its capacity to lose heat. Dairy animals feel hot sooner than humans do. Factors like high air temperature, humidity, solar

radiation and low air movement contribute to increased risk. A living body can only maintain its core temperature when heat production and heat loss are balanced. The thermoneutral zone is characterized as the range of surrounding temperatures where the body can keep up its core temperature exclusively by regulating dry heat loss, i.e., skin blood flow (Kingmaet al., 2014; Singh et al., 2020a; Singh, 2021).

Temperature humidity index (THI) is a simple method to measure and assess heat stress. It depends on humidity and ambient temperature. Present-day high-producing cattle begin to experience heat stress at a THI of 65-68. As a broad classification, the level of stress can be separated into light (68-71), moderate (72-79), severe (80- 89), very severe (>90), and deadly (>100). Signs of heat stress include reduced rumination, increased CO₂ output, increased respiration and panting, reduced feed intake, decrease in growth, reduced milk production, reduced carcass quality, decreased fertility, increased sweating, and finally, these all lead to an increase in mortality rate (Bin-Jumah et al., 2019). Older dairy animals seem to be more severely affected compared to younger animals. Heat stress reduces dry matter intake and leads to as much as a 30% drop in milk yield in dairy animals.

Strategies for the alleviation of heat stress include physical modification of the environment, genetic development of less sensitive breeds, and improved nutritional management schemes using dietary water, protein, fat, fibre, micronutrients, and feed additives (Conte et al., 2018) (Fig.1). Dietary yeast, chemical additives, fermentates, betaine, dietary cation-anion difference, propionate supplementation, and micronutrients are the main components of anti-heat stress additives. These substances are essential for improving the metabolic status of cattle as well as their immune system and energy metabolism (Zou et al., 2022). Micronutrients are essential elements needed by life in small quantities. It includes major minerals, micro/trace minerals and vitamins. Micronutrients help to maintain the production of the animals, improve nutrient usage, effectively neutralize oxidant stress and strengthen the compromised immune system (Mir et al., 2018). Based on different relevant studies list of micronutrients that are important for heat stress alleviation is given in Fig. 2.

During heat stress, there is an extra need for micronutrients. This is because the reduced feed intake and heat stress increase mineral excretion, whereas it decreases serum and liver concentrations of vitamins and minerals. Moreover, the excretion of vitamins and minerals and their mobilization from tissues are increased under stress conditions (Siegel., 1995). Recent research has demonstrated that combining micronutrients (Zn, Se, Vitamin E,

and Vitamin C) has improved animals' resistance to the harmful effects of HS. Animal performance under HS may be improved by the substances' capacity to prevent cell damage and enhance intestinal integrity and renal function (Ortega et al., 2022). The metabolic and physiological processes of several species of animals can also be improved by adding vitamins and minerals (Singh, 2019; Singh et al., 2020e). This may have an impact on the mineral digestibility and electrolyte balance of HS-affected animals. (Sivakumar et al., 2010; Liu et al 2018). Stress may aggravate a marginal vitamin and mineral deficiency or lead to increased vitamin and mineral requirements (Sahin et al., 2005). National Research Council recommendation says that mild to severe heat stress in dairy cattle has been estimated to cause an increase in maintenance requirements of micronutrients by 7 to 25%.




Fig.1 Improved nutritional management schemes during heat stress in animals




Fig. 2 Classification of important micronutrients helps in heat stress alleviation

MACROMINERALS

Heat stress-induced disturbances of acid-base status are related significantly to macromineral elements. The main change occurring during heat stress is an increase in the respiratory rate which causes carbon dioxide expulsion. That will lead to altered blood carbonic acid to bicarbonate ratio and the pH level increases. This condition is known as respiratory alkalosis. To maintain the pH, the bicarbonate will be expelled through urine. Rumen needs bicarbonate for buffering acids during rumen fermentation. Since the bicarbonate are excreted through urine, the rumen will not get enough bicarbonate ion and this ultimately leads to subacute ruminal acidosis (Sanchez et al., 1994). The demand for cations by the kidney is increased because the excretion of bicarbonate ions must be accompanied by the excretion of the cation. Potassium or sodium is a possibility however sodium is more likely to be excreted (El-Nouty et al., 1980). Sodium and potassium help to increase dry matter intake, increase in milk yield, increase in milk fat percentage and dietary Na and K are very important in the maintenance of water balance and acid-base physiology of heat-stressed dairy animals (Mallonee et al., 1985).

The increased amount of dietary chlorine in feed is much more detrimental to dry matter intake during summer. A study shows that ingestion of CaCl_2 increases plasma

chlorine concentration, which exceeds the bicarbonate buffering power capacity and produces metabolic acidosis, thereby appearing to suppress appetite in pigs (Yen et al., 1981). When animals, particularly cattle, are given diets with negative cation to anion balance (i.e. more anions than cations) or calcium chloride causes an acidifying effect, it is known as hyperchloremic metabolic acidosis.

Dietary cation-anion difference (DCAD), has been defined as milliequivalents of $(\text{Na} + \text{K}) - (\text{Cl} + \text{S})$ per kilogram of dry matter and has a direct impact on blood acid-base metabolism (Block et al., 1994). Dietary cation-anion difference calculated using Na^+ , K^+ , and Cl^- concentrations, has a significant effect on health status and productivity by influencing acid-base balance (Hu et al., 2004). Serum total amino acid and essential amino acid concentrations and the ratio of essential amino acid to total amino acid were greater for high DCAD. These results suggest that increasing DCAD improves amino acid availability for protein synthesis, which would otherwise be taken for the maintenance of acid-base balance (Wildman et al., 2007). A positive Dietary cation-anion difference diet of +350 mEq/kg dry matter improved the immunity status and nutrient intake by ameliorating heat stress (Suman et al., 2018). Keeping the dietary cation-anion difference at a healthy lactating level remains a good strategy for reducing thermal stress during the warm summer months.

Serum calcium is lowered in heat-stressed lactating dairy animals. This may be due to respiratory alkalosis. During respiratory alkalosis, the relative amount of bicarbonate ions in the blood will be increasing, along with which pH also increases. To neutralize it, the blood proteins release H^+ ions and they react with bicarbonate ions to form water and CO_2 . The free form of calcium present in the blood will get attracted to these negatively charged blood proteins and form calcium-bound proteins. Hence free calcium ions in the blood decrease during heat stress. So, there is a need for extra allowance of calcium for animals under heat stress (Odom, 1989; Sanchez et al., 1994).

MICRO/TRACE MINERALS

Supplementation of micro/trace minerals in heat-stressed dairy animals helps in improved performance and productivity, maintenance of proper immunity, udder health and reproduction, and reducing oxidative damage of tissues.

Zinc

Zinc is an important micromineral, involved in productive performances like growth (Underwood et al, 1999), immune system, and reproduction (Patel et al., 2017; Abdel et al., 2011), and involved in a wide range of metabolic activities. By using transcellular transport mechanisms, Zn is absorbed by the small intestine and activates antioxidant peptides and enzymes by promoting the expression of metallothionein, which is essential for shielding cells from ROS (Jarosz et al., 2017; Maares et al., 2020). Zinc plays a critical role in antioxidant defense as an integral part of superoxide dismutase (SOD), which is an essential enzyme. The deficiency of zinc has been reported to cause an increase in oxidative DNA damage and impair antioxidant functions (Underwood., 1999; Jing et al., 2007). Zinc supplementation lessens the heat shock protein response and enhances immunity in heat-stressed peripheral blood mononuclear cells of periparturient dairy cows (Sheikh et al., 2017). Inorganic salts like zinc sulfate ($ZnSO_4$), zinc oxide (ZnO), and zinc chloride ($ZnCl_2$) are the major sources of zinc in mineral supplements formulated for animal feeding (Batal et al., 2001). Mammary integrity should be maintained in high-producing cattle during heat stress to prevent a decrease in milk production. Studies indicate that dietary organic Zn complex (40 ppm) can improve mammary epithelial integrity, and the mammary cells displayed more integrity (Weng et al., 2018). High-yielding HF cows received a base diet containing 0.131 percent of diet DM of the zinc-methionine (Zn-Met) complex, which improved their oxidative status as evidenced by higher total antioxidant status and lowers malondialdehyde concentrations. Overall, the findings of this study demonstrated that feeding RPZM to animals during times of heat stress could maintain their ability to produce milk and the composition of their milk. Animals' improved performance after taking Zn-Met complex supplements may have improved oxidative and immune status (Kargar et al., 2022). Supplementing multiparous lactating Holstein with 35 mg of Zn/kg of DM from Zn hydroxy chloride and 40 mg of Zn/kg of DM from Zn-Met complex had a significant impact on mammary cell turnover (Rivas et al., 2021). These findings imply that cows receiving Zn supplements have increased milk secretion capacity and the ability to maintain milk yield under heat stress by safeguarding mammary epithelial cells and their capacity for secretion.

Chromium

Chromium is another favourable micronutrient for defying the adverse effects of heat stress in animals. It is a necessary mineral and is crucial for the metabolism of glucose as glucose tolerance factor (Lashkari et al., 2018). It acts as an excellent antioxidant that prevents heat-stress-induced lipid peroxidation. Chromium improves cortisol hormone

activity and nutrient metabolism. It promotes insulin action in responsive tissues, thereby increasing farm animal productivity (Bin-Jumah et al., 2020). Appropriate amounts of chromium-containing additives should be added to the ration for dairy cows during hot summers due to the low concentration of chromium in the ruminant feed ingredients. (Shan et al., 2020). 10 mg/day Cr given to dairy animals from 21 days prepartum to 35 days postpartum showed an increase in milk production which was associated with decreased lipolysis and increased glucose uptake (McNamara et al., 2005). Later studies also demonstrated that oral administration of various Cr concentrations increased dairy cows' DMI while not affecting milk yield (Vargas-Rodriguez et al., 2014; Leiva et al. 2017). Cr-methionine at 0.05 mg /kg increases feed efficiency and dry matter intake in cows (Arthington et al., 1997) and significantly reduced the respiration rate in dairy calves (Mousavi et al., 2019). A study showed that supplementation of inorganic Cr to the diet of buffalo calves reared under high ambient temperature improved heat tolerance, and immune status without affecting nutrient intake, and growth performance (Kumar et al., 2015). The estimated requirement of chromium of buffalo calves in summer conditions was calculated to be 0.044 mg/kg body mass and 10.37 ppm per day and the supplementation of Cr had a positive effect on its balance and plasma concentration without interacting with other trace minerals (Kumar et al., 2013). By lowering respiration rates and rectal temperatures, increasing dry matter intake and milk lactose content, and boosting antioxidant and immune function, chromium yeast (CY) supplementation at 0.36 mg Cr/kg DM improved the welfare of mid-lactation dairy cows (Shan et al., 2020). Additionally, it is said that Cr supplementation can guard against stress-related losses of Zn, Fe, Mn, etc (Sirirat et al., 2012).

Selenium

Selenium is an essential trace mineral that is an indispensable component of the antioxidant system (Oltamari et al., 2014). It has been demonstrated that dietary Se supplementation under heat stress supports reproductive physiology, production performance, and gastrointestinal health. The underlying mechanisms involve antioxidant status, immunocompetence, and regulation of nutrient digestibility influenced by gastrointestinal microorganisms (Zheng et al., 2021). Se, which is absorbed by active transport via a sodium pump in the duodenum and cecum, functions as a dietary antioxidant by forming selenoproteins and controlling the activity of endogenous enzymes (Mehdi et al., 2013; Kielczykowska et al., 2018). Selenium decreases the adverse effects of heat stress on metabolism and redox balance, resulting in improved dairy animal health, immune function,

and milk quality (Sejian et al., 2012). Studies show that in heat-stressed animals, there is a significant reduction in plasma selenoprotein, diet supplementation with selenium can significantly raise plasma selenoprotein and selenium concentrations, which might be a potential mechanism to protect dairy animals from heat stress (Hill et al., 2012). Se controls the production of selenoproteins, which take part in several cellular defense responses and shield cells from stressors like heat shock and oxidative damage (Kumar et al., 2017). Supplementing Se can significantly lower cellular HSP production, reduce the need for HSP defense against high-temperature stress, and reduce oxidative stress. (Zou et al., 2019). The supplementation of organic selenium in the diet had a positive effect on the percentage of milk fat and it improved mammary gland health (Oltamari et al., 2014). Sun et al. (186) found that Holstein dairy cows kept stable under heat stress with the addition of 0.3 mg/kg DM organic Se to their diets in the middle of lactation, while the GPx activity in the serum of the cows in a similar inorganic Se group gradually decreased. Studies have shown that selenium supplementation can stop heat stress-induced apoptosis in cells (Tang et al., 2019).

VITAMINS

Vitamins function as enzyme cofactors (coenzymes), participate in a variety of metabolic pathways as catalysts and are essential for the normal growth and development of animals. The addition of vitamin supplements to the diet of dairy animals might also contribute to the relief of the negative effects of heat stress (Min et al., 2019).

Vitamin E

Vitamin E is an antioxidant that is essential for body functions like growth, immunity, tissue integrity, and reproduction (Dalle et al., 2011; Singh et al., 2020b) and to prevent oxidative stress. In membrane and plasma lipoproteins, vitamin E functions as a free radical-suppressing antioxidant (Reboul et al., 2017; Singh et al., 2020c). Elevated temperature and humidity in summer will lead to greater oxidative stress for animals, so feeding of additional vitamin E is required. Vitamin E includes a group of tocopherols and tocotrienols, among them, α -tocopherol has greater biological activity. Due to the potent antioxidant properties of tocopherols, the impact of α -tocopherol in the prevention of chronic diseases believed to be associated with oxidative stress has often been studied and beneficial effects have been observed (BrigeliusFlohé et al., 1999; Singh, 2019; Singh et al., 2020d; Singh et al., 2020e).supplementation of vitamin E along with selenium (500 IU Vitamin E + 0.3mg selenium per kg DM) alleviated the heat stress in dairy cows. Supplementation of chromium

along with vitamin E and selenium (chromium propionate (0.5 mg) + vitamin E (500 IU) + selenium (0.3 mg)) in calves during heat stress decreased significantly ($P < 0.05$) the cortisol levels and rectal temperature (Sultana et al., 2022).

Vitamin C

According to Padilla et al. (2006), heat stress reduces the plasma vitamin C concentration in lactating cows; therefore, vitamin C supplements are required during the HS period. Reactive oxygen species (ROS) can damage cell membranes, DNA, proteins, and lipids during OS. Vitamin C is also crucial for the regeneration of other antioxidants, such as glutathione and alpha-tocopherol (vitamin E) (Padayatty et al., 2016). Dairy cattle are more susceptible to oxidative stress after calving and when under heat stress, and total ascorbic acid in plasma significantly drops during the peripartum period (Tanaka et al., 2011). According to experimental data with cattle, dairy cows need to have access to vitamin C to successfully adapt to stressful situations when there is heat stress (Ranjan et al., 2012). By lowering IL-6 levels and reducing ROS production, polyherbal vitamin C (20 g/d) can reduce inflammation. It also enhanced HF cow performance under heat stress.

Niacin

Niacin has been reported to be a vitamin that resists heat stress in cattle by increasing evaporative heat loss *in vivo* and cellular heat shock response by increasing gene expression of heat shock proteins during thermal stress *in vitro* (Zimbelman et al., 2013). Niacin (vitamin B3) supplementation increases resistance to heat stress by inducing greater cutaneous vasodilatation and blood flow and the increase in the rate of blood flow to the skin after niacin supplementation is associated with an increase in the evaporative heat loss from the skin surface and sweating rate (Di et al., 1997; Khan et al., 2012). The greater cutaneous vasodilatation after niacin supplementation is caused by prostaglandin D produced by epidermal Langerhans cells, which acted on vascular endothelial prostaglandin D2 receptors (Maciejewski et al., 2006; Cheng et al., 2006). When rumen-protected niacin was fed to Holstein cattle (12 g/day) during heat stress, the treatment group showed a small but detectable reduction in the rectal and vaginal temperature over the control group (Zimbelman et al., 2010). A study shows that 800 ppm niacin supplementation to lactating crossbred cows resulted in better heat stress alleviation (Khan et al., 2018). Another study conducted on Jinjiang cattle under heat-stress conditions showed that adding niacin reduced the body

temperature and respiratory rate ($P < 0.05$), and the reason is considered as niacin supplementation may alleviate heat stress by proliferating those bacteria belonging to the phylum *Succiniclaticum*, which may further contribute to the digestion of cellulose and the improvement of the metabolic function (Zou et al., 2022).

A combination of these micronutrients in adequate quantity will be a better alternative for reducing the detrimental effects of heat stress. Supplementation of trace elements (zinc, selenium, copper, cobalt, iodine, and manganese) and vitamins (Vitamin A, Vitamin D₃, and Vitamin E) above the National Research Council recommendation using a sustained-release source had positive effects on reproductive and lactation performance of dairy animals kept under thermal stress condition (Khorsandi et al., 2016).

CONCLUSION

The most adverse effects in heat-stressed dairy animals are pronounced reduction in feed intake and milk production. Concerning these issues, several nutritional strategies using micronutrients summarized in this article are capable of significantly increasing feed intake and milk production in heat-stressed dairy animals. Supplementation of micronutrients will also play an auxiliary function in heat-stressed dairy animals, such as the amelioration of defects in immune function, reproductive performance, heat dissipation, water intake, energy balance, and mammary health. So adequate amount of micronutrient supplementation in feed improves the growth, productivity, and overall well-being of the animal during heat stress.

REFERENCE

- 1) Arthington, J. D., Corah, L. R., Minton, J. E., Elsasser, T. H., & Blecha, F. (1997). Supplemental dietary chromium does not influence ACTH, cortisol, or immune responses in young calves inoculated with bovine herpesvirus-1. *Journal of animal science*, 75(1), 217-223.
- 2) Batal, A. B., Parr, T. M., & Baker, D. H. (2001). Zinc bioavailability in tetrabasic zinc chloride and the dietary zinc requirement of young chicks fed a soy concentrate diet. *Poultry Science*, 80(1), 87-90.
- 3) Bin-Jumah, M., Abd El-Hack, M. E., Abdelnour, S. A., Hendy, Y. A., Ghanem, H. A., Alsafy, S. A., ... & Momenah, M. A. (2020). Potential use of chromium to combat thermal stress in animals: A review. *Science of The Total Environment*, 707, 135996.

- 4) Bin-Jumah, M., Abd El-Hack, M. E., Abdelnour, S. A., Hendy, Y. A., Ghanem, H. A., Alsafy, S. A., ... & Momenah, M. A. (2020). Potential use of chromium to combat thermal stress in animals: A review. *Science of The Total Environment*, 707, 135996.
- 5) Block, E. (1994). Manipulation of dietary cation-anion difference on nutritionally related production diseases, productivity, and metabolic responses of dairy cows. *Journal of Dairy Science*, 77(5), 1437-1450.
- 6) Brigelius-Flohé, R., & Traber, M. G. (1999). Vitamin E: function and metabolism. *The FASEB Journal*, 13(10), 1145-1155.
- 7) C.F. Vargas-Rodriguez, K. Yuan, E.C. Titgemeyer, L.K. Mamedova, K.E. Griswold, B.J. Bradford. Effects of supplemental chromium propionate and rumen-protected amino acids on productivity, diet digestibility, and energy balance of peak-lactation dairy cattle *J. Dairy Sci.*, 97 (2014), pp. 3815-3821
- 8) Conte, G., Ciampolini, R., Cassandro, M., Lasagna, E., Calamari, L., Bernabucci, U., & Abeni, F. (2018). Feeding and nutrition management of heat-stressed dairy ruminants. *Italian Journal of Animal Science*, 17(3), 604-620.
- 9) DalleZotte, A., & Szendrő, Z. (2011). The role of rabbit meat as functional food. *Meat science*, 88(3), 319-331.
- 10) Di Costanzo, A., Spain, J. N., & Spiers, D. E. (1997). Supplementation of nicotinic acid for lactating Holstein cows under heat stress conditions. *Journal of Dairy Science*, 80(6), 1200-1206.
- 11) El-Nouty, F. D., Elbanna, I. M., Davis, T. P., & Johnson, H. D. (1980). Aldosterone and ADH response to heat and dehydration in cattle. *Journal of Applied Physiology*, 48(2), 249-255.
- 12) Hill, K. E., Wu, S., Motley, A. K., Stevenson, T. D., Winfrey, V. P., Capecchi, M. R., ... & Burk, R. F. (2012). Production of selenoprotein P (Sepp1) by hepatocytes is central to selenium homeostasis. *Journal of biological chemistry*, 287(48), 40414-40424.
- 13) Hu, W., & Murphy, M. R. (2004). Dietary cation-anion difference effects on performance and acid-base status of lactating dairy cows: A meta-analysis. *Journal of Dairy Science*, 87(7), 2222-2229.
- 14) Jarosz, M.; Olbert, M.; Wyszogrodzka, G.; Młyniec, K.; Librowski, T. Antioxidant and anti-inflammatory effects of zinc. Zinc-dependent NF-κB signaling. *Inflammopharmacology* 2017, 25, 11–24.

- 15) Jing, M. Y., Sun, J. Y., Zi, N. T., Sun, W., Qian, L. C., & Weng, X. Y. (2007). Effects of zinc on hepatic antioxidant systems and the mRNA expression levels assayed by cDNA microarrays in rats. *Annals of Nutrition and Metabolism*, 51(4), 345-351.
- 16) Kargar, H., Janssen, R., Ghesmati, A., & Vatankhah, A. (2022). Rumen-protected zinc–methionine dietary inclusion alters dairy cow performances, and oxidative and inflammatory status under long-term environmental heat stress. *Frontiers in veterinary science*, 9.
- 17) Khan, N. A. Z. A. M., Kewalramani, N., Mahajan, V. I. K. A. S., Haq, Z., & Kumar, B. (2018). Effect of supplementation of niacin on physiological and blood biochemical parameters in crossbred cows during heat stress. *Indian Journal of Animal Sciences*, 88(1), 68-75.
- 18) Khan, N., Kewalramani, N., and Kant, K. (2012). Niacin supplementation for lactating cows under heat stress conditions - a review. *Agric. Rev.* 33, 143–149.
- 19) Khorsandi, S., Riasi, A., Khorvash, M., Mahyari, S. A., Mohammadpanah, F., & Ahmadi, F. (2016). Lactation and reproductive performance of high producing dairy cows given sustained-release multi-trace element/vitamin ruminal bolus under heat stress condition. *Livestock Science*, 187, 146-150.
- 20) Kielczykowska, M.; Kocot, J.; Paździor, M.; Musik, I. Selenium—A fascinating antioxidant of protective properties. *Adv. Clin. Exp. Med.* 2018
- 21) Kingma, B. R., Frijns, A. J., Schellen, L., & van MarkenLichtenbelt, W. D. (2014). Beyond the classic thermoneutral zone: including thermal comfort. *Temperature*, 1(2), 142-149.
- 22) Kumar N, Krishnani KK, Gupta SK, Singh NP. Selenium Nanoparticles Enhanced Thermal Tolerance and Maintain Cellular Stress Protection of Pangasius Hypophthalmus Reared Under Lead and High Temperature. *Respir Physiol Neurobiol* (2017) 246:107–16. doi: 10.1016/j.resp.2017.09.006
- 23) Kumar, M., Kaur, H., Deka, R. S., Mani, V., Tyagi, A. K., & Chandra, G. (2015). Dietary inorganic chromium in summer-exposed buffalo calves (*Bubalus bubalis*): effects on biomarkers of heat stress, immune status, and endocrine variables. *Biological trace element research*, 167(1), 18-27.
- 24) Kumar, M., Kaur, H., Tyagi, A., Mani, V., Deka, R. S., Chandra, G., & Sharma, V. K. (2013). Assessment of chromium content of feedstuffs, their estimated requirement, and effects of dietary chromium supplementation on nutrient utilization, growth

performance, and mineral balance in summer-exposed buffalo calves (*Bubalus bubalis*). *Biological trace element research*, 155(1), 29-37.

- 25) Liu, F.; Celi, P.; Chauhan, S.S.; Cottrell, J.J.; Leury, B.J.; Dunshea, F.R. A short-term supranutritional vitamin E supplementation alleviated respiratory alkalosis but did not reduce oxidative stress in heat stressed pigs. *Asian Australas J. Anim. Sci.* 2018, 31, 263–269.
- 26) Maares, M.; Haase, H. A Guide to Human Zinc Absorption: General Overview and Recent Advances of In Vitro Intestinal Models. *Nutrients* 2020, 12, 762.
- 27) Maciejewski-Lenoir, D., Richman, J. G., Hakak, Y., Gaidarov, I., Behan, D. P., & Connolly, D. T. (2006). Langerhans cells release prostaglandin D2 in response to nicotinic acid. *Journal of Investigative Dermatology*, 126(12), 2637-2646.
- 28) Mallonee, P. G., Beede, D. K., Collier, R. J., & Wilcox, C. J. (1985). Production and physiological responses of dairy cows to varying dietary potassium during heat stress. *Journal of dairy science*, 68(6), 1479-1487.
- 29) McNamara, J. P., & Valdez, F. (2005). Adipose tissue metabolism and production responses to calcium propionate and chromium propionate. *Journal of Dairy Science*, 88(7), 2498-2507.
- 30) Mehdi, Y.; Hornick, J.-L.; Istasse, L.; Dufrasne, I. Selenium in the Environment, Metabolism and Involvement in Body Functions. *Molecules* 2013, 18, 3292–3311.
- 31) Min, L., Li, D., Tong, X., Nan, X., Ding, D., Xu, B., & Wang, G. (2019). Nutritional strategies for alleviating the detrimental effects of heat stress in dairy cows: a review. *International Journal of Biometeorology*, 1-20.
- 32) Mir, S. H., Pal, R. P., Mani, V., Malik, T. A., Ojha, L., & Yadav, S. (2018). Role of dietary minerals in heat-stressed poultry: A review.
- 33) Monem, U. A., & El-Shahat, K. H. (2011). Effect of different dietary levels of inorganic zinc oxide on ovarian activities, reproductive performance of Egyptian Baladi ewes and growth of their lambs. *Bulgarian Journal of Veterinary Medicine*, 14(2), 116-123.
- 34) Mousavi, F., Karimi-Dehkordi, S., Kargar, S., & Khosravi-Bakhtiari, M. (2019). Effects of dietary chromium supplementation on calf performance, metabolic hormones, oxidative status, and susceptibility to diarrhea and pneumonia. *Animal Feed Science and Technology*, 248, 95-105.

- 35) N. Sirirat, J.J. Lu, A.T.Y. Hung, S.Y. Chen, T.F. Lien, Effects different levels of nanoparticles chromium picolinate supplementation on growth performance, mineral retention, and immune responses in broiler chickens *J. Agric. Sci.*, 4 (2012), p. 12,
- 36) Odom, T. (1989). Thin egg shells in hot weather. A matter of survival. Feedstuffs. *The Miller Publishing Company, Minnetonka, MN*, 24, 20-21.
- 37) Oliveira, R. C., Guerreiro, B. M., Junior, N. M., Araujo, R. L., Pereira, R. A. N., & Pereira, M. N. (2015). Supplementation of prepartum dairy cows with β -carotene. *Journal of dairy science*, 98(9), 6304-6314.
- 38) Oltramari, C. E., Pinheiro, M. D. G., de Miranda, M. S., Arcaro, J. R., Castelani, L., Toledo, L. M., ... & Júnior, I. A. (2014). Selenium sources in the diet of dairy cows and their effects on milk production and quality, on udder health and on physiological indicators of heat stress. *Italian Journal of Animal Science*, 13(1), 2921.
- 39) Ortega, A. D. S. V., Babinszky, L., Ozsváth, X. E., Oriedo, O. H., & Szabó, C. (2022). The Effect of Heat Stress and Vitamin and Micro-Mineral Supplementation on Some Mineral Digestibility and Electrolyte Balance of Pigs. *Animals*, 12(3), 386.
- 40) Padayatty, S.J.; Levine, M. Vitamin C: The known and the unknown and Goldilocks. *Oral Dis.* 2016, 22, 463–493.
- 41) Padilla, L., Matsui, T., Kamiya, Y., Kamiya, M., Tanaka, M., & Yano, H. (2006). Heat stress decreases plasma vitamin C concentration in lactating cows. *Livestock Science*, 101(1-3), 300-304.
- 42) Patel, B., Kumar, N., Jain, V., Ajithakumar, H. M., Kumar, S., Raheja, N., ... & SINGH, S. (2017). Zinc supplementation improves reproductive performance of Karan-Fries cattle. *Indian Journal of Animal Reproduction*, 38(1), 20-22.
- 43) Ranjan, R., Ranjan, A., Dhaliwal, G. S., & Patra, R. C. (2012). L-Ascorbic acid (vitamin C) supplementation to optimize health and reproduction in cattle. *Veterinary Quarterly*, 32(3-4), 145-150.
- 44) Reboul, E. Vitamin E Bioavailability: Mechanisms of Intestinal Absorption in the Spotlight. *Antioxidants* 2017, 6, 95.
- 45) Rivas, R. O., Marins, T. N., Weng, X., Monteiro, A. P. A., Guo, J., Gao, J., ... & Tao, S. (2021). Effects of evaporative cooling and dietary zinc source on heat shock responses and mammary gland development in lactating dairy cows during summer. *Journal of Dairy Science*, 104(4), 5021-5033.
- 46) S. Lashkari, M. Habibian, S.K. Jensen. A Review on the role of chromium supplementation in ruminant nutrition—effects on productive performance, blood

- metabolites, antioxidant status, and immunocompetence *Biol. Trace Elem. Res.*, 186 (2018), pp. 305-321
- 47) Sahin, K., Sahin, N., Kucuk, O., Hayirli, A., & Prasad, A. S. (2009). Role of dietary zinc in heat-stressed poultry: A review. *Poultry science*, 88(10), 2176-2183.
- 48) Sanchez, W. K., McGuire, M. A., & Beede, D. K. (1994). Macromineral nutrition by heat stress interactions in dairy cattle: Review and original research. *Journal of Dairy Science*, 77(7), 2051-2079.
- 49) Sejian, V., Valtorta, S., Gallardo, M., & Singh, A. K. (2012). Ameliorative measures to counteract environmental stresses. In *Environmental stress and amelioration in livestock production* (pp. 153-180). Springer, Berlin, Heidelberg.
- 50) Shan, Q., Ma, F. T., Jin, Y. H., Gao, D., Li, H. Y., & Sun, P. (2020). Chromium yeast alleviates heat stress by improving antioxidant and immune function in Holstein mid-lactation dairy cows. *Animal Feed Science and Technology*, 269, 114635.
- 51) Sheikh, A. A., Aggarwal, A., Indu, B., & Aarif, O. (2017). Inorganic zinc supplementation modulates heat shock and immune response in heat stressed peripheral blood mononuclear cells of periparturient dairy cows. *Theriogenology*, 95, 75-82.
- 52) Siegel, H. S. 1995. Stress, strains and resistance. *Br. Poult. Sci.* 36:3–22.
- 53) Sivakumar, A.V.N.; Singh, G.; Varshney, V.P. Antioxidants Supplementation on Acid Base Balance during Heat Stress in Goats. *Asian Australas. J. Anim. Sci.* 2010, 23, 1462–1468.
- 54) Sultana, J. R., Chandra, A. S., Ramana, D. B. V., Raghunandan, T., Prakash, M. G., & Venkateswarlu, M. (2022). Effect of Supplemental Chromium, Vitamin E and Selenium on Biochemical and Physiological Parameters of Holstein Friesian Calves under Heat Stress. *Indian Journal of Animal Research*, 56(8), 921-927.
- 55) Suman, M., Kaur, H., Phondba, B. T., Meena, B. S., & Khan, N. (2018). Influence of varying dietary Cation-Anion difference based diets on blood biochemical and immunological parameters in crossbred calves during summer. *Indian Journal of Animal Nutrition*, 35(3), 290-297.
- 56) Sun LL, Gao ST, Wang K, Xu JC, Sanz-Fernandez MV, Baumgard LH, et al. Effects of Source on Bioavailability of Selenium, Antioxidant Status, and Performance in Lactating Dairy Cows During Oxidative Stress-Inducing Conditions. *J Dairy Sci* (2019) 102:311–9. doi: 10.3168/jds.2018-14974

- 57) T. Leiva, R.F. Cooke, A.P. Brandão, U. Pardelli, R.O. Rodrigues, F.N. Corrá, J. Vasconcelos. Effects of concentrate type and chromium propionate on insulin sensitivity, productive and reproductive parameters of lactating dairy cows consuming excessive energy *Animal*, 11 (2017), pp. 436-444
- 58) Tanaka, M., Kamiya, Y., Suzuki, T., & Nakai, Y. (2011). Changes in oxidative status in periparturient dairy cows in hot conditions. *Animal science journal*, 82(2), 320-324.
- 59) Tang J, Cao L, Jia G, Liu G, Chen X, Tian G, et al. The Protective Effect of Selenium From Heat Stress-Induced Porcine Small Intestinal Epithelial Cell Line (IPEC-J2) Injury is Associated With Regulation Expression of Selenoproteins. *Brit J Nutr* (2019) 122:1081–90. doi: 10.1017/S0007114519001910
- 60) Underwood, E. J. and Suttle, N. F. 1999. In *The mineral nutrition of livestock*, 3rd ed. CABI Publishing, CAB International, Wallingford, Oxon, UK. Technology 93: 193-203.
- 61) Weng, X., Monteiro, A. P. A., Guo, J., Li, C., Orellana, R. M., Marins, T. N., ... & Tao, S. (2018). Effects of heat stress and dietary zinc source on performance and mammary epithelial integrity of lactating dairy cows. *Journal of dairy science*, 101(3), 2617-2630.
- 62) Wildman, C. D., West, J. W., & Bernard, J. K. (2007). Effect of dietary cation-anion difference and dietary crude protein on performance of lactating dairy cows during hot weather. *Journal of dairy science*, 90(4), 1842-1850.
- 63) Yen, J. T., Pond, W. G., & Prior, R. L. (1981). Calcium chloride as a regulator of feed intake and weight gain in pigs. *Journal of Animal Science*, 52(4), 778-782.
- 64) Zheng, Y., Xie, T., Li, S., Wang, W., Wang, Y., Cao, Z., & Yang, H. (2021). Effects of selenium as a dietary source on performance, inflammation, cell damage, and reproduction of livestock induced by heat stress: a review. *Frontiers in Immunology*, 12.
- 65) Zimbelman, R. B., Baumgard, L. H., & Collier, R. J. (2010). Effects of encapsulated niacin on evaporative heat loss and body temperature in moderately heat-stressed lactating Holstein cows. *Journal of Dairy Science*, 93(6), 2387-2394.
- 66) Zimbelman, R. B., Collier, R. J., & Bilby, T. R. (2013). Effects of utilizing rumen protected niacin on core body temperature as well as milk production and composition in lactating dairy cows during heat stress. *Animal feed science and technology*, 180(1-4), 26-33.

- 67) Zou Bicheng, Long Fan, XueFuguang, Qu Mingren, Chen Chuanbin, Zhang Xian, Xu Lanjiao, (2022). Alleviation effects of niacin supplementation on beef cattle subjected to heat stress: A metagenomic insight *Frontiers in Microbiology*13
- 68) Zou Y, Shao J, Li Y, Zhao FQ, Liu JX, Liu H. Protective Effects of Inorganic and Organic Selenium on Heat Stress in Bovine Mammary Epithelial Cells. *Oxid Med Cell Longe* (2019) 2019:1503478–1503488. doi: 10.1155/2019/1503478
- 69) Singh, A.K., Bhakat, C., Yadav, D.K., Kumari, T., Mandal, D.K., Rajput, M.S., Bhatt, N., 2020b. Effect of pre and postpartum Alphatocopherol supplementation on body measurements and its relationship with body condition, milk yield, and udder health of Jersey crossbred cows at tropical lower Gangetic region. *Journal of Entomology and Zoology Studies*, 8(1): 1499-1502.
- 70) Singh, A.K., Bhakat, C., Mandal, D.K., Chatterjee, A., 2020c. Effect of pre and postpartum alpha-tocopherol supplementation on body condition and some udder health parameters of Jersey crossbred cows at tropical lower Gangetic region. *Journal of Animal Research*, 10(5): 697-703. <http://doi:10.30954/2277-940X.05.2020.4>
- 71) Singh, A.K., Bhakat, C., Kumari, T., Mandal, D.K., Chatterjee, A., Karunakaran, M., Dutta, T.K., 2020d. Influence of pre and postpartum alpha-tocopherol supplementation on milk yield, milk quality and udder health of Jersey crossbred cows at tropical lower Gangetic region. *Veterinary World*, 13(9), 2006-2011. <http://doi:10.14202/vetworld.2020.2006-2011>
- 72) Singh, A.K., 2021. Advancements in Management Practices from Far-off Dry Period to Initial Lactation Period for Improved Production, Reproduction, and Health Performances in Dairy Animals: A Review. *International Journal of Livestock Research*, 11(3): 25-41. <http://doi:10.5455/ijlr.20200827114032>
- 73) Singh, A.K., Yadav, D.K., Bhatt, N., Sriranga, K.R., Roy, S., 2020a. Housing Management for Dairy Animals under Indian Tropical Type of Climatic Conditions-A Review. *Veterinary Research International*, 8(2), 94-99.
- 74) Singh, A.K., 2019. Influence of alteration of management practice on performances of dairy cows at lower Gangetic region. M.Sc. Thesis submitted to National Dairy Research Institute, Karnal, India.
- 75) Singh, A.K, Bhakat, C., Mohhamad, A., Chatterjee, A., Karunakaran, M., Ghosh, M.K., 2020e. Economic Analysis of Pre and Postpartum Alphatocopherol Supplementation for Milk Performance and Dry Matter Intake of Dairy Cows in

Tropical Region. International Journal of Livestock Research, 10(10): 137-143.

<http://doi:10.5455/ijlr.20200710040840>

UNDER PEER REVIEW