

## **Impact of Nitrogen and Silicon levels on yield and relative economics of transplanted rice under temperate conditions**

### **Abstract**

The field experiment entitled “Impact of Nitrogen and Silicon application on yield and relative economics of transplanted rice under temperate conditions” was conducted at Agronomy Research Farm of Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir during Kharif seasons 2014-2015. The soil of the experiment was silty clay loam in texture, neutral in reaction with medium available nitrogen (442.88kg ha<sup>-1</sup>), phosphorus (9.3kg ha<sup>-1</sup>), potassium (221.60kg ha<sup>-1</sup>) and low available silicon (280.23kg ha<sup>-1</sup>). The experiment was laidout in Factorial Randomized Completely Block Design assigning combinations of three nitrogen levels viz. 120, 150 and 180N kg ha<sup>-1</sup> and four silicon levels viz. Control, 5%, 10% and 15% Si with four replications. The results revealed that higher grain yield (76.00q ha<sup>-1</sup>), straw yield (93.97q ha<sup>-1</sup>) and harvest index (44.84%) were recorded with 120Nkg ha<sup>-1</sup>. So far as silicon levels are concerned, 15%Si observed higher grain yield (76 q ha<sup>-1</sup>), straw yield (93.97 q ha<sup>-1</sup>) and harvest index (44.84%). Relative economics revealed that 120Nkg ha<sup>-1</sup> with 15%Si realized higher benefit cost ratio (Rs. 2.12).

**Keywords:** Benefit Cost Ratio, Nitrogen, Rice, Silicon and Yield.

### **Introduction**

“Rice is a staple food for more than half of the world’s population. Globally, it is grown on acreage of 164.19mha with total production of 509.87 million tonnes and productivity 3.10 ton per hectare” (DES, 2020). “In Jammu and Kashmir rice is grown on an area of 0.3million ha with a production of 0.6 million tonnes and productivity of 2 tonnes per hectare” (DES, 2020). “It is essential to the rice plant, with about 75% of leaf nitrogen associated with chloroplasts, which are physiologically important in dry matter production” (Dalling, 1995). “There are many factors affecting rice yield and quality, such as variety, accumulation temperature during the growth period, and relatively minor factors such as cultivation techniques (fertilizer rates, fertilization period, water management, etc.)” Laenoi *et al.*, 2018; Zhu *et al.*, 2017; Cao *et al.*, 2017 and Zhou *et al.*, 2018. Due to the effect of increased yield brought by the application of nitrogen fertilizer, an increasing number of nitrogen fertilizers have been applied in rice production.

“Silicon is the second most abundant element in soil after oxygen. Rice is considered to be a silicon accumulator plant and tends to actively accumulate Si to tissue concentrations of 5% or higher” (Epstein, 2002). “Reduced amount of silicon in plant produces necrosis, disturbance in leaf photosynthetic efficiency, growth retardation and reduces grain yield in cereals especially rice” (Shashidhar *et al.*, 2008). “The positive effects of silicon on rice growth and production were manifested when it was specifically supplied during the reproductive growth stage (panicle

initiation to heading) as compared to vegetative and ripening stages. Panicle initiation stage application exerted a feed-forward effect on photosynthesis coupled with increase in both stomatal conductance and biochemical capacity to fix CO<sub>2</sub> as reported” by Lavinsky *et al.*, 2016, who also surmised that “proper levels of Si in reproductive structures played an unidentified role in increasing the yields of rice. Silicon also has a major role in increasing yield attributing characters like number of spikelets, filled spikelet percentage, test weight and total grain yield” (Rani *et al.*, 1997; Ahmad *et al.*, 2013 and Jawahar *et al.*, 2015) in rice. Keeping in view the above facts, the study was designed with the objective to study the impact of Nitrogen and Silicon levels on yield and relative economics of transplanted rice under temperate conditions.

## Material and methods

The field experiment was carried out at Research Farm of Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar during 2014-2015. The mean maximum (27.50<sup>0</sup>C and 27.30<sup>0</sup>C), mean minimum temperatures (12.60<sup>0</sup>C and 13.20<sup>0</sup>C), mean maximum relative humidities (82.50% and 82.80%) and mean minimum relative humidities (55.30% and 56.20%) were recorded for the entire cropping seasons during 2014 and 2015 respectively. The total rainfall received during the experimentation period was 33.3mm during 2014 and 29.5mm during 2015. The experiment comprised two factors ( three nitrogen levels and four silicon levels) was laid out in a randomized completely block design with four replications. The factors included three nitrogen levels: N<sub>1</sub>: 120kg/ha, N<sub>2</sub>:150kg/ha and N<sub>3</sub>:180kg/ha and four silicon levels: Si<sub>0</sub>: Control, Si<sub>1</sub>:5%, Si<sub>2</sub>:10% and Si<sub>3</sub>:15%. The following observations were recorded during the experiment are:

**Grain yield:** Grain yield of each plot was recorded separately as kg plot<sup>-1</sup> and then converted into q ha<sup>-1</sup>.

**Straw yield:** Straw yield of each plot was computed by deducting the grain yield from biological yield and expressed as q ha<sup>-1</sup>.

**Harvest index:** It was calculated by the following formula:

$$H.I = \text{Economic yield} / \text{Biological yield} \times 100$$

## Relative economics

Economics of different treatments was worked out on the basis of grain and straw yield per hectare. The cost of input and output was estimated as per prevailing market rates. The benefit : cost ratio (returns per rupee invested) was determined as:

$$\text{Benefit cost ratio} = \frac{\text{Net returns}}{\text{Total cost of cultivation}}$$

## Statistical analysis

The data obtained in respect of various observations were statistically analysed by the method described by Cochran and Cox (1964) using Ne-software. The significance of 'F' and 'T' was tested at 5% level of significance. The critical difference value was determined wherever 'F' test was found to be significant.


## Result and Discussion

From the table 1 and fig. 1 indicated that the grain yield, straw yield and harvest index showed the significant variation among the nitrogen and silicon levels. Significantly higher grain yield (76.00q ha<sup>-1</sup>), straw yield (93.97q ha<sup>-1</sup>) and harvest index (44.84%) were observed with 120kg N ha<sup>-1</sup> followed by 150kg N ha<sup>-1</sup> (grain yield: 74.03q ha<sup>-1</sup>, straw yield: 91.66q ha<sup>-1</sup> and harvest index: 44.51%). It is because of higher yield attributing characters which ultimately caused the increase of rice grain yield. Similar results were reported by Huang *et al.*, 2011. "Also, the increment of grain yield by the application of nitrogen upto a certain level was reported" by Zhaowen *et al.*, 2013. Similar trend of the effect of nitrogen levels on straw yield were also reported by Pramanik and Bera (2013). However, harvest index were significantly higher with 120kgN ha<sup>-1</sup>. Rahman *et al.*, 2007, also reported similar trend. Significantly lower grain yield (68.83q ha<sup>-1</sup>) straw yield (88.19q ha<sup>-1</sup>) and harvest index (43.96%) were recorded with 180kg N ha<sup>-1</sup>. Among different silicon levels, 15% Si recorded the higher grain yield (76.76q ha<sup>-1</sup>), straw yield (94.72q ha<sup>-1</sup>) and harvest index (44.80%) followed by 10%Si (grain yield: 74.58q ha<sup>-1</sup>, straw yield: 92.89q ha<sup>-1</sup>, and harvest index: 44.40%) being at par with 5%Si (grain yield:73.21q ha<sup>-1</sup>, straw yield:91.96q ha<sup>-1</sup>, and harvest index:44.24%). It is because of silicon is responsible to control stomatal activity, photosynthesis, water use efficiency which ultimately results in better vegetative growth results in higher yield. And it may also be attributed to leaf erectness which facilitated better penetration of sunlight leading to higher photosynthetic activity of plant and higher production of carbohydrates. Similar results were also noticed by Singh *et al.*, 2006 and Pati *et al.*, 2015. Significantly lower grain yield (67.27q ha<sup>-1</sup>) straw yield (85.52q ha<sup>-1</sup>) and harvest index (43.84%) were observed with control.


**Table 1. Impact of nitrogen and silicon levels on grain yield, straw yield and harvest index of transplanted rice (Pooled data of two years)**

Treatments	Grain yield (q ha <sup>-1</sup> )	Straw yield (q ha <sup>-1</sup> )	Harvest index (%)
<b>Nitrogen levels (kg ha<sup>-1</sup>)</b>			
<b>N<sub>1</sub> :120</b>	76.00	93.97	44.84
<b>N<sub>2</sub> :150</b>	74.03	91.66	44.51
<b>N<sub>3</sub> :180</b>	68.83	88.19	43.96
<b>SEm±</b>	0.51	0.54	0.09
<b>CD(p≤0.05)</b>	1.52	1.59	0.28
<b>Silicon levels</b>			

<b>Si<sub>0</sub> :Control</b>	67.27	85.52	43.84
<b>Si<sub>1</sub> : 5%</b>	73.21	91.96	44.24
<b>Si<sub>2</sub> :10%</b>	74.58	92.89	44.40
<b>Si<sub>3</sub> :15%</b>	76.76	94.72	44.80
<b>SEm±</b>	0.57	0.63	0.12
<b>CD(p≤0.05)</b>	1.71	1.84	0.37



**Fig. 1 (a)**



**Fig. 1 (b)**

**Fig. 1 (a and b). Impact of nitrogen and silicon levels on grain yield, straw yield and harvest index of transplanted rice (Pooled data of two years)**

The data relative economics presented in table 2 revealed that benefit cost ratio remained higher (Rs.2.13) when 120 kg N ha<sup>-1</sup> and 15% Si were applied. The same treatment also recorded the higher net returns of Rs.106107. The efficiency of a treatment is finally decided in terms of economics (benefit cost ratio) of the treatment and it might be due to the higher grain and straw yield and minimum cost of nitrogen fertilizers as cost of silicon does not have a greater effect.

**Table 2. Relative economics of transplanted rice as influenced by nitrogen and silicon levels (pooled data of two years)**

Treatments	Cost of Cultivation (Rs. ha <sup>-1</sup> )	Gross returns (Rs. ha <sup>-1</sup> )	Net returns (Rs. ha <sup>-1</sup> )	B:C
N <sub>1</sub> Si <sub>0</sub>	48522	138929	90407	1.86
N <sub>1</sub> Si <sub>1</sub>	49146	148280	99134	2.01
N <sub>1</sub> Si <sub>2</sub>	49490	150174	100684	2.03
N <sub>1</sub> Si <sub>3</sub>	49834	155941	106107	2.13
N <sub>2</sub> Si <sub>0</sub>	48687	132364	83677	1.71
N <sub>2</sub> Si <sub>1</sub>	49311	145581	96270	1.95
N <sub>2</sub> Si <sub>2</sub>	49655	147330	97675	1.96
N <sub>2</sub> Si <sub>3</sub>	49999	153206	103207	2.06
N <sub>3</sub> Si <sub>0</sub>	48852	128447	79595	1.62
N <sub>3</sub> Si <sub>1</sub>	49476	137549	88073	1.78

<b>N<sub>3</sub> Si<sub>2</sub></b>	49820	139223	89403	1.79
<b>N<sub>3</sub> Si<sub>3</sub></b>	50164	140566	90402	1.80

## Conclusion

From this study we concluded that silicon applications can significantly improve rice yield if applied at proper time with feasible concentration. Optimum nitrogen fertilizer is an important practice for increasing rice yield. Considering the influence of nitrogen and silicon levels, it can be suggested that for realizing economically higher grain yield of rice with 120Nkg ha<sup>-1</sup> and 15% Si based nutrient management, as it helped to achieve the higher net returns (Rs. 106107) and the higher B:C ratio (Rs. 2.13).

## References

- Ahmad, A., Afzal, M., Ahmad, A. U.H. and Tahir, M. 2013. Effect of foliar application of silicon on yield and quality of rice (*Oryza sativa* L.). *Cercetari Agronomic in Moldova*. **XLVI** (3):155.
- Cao, X.M., Sun, H.Y., Wang, C.G., Ren, X.J., Liu, H.F. and Zhang, Z.J. 2017. Effects of Late-Stage Nitrogen Fertilizer Application on the Starch Structure and Cooking Quality of Rice. *Journal of the Science of Food and Agriculture*. **98**: 2332-2340.
- Cochran, W.G. and Cox, G.M. 1964. *Experimental Designs*.2. Ufl. John Wiley & Sons, London 1964. XIV+617S., Preis 68 s
- Dalling, M.J. 1995. The physiological basis of nitrogen redistributing during filling in cereals pp.55-71. In: Exploitation of physiological and genetic variability to enhance crop productivity. *American Society of Plant Physiologists*, Rockville, MD.
- DES, 2020. Economic Survey, Directorate of Economics and Statistics. Government of Jammu and Kashmir. **123**: 129-130.
- Epstein, E. 2002. Silicon in plants: Facts vs. concepts. In: Silicon in Agriculture. (Ed. L.E. Datnoff, G.H. Snyder and G.H. Korndorfer) *Elsevier, Amsterdam* pp. 1-15.
- Huang, M., Zou, Y.B., Jiang, P., Bing, X.I.A., Md. I. and Ao, H.J. 2011. Relationship between grain yield and yield components in super hybrid rice. *Agricultural Sciences in China*. **10**: 1537-1544.
- Jawahar, S., Vijayakumar, D., Bommera, R., Neeru Jain and Jeevanandham. 2015. Effect of silixol granules on growth and yield of rice. *International Journal of Current Research and Academic Review*. **3**(1): 168-174.

- Laenoi, S., Rerkasem, B. and Lordkaew, S. 2018. Seasonal Variation in Grain Yield and Quality in Different Rice Varieties. *Field Crops Research*. **221**: 350-357.
- Lavinsky, A.O., Detmann, K.C., Josimar, V.R., Avila, R.T., Sanglard, M.L., Pereira, L.F., Sanglard, M.V.P.L., Rodrigues, A.F., Wagner, L.A and DaMatta, F.M. 2016. Silicon improves rice grain yield and photosynthesis specifically when supplied during the reproductive growth stage. *Journal of Plant Physiology*. **206**: 125-132.
- Pati, S., Pal, B. and Badole, S. 2015. Effect of silicon fertilization on growth, yield and nitrogen uptake of rice. *Communication in Soil Science and Plant Analysis*. **47**(3):1081-1086.
- Pramanik, K. and Bera, A.K. 2013. Effect of seedling age and nitrogen fertilizer on growth, chlorophyll content, yield and economics of hybrid rice (*Oryza sativa* L. ). *International Journal of Agronomy and Plant Production*. **4**:3489-3499.
- Rahman, M.H., Khatun, M.M., Mamun, M.A.A., Islam, M.Z. and Islam, M.R. 2007. Effect of number of seedling hill-1 and nitrogen levels on growth and yield of BRRI dhan 32. *Journal of Soil Natural*. **1**(2): 01-07.
- Rani, Y.A., Narayanan, A., Devi, V. and Subbaramamma, P. 1997. The effect of silicon application on growth and yield of rice plants. *Annals in Plant Physiology*. **11**:125-128.
- Shashidhar, H.E., Chandrashekhar, N., Narayanaswamy, C., Mehendra, A.C. and Prakash, N.B. 2008. Calcium silicate as silicon source and its interaction with nitrogen in aerobic rice. Silicon in Agriculture: 4<sup>th</sup> International Conference 26-31 October, South Africa pp. 93.
- Singh, R., Ahuja, U., and Ahuja, S.C. 2006. Basmati for prosperity. *Indian Farming*. **56**:33-36.
- Zhaowen, M.O., Shenggang, P., Bingshun, L., Hua, T., Meiyang, D., Qingshan, L., Yijun, J. and Xiangru, T. 2013. Growth and Yield response of five super rice (*Oryza sativa* L.) varieties to nitrogen fertilizer rates. *Journal of Food Agriculture and Environment*. **11** (3-4):1257-1262.
- Zhou, C.C., Huang, Y.C., Jia, B.Y., Wang, Y., Wang, Y., Xu, Q., Li, R.F., Wang, S. and Dou, F.G. 2018. Effects of Cultivar, Nitrogen Rate, and Planting Density on Rice-Grain Quality. *Agronomy*. **8**: 246.
- Zhu, D.W., Zhang, H.C., Guo, B.W., Xu, K., Dai, Q.G., Wei, H.Y., Gao, H., Hu, Y.J., Cui, P.Y. and Huo, Z.Y. 2017. Effects of Nitrogen Level on Yield and Quality of Japonica Soft Super Rice. *Journal of Integrative Agriculture*. **16**: 1018-1027.