

Chemical constituents from the stem bark of *Ficus thonningii* and their chemotaxonomic significance

Abstract

Background of the Study: Tropical plants of the *Ficus* genus (Moraceae) are among the earliest fruit trees that humans have cultivated. Since ancient times, many folk medicines have used species of this genus to treat a variety of ailments. Evidence from earlier investigations has shown these plants contain abundant secondary metabolites with a variety of structural properties and biological functions.

Place and Duration of Study: The research was carried out at the University of Nairobi (Faculty of Science and Technology, Department of Chemistry) from January to June 2022.

Aim: The study focuses on isolating and identifying secondary metabolites from the stem bark of *Ficus thonningii* Blume found in Kenya and their chemotaxonomic significance.

Methodology: Dried powdered stem bark of *Ficus thonningii* was extracted by maceration at room temperature using CH₂Cl₂/CH₃OH (1:1) to yield a crude extract which was fractionated in a chromatographic column (CC) using silica gel (60 – 120 mesh) as an adsorbent eluting with EtOAc/*n*-hexane followed by CH₃OH/EtOAc. The fractions were purified using silica gel (70 – 230, 230 – 400 mesh) CC and chromatotron eluting with solvents of different polarity, as well as a crystallization technique. Structures of the isolated compounds were elucidated and identified using the spectroscopic method (NMR (1D and 2D)) and by comparison with reported literature data.

Results: Yukovanol (1), 5,7,4'-trihydroxy-3'-(2-hydroxy-3-methyl-3-butenyl)isoflavone (2), cajanin (3), taxifolin (4), protocatechuic acid (5), saccharose (6), and stigmasterol (7) were isolated.

Conclusion: The chemotaxonomic significance of the isolated phytochemicals demonstrates the taxonomic position of *F. thonningii* in the genus *Ficus* and explains its multiple ethnomedicinal applications.

Keywords: *Ficus thonningii*, Moraceae, Flavonoids, Phenolic acid, Chemotaxonomy

1. INTRODUCTION

The genus *Ficus* (Moraceae) consists of over 850 species widely distributed in tropical and subtropical countries across the globe [1–3]. *Ficus* is among the leading diversified plant genera including creepers, climbers, and stranglers. It also has free-standing deciduous and evergreen trees. *Ficus* species are distinguished by their distinctive syconium-like inflorescence and symbiotic connection with Agaonidae wasps, which pollinate their species exclusively [4,5]. In the African region, 112 species of *Ficus* are recognized currently [6], with 37 being distributed in Kenya within 0 – 2300 m altitude, including *Ficus thonningii* [7,8].

Ficus thonningii (commonly called 'Mugumo' by Kikuyu in Kenya) has a dense, rounded to spreading crown, often epiphytically initially, and is multi-stemmed, evergreen, or short deciduous. The shiny green leaves of *F. thonningii* are alternate, oval (up to 12 cm) with rounded tip and tapering base, whereas the young leaves are pale and finely hairy. The arial roots are frequently present, and the bark is grey, thin, and smooth [8,9]. *Ficus* species such as *Ficus thonningii* have long been used in indigenous medical systems to treat diarrhea, gonorrhoea, diabetes mellitus, inflammation, and induce lactation [10–13].

Phytochemical investigations on the genus *Ficus* have revealed flavonoids (especially isoflavones) [14–18] and terpenoids [19–25] as the major chemical constituents. Phytochemicals such as alkaloids [26–28], coumarins [29–31], and phenolic acids [32,33] have also been reported from various species of *Ficus*. However, the systematic phytochemical studies of *F. thonningii* from East Africa has hitherto not been reported. Therefore, we herein report the isolation and identification of phytoconstituents from the stem bark of *F. thonningii* found in Kenya, and their chemotaxonomic significance.

2. MATERIALS AND METHODS

2.1 Plant collection

The stem bark of *Ficus thonningii* was collected from the Riverside drive in September 2020 (1°16'19.2"S 36°48'07.6"E) in Nairobi County, Kenya. The plant was identified by Mr. Patrick C. Mutiso, a taxonomist from the Faculty of Science and Technology (FST), University of Nairobi, Kenya, where a voucher specimen (HIUON 2021/004) was deposited.

2.2 Experimental Procedures

Silica gel 60 – 120, 70 – 230, and 230 – 400 meshes as solid phases for column chromatography (CC), and Sephadex LH-20 (25–100 μm , Sigma Aldrich) were used. Thin Layer Chromatography (TLC) was carried out on pre-coated silica gel 60 plates (0.25 mm; Merck, Darmstadt, Germany). Compounds were visualized under UV light and further by spraying with $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$ (5 %, v/v). NMR spectra were performed on Bruker Advance Neo 500 MHz spectrometer using standard pulse sequences and referenced to residual solvent signals.

2.3 Extraction and isolation

Dried powdered stem bark of *Ficus thonningii* (1.7 Kg) was extracted at room temperature with $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ (1:1, 6 L, 24 h \times 3) by maceration to afford 85.7 g of crude extract. 80 g of the stem bark crude extract was fractionated in a chromatographic column using silica gel (60 – 120 mesh) as an adsorbent eluting with EtOAc/*n*-hexane (0:10, 1:9, 1.5:8.5, 2:8, 3:7, 3.5:6.5; 4:6, 4.5:5.5, 1:1 and 10:0) followed by $\text{CH}_3\text{OH}/\text{EtOAc}$ (1:9 and 2:8) to yield twelve fractions ($\text{HIF}_{\text{A-L}}$). Fraction HIF_{I} (2.8 g) was subjected to silica gel (70 – 230 mesh) CC eluting with EtOAc/*n*-hexane (1:9 to 10:0), resulting in the isolation of compounds **1** (5.1 mg) and **2** (3.0 mg). Purification of fraction HIF_{J} (2.81 g) using silica gel (70 – 230 mesh) CC eluting with a gradient of EtOAc/*n*-hexane (0.5:9.5 to 10:0), resulted in 85 fractions of 30 mL each. The fractions were pooled based on their TLC profiles into two main subfractions ($\text{HIF}_{\text{J1-2}}$).

Subfraction HIF_{J1} (700 mg) was purified on a silica gel (70 -230 mesh) CC eluting with a gradient polarity of EtOAc/*n*-hexane (0:10 to 1.5:8.5) to afford a semi-pure compound. The semi-pure compound was purified on a chromatotron using a gradient of EtOAc/*n*-hexane (2:8 to 10:0) to yield compound **3** (13.4 mg). Subfraction HIF_{J2} (900 mg) was also purified using silica gel (230 – 400 mesh) CC eluting with a gradient of EtOAc/*n*-hexane (1:9 to 6:4) to afford compounds **4** (1.06 mg) and **5** (2.82 mg). Fraction HIF_{L} (3.5 g) afforded brown crystals which, after filtration and recrystallization in CH_3OH , compound **6** (43.2 mg) was obtained. Similarly, compound **7** (33.6 mg) was crystallized in fraction HIF_{F} (1.02 g), and the crystals were repeatedly washed with *n*-hexane to obtain the compound.

3. RESULTS AND DISCUSSION

Based on the NMR (1D and 2D) and in comparison with the previously published data, the isolated compounds were identified as yukovanol (**1**) [34,35], 5,7,4'-trihydroxy-3'-(2-hydroxy-3-methyl-3-butenyl)isoflavone (**2**) [36], cajanin (**3**) [37], taxifolin (**4**) [38], protocatechuic acid (**5**) [39], saccharose (**6**) [40], and stigmasterol (**7**) [41], (Fig. 1). The NMR data of the isolated compounds are presented below.

Yukovanol (**1**): Yellow powder; ^1H NMR (CD_3OD , 500 MHz): δ (ppm) 5.02 (1H, d, $J = 11.6$ Hz, H-2), 4.60 (1H, d, $J = 11.6$ Hz, H-3), 5.91 (1H, s, H-6), 7.37 (2H, d, $J = 8.6$ Hz, H-2'/6'), 6.85 (2H, d, $J = 8.6$ Hz, H-3'/5'), 6.62 (1H, d, $J = 10.1$ Hz, H-4''), 5.62 (1H, d, $J = 10.1$ Hz, H-5''), 1.44 (6H, d, $J = 2.4$ Hz, H-7''/8''); ^{13}C NMR (CD_3OD , 125 MHz): δ : 85.1 (CH, C-2), 73.7 (CH, C-3), 199.2 (C, C-4), 102.4 (C, C-4a), 163.9 (C, C-5), 97.1 (CH, C-6), 163.6 (C, C-7), 104.1 (C, C-8), 159.2 (C, C-8a), 129.1 (C, C-1'), 130.4 (CH, C-2'/6'), 116.2 (CH, C-3'/5'), 159.3 (C, C-4'), 116.0 (CH, C-4''), 127.7 (CH, C-5''), 79.5 (C, C-6''), 28.6 (CH_3 , C-7''), 28.5 (CH_3 , C-8'') [34,35].

5,7,4'-trihydroxy-3'-(2-hydroxy-3-methyl-3-butenyl)isoflavone (**2**): Yellow powder; ^1H NMR (CD_3OD , 500 MHz): δ (ppm) 7.87 (1H, s, H-2), 6.28 (1H, d, $J = 2.2$ Hz, H-6), 6.37 (1H, d, $J = 2.2$ Hz, H-8), 7.28 (1H, dd, $J = 8.3, 2.3$ Hz, H-2'), 6.94 (1H, d, $J = 8.3$ Hz, H-3'), 7.23 (1H, d, $J = 2.3$ Hz, H-6'), 2.86 (1H, dd, $J = 14.7, 2.3$ Hz, H-1a''), 2.99 (1H, dd, $J = 14.7, 8.77$ Hz, H-1b''), 4.44 (1H, m, H-2''), 4.89 (2H, s, H-4''), 1.83 (3H, s, H-5''); ^{13}C NMR (CD_3OD , 125 MHz): δ (ppm) 153.0 (CH, C-2), 123.5 (C, C-3), 180.9 (C, C-4), 106.2 (C, C-4a), 162.7 (C, C-5), 99.6 (CH, C-6), 163.4 (C, C-7), 94.3 (CH, C-8), 158.2 (C, C-8a), 123.0 (C, C-1'), 129.4 (CH, C-2'), 117.5 (CH, C-3'), 156.7 (C, C-4'), 126.2 (C, C-5'), 132.5 (CH, C-6'), 38.4 (CH_2 , C-1''), 78.4 (CH, C-2''), 147.2 (C, C-3''), 111 (CH_2 , C-4''), 18.4 (CH_3 , C-5'') [36].

Cajanin (**3**): Pale yellow powder; ^1H NMR (CD_3OD , 500 MHz): δ (ppm) 8.07 (1H, s, H-2), 6.38 (1H, m, H-6), 6.57 (1H, d, $J = 2.3$ Hz, H-8), 3.89 (3H, s, 7- OCH_3), 6.40 (1H, m, H-3'), 6.36 (1H, m, H-5'), 7.05 (1H, d, $J = 8.2$ Hz, H-6'); ^{13}C NMR (CD_3OD , 125 MHz): δ (ppm) 157.8 (CH, C-2), 122.8 (C, C-3), 182.8 (C, C-4), 107.1 (C, C-4a), 163.3 (C, C-5), 99.3 (CH, C-6), 167.3 (CH, C-7), 93.2 (CH, C-8), 159.7 (C, C-8a), 56.5 (CH_3 , 7- OCH_3), 110.6 (C, C-1'), 157.0 (C, C-2'), 104.2 (CH, C-3'), 160.3 (C, C-4'), 108.1 (CH, C-5'), 133.2 (CH, C-6') [37].

Taxifolin (**4**): Yellow solid; ^1H NMR (CD_3OD , 500 MHz): δ (ppm) 4.91 (1H, d, $J = 11.6$ Hz, H-2), 4.50 (1H, d, $J = 11.6$ Hz, H-3), 5.92 (1H, d, $J = 2.1$ Hz, H-6), 5.88 (1H, d, $J = 2.1$ Hz, H-8), 6.85 (1H, dd, $J = 8.1, 2.1$ Hz, H-2'), 6.80 (1H, d, $J = 8.1$ Hz, H-3'), 6.96 (1H, d, $J = 2.0$ Hz, H-6'); ^{13}C NMR (CD_3OD , 125 MHz): δ (ppm) 85.1 (CH, C-2), 73.7 (CH, C-3), 198.4 (C, C-4), 101.8 (C, C-4a), 164.5 (C, C-5), 97.4 (CH, C-6), 165.3 (C, C-7), 96.3 (CH, C-8), 164.5 (C, C-8a), 129.9 (C, C-1'), 120.9 (CH, C-2'), 116.1 (CH, C-3'), 147.2 (C, C-4'), 146.3 (C, C-5'), 115.9 (CH, C-6') [38].

Protocatechuic acid (**5**): White powder; ^1H NMR (CD_3OD , 500 MHz): δ (ppm) 7.43 (1H, d, $J = 1.9$ Hz, H-2), 6.77 (1H, d, $J = 8.2$ Hz, H-5), 7.41 (1H, dd, $J = 8.2, 1.9$ Hz, H-6); ^{13}C NMR (CD_3OD , 125 MHz): δ (ppm) 125.0 (C, C-1), 117.7 (CH, C-2), 145.9 (C, C-3), 150.9 (C, C-4), 115.6 (CH, C-5), 123.7 (CH, C-6), 170.0 (CO) [39].

Saccharose (**6**): Colorless crystals; ^1H NMR (D_2O , 500 MHz): δ (ppm) 5.40 (1H, d, $J = 3.8$ Hz, H-1), 3.54 (1H, dd, $J = 10.0, 3.8$ Hz, H-2), 3.75 (1H, t, $J = 10.0, 9.1$ Hz, H-3), 3.46 (1H, t, $J = 9.5$ Hz, H-4), 3.83 (1H, m, H-5), 3.80 (2H, m, H-6), 3.66 (2H, s, H-1'), 4.20 (1H, d, $J = 8.8$ Hz, H-3'), 4.04 (1H, t, $J = 8.6$ Hz, H-4'), 3.87 (1H, m, H-5'), 3.80 (2H, m, H-6'); ^{13}C NMR (CD_3OD , 125 MHz): δ (ppm) 92.1 (CH, C-1), 71.0 (CH, C-2), 72.5 (CH, C-3), 69.1 (CH, C-4), 72.3 (CH, C-5), 60.0 (CH_2 , C-6), 61.2 (CH_2 , C-1'), 103.6 (C, C-2'), 76.3 (CH, C-3'), 73.9 (CH, C-4'), 81.3 (CH, C-5'), 62.3 (CH_2 , C-6') [40].

Stigmasterol (**7**): White amorphous solid; ^1H NMR (CDCl_3 , 500 MHz): δ (ppm) 1.84 (2H, m, H-1), 1.99 (2H, m, H-2), 3.52 (1H, tt, $J = 11.6, 4.9$ Hz, H-3), 2.28 (2H, m, H-4), 5.34 (1H, dt, $J = 5.5, 1.9$ Hz, H-6), 1.50 (2H, m, H-7), 1.43 (1H, m, H-8), 0.92 (1H, m, H-9), 1.47 (2H, m, H-11), 2.01 (2H, m, H-12), 0.99 (1H, m, H-14), 1.84 (2H, m, H-15), 1.25 (2H, m, H-16), 1.10 (2H, m, H-17), 0.68 (3H, s, H-18), 1.00 (3H, s, H-19), 2.03 (1H, m, H-20), 1.02 (3H, d, $J = 6.7$ Hz, H-21), 5.14 (1H, dd, $J = 15.1, 8.6$ Hz, H-22), 5.02 (1H, dd, $J = 15.1, 8.6$ Hz, H-23), 1.52 (1H, m, H-24), 1.46 (1H, m, H-25), 0.92 (3H, d, $J = 6.4$), 0.81 (3H, dd, $J = 8.3, 6.8$ Hz, H-27), 1.60 (2H, m, H-28), 0.84 (3H, dd, $J = 8.3, 6.8$ Hz, H-29); ^{13}C NMR (CD_3OD , 125 MHz): δ (ppm) 37.4 (CH_2 , C-1), 31.8 (CH_2 , C-2), 72.0 (CH, C-3), 42.4 (CH_2 , C-4), 140.9 (C, C-5), 121.9 (CH, C-6), 31.8 (CH_2 , C-7), 32.1 (CH, C-8), 50.3 (CH, C-9), 36.6 (C, C-10), 21.2 (CH_2 , C-11), 39.9 (CH_2 , C-12), 42.5 (C, C-13), 56.9 (CH, C-14), 24.4 (CH_2 , C-15), 29.9 (CH_2 , C-16), 56.2 (CH, C-17), 12.0 (CH_3 , C-18), 19.5 (CH_3 , C-19), 40.6 (CH, C-20), 21.4 (CH_3 , C-21), 138.5 (CH,

C-22), 129.4 (CH, C-23), 51.4 (CH, C-24), 32.0 (CH, C-25), 18.9 (CH₃, C-26), 19.2 (CH₃, C-27), 26.2 (CH₂, C-28), 12.1 (CH₃, C-29) [41].




Figure 1: Structures of compounds 1 – 7.

The present phytochemical investigation led to the isolation of seven compounds from the stem bark of *F. thonningii*, including four flavonoids (1 – 4), one phenolic acid (5), one sugar (6), and a sterol (7). Compound 6 was isolated from the genus *Ficus* for the first time, while this is the first report of compounds 1 – 2 and 4 – 7 from *F. thonningii*. Therefore, these findings improve the *F. thonningii*'s chemical profile and provide additional data for the chemotaxonomic research of the genus *Ficus*.

Whereas, compound 1, which has only been reported previously from *F. tikoua* [42], in this study it has been isolated in the genus *Ficus* and Moraceae family for the second time, revealing a close phylogenetic relationship between *F. thonningii* and *F. tikoua*. As a result, compound 1 might be used as a chemotaxonomic marker to distinguish the genus *Ficus*. In addition, compound 1 (which its

occurrence is very rare) was also reported from other species of *Desmodium*, such as *Desmodium caudatum* [43] and *Desmodium triquetrum* [44], which demonstrated that the two genera (*Ficus* and *Desmodium*) might possibly be related. Furthermore, Compound **6** is yet to be reported from other species of *Ficus*; as such, it can act together with compound **1** as chemotaxonomic markers for *F. thonningii*.

Isoflavones are the major class of flavonoids reported from *Ficus* species [14,16,42]; therefore, the isolation of compounds **2** – **3** agrees with the previous reports and supports the taxonomic position of *F. thonningii* in the genus. Compounds **2** and **3** were previously isolated from *F. pumila* [45], *F. nervosa* [14], and *F. ovata* [46]. The current findings indicate chemotaxonomic relationships between *F. thonningii* and these species. Compound **5** was reported from the fruits of *F. aurata* and *F. hispida* [32], leaves of *F. trigonata* [47], and stem bark of *F. pandurata* [22], indicating the chemotaxonomic relationship between *F. thonningii* and the other species. Taxifolin (**4**) and stigmasterol (**7**) are often present in plants and so have a less taxonomic importance.

4. CONCLUSION

In conclusion, our study provides new phytochemical information for *F. thonningii* and revealed that *F. thonningii* shares certain flavonoids and phenolic acids with other *Ficus* species such as *F. hispida*, *F. tikoua*, *F. pumila*, *F. tsiangii*, *F. ovata*, *F. aurata* and *F. nymphaefolia* supporting its taxonomic assignment to the genus *Ficus*. The isolated compounds in this work can potentially be chemotaxonomic markers for *F. thonningii*. Our findings have increased the chemical diversity of *F. thonningii* compounds and demonstrated their chemotaxonomic relevance. Besides, the isolation of different classes of secondary metabolites that possess diverse pharmacological potentials from *F. thonningii* explained its multiple ethnomedicinal applications.

REFERENCES

1. Ahmed F, Urooj A. Traditional uses, medicinal properties, and phytopharmacology of *Ficus racemosa*: A review. *Pharm. Biol.* 2010; 48(6): 672-681.
2. Al-Musayeb N, Ebada S, Gad H, Youssef F, Ashour M. Chemotaxonomic diversity of three *Ficus* species: Their discrimination using chemometric analysis and their role in combating oxidative stress. *Phcog Mag.* 2017; 13: S613-S622.
3. Singh D, Singh B, Goel R. Traditional uses, phytochemistry and pharmacology of *Ficus religiosa*: A review. *J. Ethnopharmacol.* 2011; 134: 565-583.
4. Salehi B, Mishra AP, Nigam M, Karazhan N, Shukla I, Kie A, Sawicka B, Aleksandra G. *Ficus* plants: State of the art from a phytochemical, pharmacological, and toxicological perspective. *Phytother. Res.* 2021; 35: 1187-1217.
5. Teixeira SP, Costa MFB, Basso-alves JP, Kjellberg F, Pereira RAS. Morphological diversity and function of the stigma in *Ficus* species (Moraceae). *Acta Oecologica.* 2019; 90: 117-131.
6. Noort S, Gardiner A, Tolley K. New records of *Ficus* (Moraceae) species emphasize the conservation significance of inselbergs in Mozambique. *South African J. Bot.* 2007; 73: 642-649.
7. Karangi M. Revisiting the roots of Gikuyu culture through the sacred Mugumo tree. *J. African Cult. Studies.* 2008; 20: 117-132.
8. Maundu P, Tengnas B, Muema N, Birnie A. Useful trees and shrubs for Kenya; ICRAF-ECA, Nairobi, Kenya, 2005.
9. Danthu P, Soloviev P, Gaye A, Sarr A, Seck M, Thomas I. Vegetative propagation of some West African *Ficus* species by cuttings. *Agroforestry System.* 2002; 55: 57-63.
10. Ahur V, Madubunyi I, Adenkola A and Udem S. The effect of ethyl acetate extract of *Ficus thonningii* (Blume) leaves on erythrocyte osmotic fragility and haematological parameters in acetaminophen-treated rats. *Comp Clin Pathol.* 2012; 21: 409-413.
11. Dangarembizi R, Erlwanger K, Moyo D, Chivandi E. Phytochemistry, pharmacology and ethnomedicinal uses of *Ficus thonningii* (Blume Moraceae): A review. *Afr J Tradit Complement Altern Med.* 2013; 10(2): 203–212.
12. Kokwaro JO. Medicinal plants of East Africa. Second edi., Nairobi: University of Nairobi Press; 1993: 180-185.
13. Njoroge G, Kibunga J. Herbal medicine acceptance, sources and utilization for diarrhoea

- management in a cosmopolitan urban area (Thika, Kenya). *Afr. J. Ecol.* 2007; 45(Suppl. 1): 65-70.
14. Chen L, Cheng M, Peng C, Chen I. Secondary metabolites and antimycobacterial activities from the roots of *Ficus nervosa*. *Chem. Biodivers.* 2010; 7: 1814-1821.
 15. Lee C, Lu C, Kuo Y, Chen J, Sun G. New prenylated flavones from the roots of *Ficus beecheyana*. *J. of the Chinese Chem. Society.* 2004; 51(2): 437-441.
 16. Shao T, Liao H, Li X, Chen G, Song X. A new isoflavone from the fruits of *Ficus auriculata* and its antibacterial activity. *Nat. Prod. Res.* 2022; 36(5): 1191-1196.
 17. Wang Y, Liang H, Zhang Q, Cheng W, Yi S. Phytochemical and chemotaxonomic study on *Ficus tsiangii* Merr. ex Corner. *Biochem. Syst. Ecol.* 2014; 57: 210-215.
 18. Wei S, Lu L, Ji Z, Zhang J, Wu W. Chemical constituents from *Ficus tikoua*. *Chem. Nat. Compd.* 2012; 48(3): 484-485.
 19. Muktar B, Bello I, Sallau M. Isolation, characterization and antimicrobial study of lupeol acetate from the root bark of Fig-Mulberry Sycamore (*Ficus sycomorus* LINN). *J. Appl. Sci. Environ. Manag.* 2018; 22(7): 1129-113.
 20. Poumale HMP, Kengap RT, Claude J. Pentacyclic triterpenes and other constituents from *Ficus cordata* (Moraceae). *Zeitschrift fur Naturforschung B.* 2008; 63(11): 1335-1338.
 21. Ragasa CY, Ng VAS, Shen C. Triterpenes from *Ficus nervosa*. *J. Chem. and Pharm. Res.* 2013; 5(12): 1070-1073.
 22. Ramadan MA, Ahmad AS, Nafady AM, Mansour AI. Chemical composition of the stem bark and leaves of *Ficus pandurata* Hance. *Nat. Prod. Res.* 2009; 23(10): 1218-1230.
 23. Sarg TM, Abbas FA, El-sayed ZI, Mustafa AM. Two new polyphenolic compounds from *Ficus retusa* L. "variegata" and the biological activity of the different plant extracts. *J. Pharm. Phytother.* 2011; 3(7): 89-100.
 24. Singh A, Mukhtar HM, Kaur H, Kaur L. Investigation of antiplasmodial efficacy of lupeol and ursolic acid isolated from *Ficus benjamina* leaves extract. *Nat. Prod. Res.* 2019; 34(17): 2514-2517.
 25. Tameye S, Mbeunkeu A, Fouokeng Y, Tameye N, Tabekoueng G, Wansi J, Sewald N, Ndom J, Azebaze A. Ficusanolide A and ficusanolide B, two new cinnamic acid derivative stereoisomers and other constituents of the stem barks of *Ficus exasperata*. *Phytochem. Lett.* 2021; 43: 150-153.
 26. Jia X, Wu Y, Li J, Lei C, Hou A. Alkaloid constituents of *Ficus hispida* and their antiinflammatory

- activity. *Nat. Products Bioprospect.* 2020; 10: 45-49.
27. Shi Z, Lei C, Yu B, Wang H, Hou A. New alkaloids and α -glucosidase inhibitory flavonoids from *Ficus hispida*. *Chem. Biodivers.* 2016; 13(4): 445-450.
28. Wan C, Chen C, Li M, Yang Y, Chen M, Chen J. Chemical constituents and antifungal activity of *Ficus hirta* Vahl. fruits. *Plants.* 2017; 6(44): 1-9.
29. Dai DT, Tam TN, Thien DD, Sa HN, Thuy TT, Anh HTN, Quan DT. A New furanocoumarin glycoside from the roots of *Ficus hirta*. *Lett. Org. Chem.* 2018; 15(12): 1007-1011.
30. Amponsah I, Fleischer T, Dickson R, Annan K, Thoss V. Evaluation of anti-inflammatory and antioxidant activity of furanocoumarins and sterolin from the stem bark of *Ficus exasperata* Vahl (Moraceae). *J. Sci. Innov. Res.* 2013; 2(5): 880-887.
31. Chunyan C, Bo S, Ping L. Isolation and purification of psoralen and bergapten from *Ficus carica* L leaves by high-speed countercurrent chromatography. *J. Liq. Chromatogr. Relat. Technol.* 2009; 32(1): 136-143.
32. Nurhamidah, Firdaus ML, Suryati and Andriani Y. Antibacterial activity of secondary metabolites isolated from *Ficus aurata* (Miq.) Fruits. *CRBB.* 2021; 3(1): 157-164.
33. Wei S, Luan J, Lu L, Wu W, Ji Z. A new benzofuran glucoside from *Ficus tikoua* Bur. *Int. J. Mol. Sci.* 2011; 12: 4946-4952.
34. Ito C, Sato K, Oka T, Inoue M, Ju-ichi M, Omura M, Furukawa H. Two flavones from *Citrus* species. *Phytochem.* 1989; 28(12): 3562-3564.
35. Sasaki H, Kashiwada Y, Shibata H, Takaishi Y. Phytochemistry prenylated flavonoids from *Desmodium caudatum* and evaluation of their anti-MRSA activity. *Phytochem.* 2012; 82:136-142.
36. Li X, Joshi A, ElSohly H, Khan S, Jacob M, Zhang Z, Khan I, Ferreira D, Walker L, Broedel S, Raulli R, Cihlar R. Fatty acid synthase inhibitors from plants: Isolation, structure elucidation, and SAR studies. *J. Nat. Prod.* 2002; 65(12): 1909-1914.
37. Awouafack MD, Spiteller P, Lamshöft M, Kusari S, Ivanova B, Tane P, Spiteller M. Antimicrobial isopropenyl-dihydrofuranisoflavones from *Crotalaria lachnophora*. *J. Nat. Prod.* 2011; 74(2): 272-278.
38. Usman A, Mohammad R, Abdullahi A, Zakari A, Usman N. Isolation of dihydroquercetin glycoside from the root bark of *Calotropis procera* and antioxidant and cytotoxic screening of the crude extracts. *J. Chem. Soc. Niger.* 2021; 46(1): 0083-0093.

39. Erukainure OL, Choudhary MI, Adhikari A, Islam MS, Hafizur RM, Mesaik AM, Muhammad A, Atolani O, Banerjee P, Preissner R. Anti-diabetic effect of the ethyl acetate fraction of *Clerodendrum volubile*: protocatechuic acid suppresses phagocytic oxidative burst and modulates inflammatory cytokines. *Biomed. Pharmacother.* 2017; 86: 307-315.
40. Valizadeh H, Mahmoodi KF, Alizadeh Z, Bahadori MB. Isolation and structure elucidation of secondary metabolites from *Echinophora platyloba* DC from Iran. *J. Med. Plants.* 13(49): 15-21.
41. Ayele TT, Gurmessa GT, Abdissa Z, Kenasa G, Abdissa N. Oleanane and stigmasterol-type triterpenoid derivatives from the stem bark of *Albizia gummifera* and their antibacterial activities. *J. Chem.* 2022; 1-7.
42. Zhou Q, Lei X, Niu J, Chen Y, Shen X, Zhang N. A new hemiacetal chromone racemate and α -glucosidase inhibitors from *Ficus tikoua* Bur. *Nat. Prod. Res.* 2022; 1-9.
43. Lu G, Ye Y, Lu W, Huang Z, Lan Y, Zeng F, Huang W. Isolation, purification and determination of yukovanol in *Desmodium caudatum* (Thunb.) DC. *Nat Prod Res Dev.* 2014; 26(2): 1450-1453.
44. Xiang W, Li RT, Mao YL, Zhang HJ, Li SH, Song QS, Sun HD. Four new prenylated isoflavonoids in *Tadehagi triquetrum*. *J. Agric. Food Chem.* 2005; 53(2): 267-271.
45. Xiao W, Chen W, Song X, Chen G, Zhang J, Liu L, Han C. Chemical constituents from the stems of *Ficus pumila*. *Chin. Tradit. Pat. Med.* 2015; 37: 1734-1737.
46. Kuete V, Nana F, Ngameni B, Tsafack A, Keumedjio F, Tchaleu B. Antimicrobial activity of the crude extract, fractions and compounds from stem bark of *Ficus ovata* (Moraceae). *Journal of Ethnopharmacol.* 2009; 124: 556-561.
47. Afifi W, Ragab E, Mohammed A, El-Hela A. Bioactivities and phytoconstituents of *Ficus trigonata*. *J. Pharmacogn. Phytochem.* 2014; 3(4): 178-184.